Scientific Basis for Current Hand Hygiene Guidelines and Future Trends


John M. Boyce, MD
J.M. Boyce Consulting, LLC
Middletown, CT

www.jmboyceconsulting.com

1

Presentation Objectives

- Review of evidence used to develop the CDC and WHO Guidelines for Hand Hygiene in Healthcare Settings
- Current challenges in improving hand hygiene compliance
- Future trends in hand hygiene products and strategies for improving hand hygiene practices



Hand Hygiene Intervention


May 1847

- · Students and doctors were required to:
 - clean their hands with a chlorine solution when entering the Obstetrics ward
 - hands were washed with soap & water after examining each patient

Hand Hygiene Intervention

October 1847

- A patient located in the bed where rounds were started each day developed signs of infection
- 11/12 patients who delivered on the ward died
- Soap & water handwashing between patients was felt to be <u>inadequate</u>
- Students and doctors were subsequently required to:
 - clean their hands with chlorine solution before examining each patient




NIH-Sponsored Prospective, Controlled Trial of Handwashing No Handwashing by Nurses Handwashing with Hexachlorophene Soap

Impact of Handwashing on Transmission of *Staphylococcus aureus*

Group	% Acquiring S. aureus	Avg. Hrs Exposure
No Handwashing	92% *	35 Hrs
Hexachlorophene Handwashing	53% *	133 Hrs
* O P = 9 9	05% CI = 2.6 - 45	n = 0 0001

Mortimer EA et al. Am J Dis Child 1962;104:289

Mortimer EA et al. Am J Dis Child 1962;104:289



Factors Associated with Poor Handwashing Compliance


Observed Risk Factors

- Physician (rather than nurse)
- Nursing assistant (not nurse)
- Male sex
- Working during the week
- Wearing gowns/gloves
- Activities with high risk of cross-transmission
- High number of opportunities for hand hygiene per hour of care

Self-Reported Risk Factors

- Handwashing causes irritation and dryness of skin
- Sinks inconveniently located
- Lack of soap and paper towels
- Often too busy/insufficient time
- Patient needs take priority
- Belief that wearing gloves obviates the need for hand hygiene
- Forgetfulness
- Disagreement with guidelines

Adapted from Pittet D Infect Control Hosp Epidemiol 2000;21:381



Hand Hygiene by Handwashing

- 1995 APIC Guideline recommended use of alcoholbased handrubs primarily:
 - If handwashing facilities were insufficient or inadequate
 - Interruption of water supply
- However, hand washing with soap & water continued to be the only form of hand hygiene performed in most hospitals in the United States and many other countries



Larson EL Am J Infect Control 1995;23:251

Attitudes of Healthcare Workers in the United States about Alcohol Hand Rubs

- A few countries in Europe and Scandinavia were using alcohol-based hand rubs (ABHRs) for hand hygiene in the 1980s and 1990s
- However, in the U.S., there was a widespread belief among healthcare workers that: "alcohol will dry out my hands"





Are Recommended Handwashing Policies Practical?

• Time required for soap & water handwashing:

• 62 seconds to get to sink, wash, dry and return

• ICU with 12 nurses

• 40% compliance: 2 to 6.4 hrs/8-hr shift

• 100% compliance: 16 hrs/shift

• Time required for alcoholic hand disinfection:

• 15-second contact time · bedside dispenser

• 40% compliance: 1 to 1.6 hrs/8-hr shift

• 100% compliance: 4 hrs/shift

Voss A & Widmer AF Infect Control Hosp Epidemiol 1997;18:205-8

15

Irritant Contact Dermatitis Due to Frequent Handwashing

- Frequent use of soap & water can lead to skin irritation and damage due to irritant contact dermatitis
 - Painful skin irritation causes healthcare personnel to avoid handwashing
 - May lead to increased colonization of hands by pathogens





Skin Irritation and Dryness: Soap & Water Handwashing vs Alcohol Hand Gel


- In 1998, a 6-week prospective randomized trial with crossover design
 - Funded by GOJO Industries
- 29 nurses on 3 wards participated
- The study compared:
 - a non-medicated, "mild" soap
 - an alcohol hand gel
- Skin irritation/dryness of nurses hands were assessed:
 - self-assessment by participants
 - visual assessment by study nurse
 - measuring electrical capacitance of skin on hands



Boyce JM et al. Infect Control Hosp Epidemiol 2000;21:442







Brief Summary of Evidence Supporting the Use of Alcohol-Based Hand Rubs

- Handwashing compliance among HCWs has remained unacceptably low for decades
- · Advantages of using alcohol-based handrubs
 - can be made more accessible; faster
 - cause less skin irritation and dryness
 - more effective than washing with <u>plain soap</u>/water;
 more effective than washing with <u>antimicrobial soap</u>
 - can promote improved hand hygiene compliance

21

Why Clean Hands Before Touching Patients?

- Cleaning your hands before patient contact protects the patient
- Healthcare workers can contaminate their hands before touching patients by
 - Touching their own skin or mucous membranes
 - Touching contaminated items on the ward
 - Computer keyboards
 - Door knobs
 - Bedside rails or sheets




Courtesy of Prof. Didier Pittet








MRSA on Bed Sheets


VRE on Bedside Rail























Evidence Supporting the Need for Hand Hygiene After Removing Gloves

- HCWs often contaminate their gloves while touching patients or surfaces near patients
- Multiple studies have documented that HCWs' hands may become contaminated even though they wore gloves
- Hand contamination may occur despite glove use because:
 - Gloves may have tiny holes not apparent to HCWs
 - Hands may become contaminated during removal of gloves

Olsen RJ et al. JAMA 1993;270:350
Tenorio AR et al. Clin Infect Dis 2001;32:826
Grundmann H et al. J Infect Dis 2002;185:481
Hayden MK et al. Infect Control Hosp Epidemiol 2008;29:149
Morgan DJ et al. Infect Control Hosp Epidemiol 2010;31:716

HICPAC/SHEA/APIC/IDSA Guideline for Hand Hygiene in Health-Care Settings


- Major recommendations:
 - Alcohol-based hand rub (ABHR) was recommended as the preferred form of hand hygiene if hands are not visibly soiled
 - Indications for when to wash with with soap and water were included
 - Educate healthcare workers (HCWs) regarding the advantages of ABHRs
 - Monitor hand hygiene compliance of HCWs and provide then with feedback on their performance




Boyce JM & Pittet D et al. MMWR 2002;51 (RR-16):1-45

33

2009 World Health Organization (WHO) Guidelines on Hand Hygiene in Health Care





Developed by a team of >100 international experts, led by Prof. Didier Pittet

www.who.int/gpsc/5may/en/

Selected Recommendations from WHO Guideline

- Wash hands with soap and water when visibly dirty or visibly soiled with blood or other body fluids, or after using the toilet
- If exposure to potential spore-forming pathogens is strongly suspected or proven, including outbreaks of C. difficile, hand washing with soap & water is the preferred means
- Use alcohol-based handrub as the preferred means for routine hand antisepsis in all other clinical situations listed below, if hands are not visibly soiled. If alcohol-based handrub is not available, wash hands with soap & water

WHO Guideline for Hand Hygiene in Health Care, 2009

35

Selected Recommendations from WHO Guideline

- Perform hand hygiene:
 - Before and after touching the patient
 - Before handling an invasive device for patient care, regardless of whether or not gloves are used
 - After contact with body fluids or excretions, mucous membranes, non-intact skin, or wound dressings
 - If moving from a contaminated body site to another body site during care of the same patient
 - After contact with inanimate surfaces and objects (including medical equipment) in the immediate vicinity of the patient
 - After removing sterile or non-sterile gloves

WHO Guideline for Hand Hygiene in Health Care, 2009

Selected Recommendations from WHO Guideline

- Before handling medication or preparing food, perform hand hygiene using an alcohol-based handrub or wash hands with either plain or antimicrobial soap & water
- Soap and alcohol-based handrub should not be used concomitantly
- Apply a palmful of alcohol-based handrub and cover all surfaces of the hands. Rub hands until dry. This should take <u>20-30 seconds</u>
- When washing hands with soap & water, wet hands with water and apply the amount of product necessary to cover all surfaces. Rinse hands with water and dry thoroughly with a single-use towel. This should take 40-60 seconds

WHO Guideline for Hand Hygiene in Health Care, 2009


37


Selected Recommendations from WHO Guideline


- Educate HCWs about the type of patient-care activities that can result in hand contamination and about the advantages and disadvantages of various methods to clean their hands
- Monitor HCWs' adherence to recommended hand hygiene practices and provide them with performance feedback
- Encourage partnerships between patients, their families, and HCWs to promote hand hygiene in health care settings

WHO Guideline for Hand Hygiene in Health Care, 2009









SHEA 2014 Compendium: Strategies to Prevent Healthcare-Associated Infections through Hand Hygiene

In 2014, SHEA published a document intended to:

- Highlight practical recommendations in a concise format
- Update recommendations with the most current evidence
- Elucidate topics that warrant clarification or more research
- Assist healthcare facilities in implementing hand hygiene adherence improvement programs
 - including efforts to optimize hand hygiene product use
 - monitor and report back hand hygiene adherence data
 - promote behavior change

Ellingson K et al. Infect Control Hosp Epidemiol 2014;35:937

43

Hand Hygiene in LTCFs

- Deficiencies in hand hygiene practices are among the top 25 deficiencies for which LTCFs are cited by CMS
- From 2000 2009, an average of 9% of nursing homes per year received a deficiency citation (F-Tag 444) for inadequate hand hygiene practices from CMS
 - In Wisconsin, 10% to 36% of nursing homes per year received a deficiency citation for hand hygiene during this time period
- These data suggest that there has been (? and still is) an opportunity to improve hand hygiene practices in LTCFs

Castle N et al. J Appl Gerontol 2014;33:24

Challenges to Improving Hand Hygiene in LTCFs

- Survey of 1143 individuals in 17 nursing facilities in 6 states identified knowledge, attitudes and barriers to hand hygiene
- 29.7% stated that they would not change their hand hygiene practices regardless of guideline recommendations
 - ~20% felt that guidelines were impractical
- ~21% of employees either did not receive training in hand hygiene during the previous year, or were uncertain if they had received training

Ashraf MS et al. Infect Control Hosp Epidemiol 2010;31:758

45

Challenges to Improving Hand Hygiene in LTCFs

- Barriers affecting hand hygiene in LTCFs
 - Staffing shortages of nurses and CNAs
 - Limited financial resources of LTCFs
 - Insufficient hand hygiene product availability
 - Limited in-house infection prevention/control resources
- HCW beliefs that affect hand hygiene practices in LTCFs
 - Too busy to wash hands
 - Senior personnel and colleagues don't wash hands
 - Absence of soap & water, sink, or alcohol-based handrub
 - Didn't wash because I wore gloves
 - Repeated handwashing will damage my skin
 - Just went into the resident's room to talk

Smith A et al. Am J Infect Control 2008;36:492 Ashraf MS et al. Infect Control Hosp Epidemiol 2010;31:758 Herzig CTA et al. J Am Med Dir Assoc 2016;17:85

Challenges to Improving Hand Hygiene in LTCFs: Attitudes of HCWs Regarding Hand Hygiene

Self-reported attitudes regarding hand hygiene among LTCF personnel

Indication for Hand Hygiene	Nurses	CNAs	Others
Wash hands when visibly dirty	60.5%	56%	45%
Wash hands when not visibly dirty	62%	57.4%	42.8%
Wash hands after removing gloves	61.2%	62%	44%

Ashraf MS et al. Infect Control Hosp Epidemiol 2010;31:758

47

Hand Hygiene Promotion Campaign in LTCF

- Comprehensive hand hygiene promotion campaign, conducted in 174-bed LTCF, included:
 - Installation of touch-free alcohol-based handrub dispensers
 - Alcohol wipes placed in common areas and on food trays
 - Educational program for HCWs and for residents
 - Posters promoting hand hygiene located throughout the facility
 - Monitoring of hand hygiene compliance during the intervention
- Results
 - Significant reduction in rate of lower respiratory tract infections
 - Slight reduction in skin and soft tissue infections
 - No change in MRSA, VRE or C. difficile infections
 - Overall hand hygiene compliance during the intervention = 54%

Schweon SJ et al. Am J Infect Control 2013;41:39

Cluster Randomized Controlled Trial of Hand Hygiene Promotion with Pocket-Sized Containers in LTCFs


- Trial involved 6 LTCFs in Hong Kong
- After a 3-month pre-intervention period, LTCFs were randomized and the intervention in 3 LTCFs included:
 - Pocket-sized containers of alcohol-based handrub
 - Reminder materials
 - Education for all HCWs
- Hand hygiene was directly observed and infections recorded
- · Results:
 - Adherence to handrubbing increased from 1.5% to 15.9%
 - Total adherence to hand hygiene increased from 25.8% to 33.3% (p = 0.01)
 - Incidence of serious infections, pneumonia and deaths due to infection decreased significantly in intervention facilities

Yeung WK et al. Infect Control Hosp Epidemiol 2011;32:67


40


Improving Meal-Time Hand Hygiene Among Residents

- 1-month baseline:
 Observations of meal-time hand hygiene by residents
- Barriers to hand hygiene
 - Inaccessible products
 - Difficult to use products
- 6-month interdisciplinary intervention to engage residents & staff
- Meal-time HH improved from 2% to 85% among residents




O'Donnell M et al. Am J Infect Control 2015;43:162






Importance of Product Formulation

- 2 novel alcohol-based hand rub (ABHR) products containing 70% ethanol were compared to
 9 other products with alcohol concentrations of 60% - 90%
- Both new products met efficacy requirements for USA and Europe
- Product efficacy did not correlate with alcohol concentration
- Conclusion: Formulation greatly influences the efficacy of products






Edmonds SL et al. Am J Infect Control 2012;40:521

Factors Influencing Efficacy of ABHRs

- Influence of the volume of ABHR used, product format, and alcohol concentration on
 - Dry-time (time it takes for product to dry on hands)
 - Antimicrobial efficacy
- Volume of ABHR applied to hands is the primary factor affecting dry-time
- Amount of time that the product remains wet on hands greatly influences antimicrobial efficacy
- · Product format did not affect dry-time

Macinga DR et al. BMC Infect Dis 2014;14:511



Compliance with WHO Recommended Hand Hygiene (HH) Technique

- Observational study was conducted in large Swiss hospital with established HH program
- HH was performed in 93.2% of 2,662 opportunities observed
- Compliance with steps 1-6:

Step 1: 92%

Step 2: 83.6%


Step 3: 48.8 %

Step 4: 21.5%

Step 5: 42.5%

Step 6: 19.5%

 Compliance with performing all 6 steps: 8.5%



Tschudin-Sutter S et al. Infect Control Hosp Epidemiol 2015;36:482

Checking Hand Hygiene Technique

- Hand hygiene technique issues include:
 - Is alcohol-based handrub applied to all surfaces of the hands?
 - What is the duration of handrubbing?
- Fluorescent dye can be added to alcohol-based handrub to check for surfaces commonly missed
 - Thumbs
 - Fingertips



Widmer A et al. Infect Control Hosp Epidemiol 2004;25:207

Improving Monitoring of Hand Hygiene Performance

- Direct observations by expert observers
- Self-report by health-care workers
- Direct observations by patients
- Consumption of hygiene products such as towels, soap, and alcohol-based handrub
- Automated monitoring systems

57

Monitoring Hand Hygiene Compliance Using Direct Observation by Trained Observers

- Advantages
 - Determine compliance with all 5 Moments for Hand Hygiene
 - Automated methods cannot monitor Moments 2 and 3
 - Evaluate hand hygiene technique
 - Duration of hand rub
 - Is hand hygiene performed at appropriate times during episode of care
 - Provide immediate feedback to healthcare personnel
 - · Real-time coaching
 - Identify barriers to hand hygiene

Boyce JM Am J Infect Control 2017 (in press)

Monitoring Hand Hygiene Compliance Using Direct Observation by Trained Observers

• Limitations

- Lack of standardized methods precludes comparison of hospitals
- Evaluates < 1% to 2% of all hand hygiene opportunities
- Hawthorne effect may overestimate compliance rates by 300%
- Time-consuming
 - Average number of hand hygiene opportunities (HHOs) that can be observed in 1 hr of observation = 18 (range 3.3 – 41.4)
 - In a hospital with a 70% compliance rate, it is estimated it would require 153 observations per nursing unit per time period (e.g. month) to accurately detect a 10% change in compliance
 - Many hospitals have difficulty providing sufficient auditors

Yin J et al. Infect Control Hosp Epidemiol 2014;35:1163 Boyce JM Am J Infect Control 2017 (in press)

59

Monitoring Product Usage

- · Manual methods of measuring volume of soap and ABHR used
 - Has been useful in establishing trends over time
 - Is used in Europe to compare nursing units and hospitals
 - Requires personnel time to measure, record and analyze results
 - In the U.S., a system for submitting volume measurements on-line for analysis has been associated with increased hand hygiene
 - McGuckin Methods International, Inc.

Limitations

- Personnel time
- Cannot tell who used dispensers (HCW, visitors, patients)
- Does not give information of hand hygiene opportunities or compliance

Pittet D et al. Lancet 2000;356:1307 Hansen S et al. Clin Micribiol Infect 2015;21:1047 McGuckin M et al. Am J Med Qual 2009;24:205

Electronic Monitoring of Product Usage

- Electronic devices can be placed inside product dispensers
- Electronic devices record each time the dispenser is accessed (HH event)
- HH events are time/date stamped
- HH Event data can be downloaded for subsequent analysis

Larson EL et al. Am J Crit Care 2005;14:304
Boyce JM et al. Infect Control Hosp Epidemiol 2009;30:1090
Marra AR et al. Infect Control Hosp Epidemiol 2010;31:796
Sodre da Costa LS Am J Infect Control 2013;41:997
Filho MA et al. Am J Infect Control 2014;42:1188
Arai A et al. Am J Infect Control 2016;44:1481





61

Electronic Monitoring of Product Usage

- Electronic system for monitoring of HH events
 - + estimated number of HH opportunities
 - Dispensers record electronically each time the dispenser is accessed (HH event) and send data to computer server
 - HH opportunities are estimated based patient census, patient-to-nurse ratio, and adjustments
 - HH compliance is estimated by software
 - # of HH events = estimated compliance # of estimated opportunities

Steed C et al. Am J Infect Control 2011;39:19
Diller T et al. Am J Infect Control 2014;42:602
Conway et al. Jt Comm J Qual Pat Saf 2014;40:408
Kwok YL et al. Am J Infect Control 2016;44:1475

Automated Group Monitoring and Feedback Systems

- More complex electronic systems with
 - Counting devices in dispensers, and
 - Sensors detect persons entering/exiting patient rooms
 - Can estimate hand hygiene compliance of groups of personnel
- Dispensers record hand hygiene events
- Room entry = proxy for Moment 1; exit = proxy for Moments 4 & 5
- # of Events / # of room entries & exits = estimated compliance
- Can provide real-time feedback to groups of HCWs
- Shortcoming: cannot tell if persons entering room are HCWs or not

Limper HM et al. Infect Control Hosp Epidemiol 2016 (Epub ahead of print)

63

Automated Badge-Based Monitoring Systems



Automated Badge-Based Monitoring Systems

- At least 20 articles have described evaluations of a variety of badgebased systems
- Systems varied in the technologies used, the settings in which evaluations were conducted, and the duration of each study
- Electronic monitoring systems utilize individual badges worn by HCWs
 - Monitor entry and exit into patient rooms, or proximity to patient
 - Record when HCW wearing badge accesses HH dispenser
 - Estimates compliance with WHO Moments 1, 4 & 5
 - Some systems can provide real-time reminders to HCW
 - Can give HCWs individual, real-time feedback on their performance

6.5

Reviews of Automated Hand Hygiene Monitoring Systems

- A 2014 systematic review by Ward et al. reviewed 42 articles on automated or electronically-assisted monitoring systems
 - < 20% of articles included data on efficiency or accuracy</p>
 - Little compelling data on impact of systems on hand hygiene compliance
 - Further studies are needed of their accuracy, cost, & cost-effectiveness
- Srigley et al. reviewed 7 automated monitoring systems in 2015
 - Most (6) were conducted on a single unit
 - None measured directly observed compliance
 - Study designs varied substantially and were considered to be of poor quality
 - Future studies should include control groups and system-independent measures of hand hygiene to validate system accuracy and predictive value

Boyce JM Infect Control Hosp Epidemiol 2011;32:1016
Ward MA et al. Am J Infect Control 2014;42:472
McGuckin M J Healthc Manag 2015;60:348
Srigley JA et al. J Hosp Infect 2015;89:51
Limper HM et al. Infect Control Hosp Epidemiol 2016:37:1002
Boyce JM Am J Infect Control 2017 (in press)

Automated Hand Hygiene Monitoring Systems

- Further studies of electronic monitoring systems to establish:
 - Accuracy in detecting HH events and estimates of compliance
 - Acceptance by HCW of electronic monitoring systems
 - Ability to accurately reflect compliance with 5 Moments for HH
 - Ability to improve HH compliance rates in a sustained manner
 - Effective ways to use data for feedback and training
 - Their impact on healthcare-associated infection rates
 - Cost-effectiveness

Boyce JM Am J Infect Control 2017 (in press)

67

Direct Observation of All 5 Moments vs Automated Monitoring Moments 1, 4 and 5

- Currently, direct observation by trained observers is the only method for establishing compliance with all 5 Moments for Hand Hygiene
- Inability of current automated HH monitoring systems to estimate compliance with Moments 2 and 3 is a frequently cited limitation
- Question: How does compliance with Moments 1, 4 & 5 compare with compliance with all 5 Moments?
- One study found that compliance with Moments 1 and 4 was 61%, while compliance with all 5 Moments was 62%

Stewardson A et al. J Hosp Infect 2011;77:358

Direct Observation of All 5 Moments vs Automated Monitoring Moments 1, 4 and 5

- Literature review identified a total of 28 studies that reported the distribution of the 5 Moments for Hand Hygiene
- A combined total of 601,988 HH opportunities were reported
- Moments 1, 4 & 5 accounted for 81.3% of all the 5 Moments
- 18 of the studies reported compliance with the 5 Moments
 - For all 5 Moments combined: 66.5%
 - For Moments 1, 4 & 5 combined: 64.9%
- Conclusion: Monitoring Moments 1, 4 & 5 may give reasonable estimate of compliance with all 5 Moments

Boyce JM Am J Infect Control 2017 (in press)

69

Combining Direct Observation with Automated Hand Hygiene Monitoring

- Given its unique capabilities, direct observation should continue to be used for years to come as a qualitative measure of hand hygiene
- As more data on their accuracy, effectiveness and costeffectiveness are generated, automated hand hygiene monitoring systems may become the main quantitative approach to measuring hand hygiene compliance
- A combination of direct observation & automated methods may
 - Provide the best information regarding hand hygiene practices
 - Become a key part of a multimodal strategy for improving hand hygiene

Boyce JM Am J Infect Control 2017 (in press)

Innovation in Hand Hygiene Products

- New formulations of hand hygiene products that are available or under development
- Non-alcohol based hand hygiene product
 - Potassium oleate as primary ingredient
 - Reduces Staphylococus aureus without skin damage
- Hand rubs with activity against Clostridium difficile spores
 - Acidified alcohol-based hand rub
 - Acidified alcohol + peracetic acid
- Hand rubs with improved activity against Norovirus

Asaoka K et al. Am J Infect Control 2016;44:e129 Nerandzic MM et al. PLoS One 2015;10:e132805 Nerandzic MM et al. Open Forum Infect Dis 2015;3:ofv206

7

Innovation: New Dispenser Technology

- Potential improvements in dispenser design and function
 - Notifies HCW of empty dispenser
 - Window shows level of product
 - · Sends electronic signal when empty
 - Auditory or visual cues that draw attention of HCWs
 - Ability to alert HCWs if HH not performed upon room entry
 - Improved electronic wearable dispensers or pocket bottles
 - ? Deliver amount of product based on hand size of HCW




Beyfus TA et al. Am J Infect Control 2016;44:496

Summary

- Hand hygiene guidelines have led to considerable improvements in HH practices in recent years
- Further improvements in compliance are needed
- Greater understanding of factors that influence individual behavior and promote institutional safety climate is needed
- Continued improvement of HH products and delivery systems should facilitate efforts to improve HH compliance
- New approaches to monitoring HH compliance and providing HCWs with feedback will be implemented in coming years

73

Questions?



Average Duration of Handwashing by Healthcare Workers

Year	Mean/median time	
1997	4.7-5.3 seconds	
1994	6.6 seconds	
1974	8-9.3 seconds	
1984	8.6 seconds	
1994	<9 seconds	
1994	9.5 seconds	
1991	<10 seconds	
1990	10 seconds	
1984	11.6 seconds	
1992	12.5 seconds	
1988	15.6-24.4 seconds	
1998	20.6 seconds	
1978	21 seconds	
1989	24 seconds	

Boyce JM & Pittet D MMWR 2002;51 (RR-16):1-45

Preference of Alcohol Hand Rub Dose by Nurses

- Some HCWs believe that 1.1 ml of ABHR is not enough to adequately cover all surfaces of hands; some nures think it is too much
- Prospective observational study included 53 nurses on 8 nursing units in 1 hospital
- Nurses were given a special bottle that recorded each time the bottle was opened.
 Nurses were given a new bottle on 3 shifts
- Bottles were weighed at the end of each of 3 shifts, and average amount (ml) per application was calculated for each nurse
- Hand size was estimated using a validated method




Bottle Provided by GOJO

Martinello R et al. SHEA Spring conference 2017

77


Association Between Hand Size and ABHR Dose Volume per Application by Nurses with Small, Medium and Large Hand Size



• Average dose/application was 1.09 ml

Martinello R et al. SHEA Spring conference 2017





Mody L et al. Infect Control Hosp Epidemiol 2003;24:165

70

Electronic Monitoring of Product Usage

- Electronic system for monitoring of HH events
 - + estimated number of HH opportunities
- Early studies of impact on HH compliance rates:
 - A) In-patient and out-patient units in community hospital (USA)
 - Minimal, temporary increase in HH compliance
 - Number of logistical and implementation problems identified
 - B) Medical ward and surgical ward in large hospital (Australia)
 - Temporary increase in compliance on surgical ward
 - No increase in compliance on medical ward
 - Ward culture and personnel greatly affected results

Conway et al. Jt Comm J Qual Pat Saf 2014;40:408 Kwok YL et al. Am J Infect Control 2016;44:1475

Innovations in Hand Hygiene

- New research will provide additional insights into:
 - Improved methods for educating HCWs regarding HH
 - Novel HH promotional methods based on
 - Behavioral theories
 - · Social marketing techniques
 - Improved institutional interventions to sustain HH promotion
 - Increased use of human factors and ergonomic methods:
 - Influence hospital design to facilitate HH practices

81

Innovation: Electronic Monitoring Systems

- Placing sensors on the patient or on medical equipment to monitor compliance with Moments 2 & 3
 - Example: place sensors intravenous catheters, urinary catheters, wound dressings
- Technology for using video cameras to track position of HCWs in patient rooms, without identifying the patient or HCWs may be further developed and used for monitoring compliance