

This information is from the CounsellingResource.com web site.  
<http://counsellingresource.com/quizzes/adhd-asrs/index.html>

## **Welcome to the Adult ADHD Self-Report Scale (ASRS-v1.1)**

This quick ADHD screening quiz includes just 6 questions which have been found to be the most predictive of DSM-IV-TR symptoms consistent with ADHD. This quiz is adapted from the 18-question symptom checklist developed in conjunction with the World Health Organization and the Workgroup on Adult ADHD. Please read our disclaimer on psychological testing and our psychological testing privacy guarantee.

### ***Completing this Psychological Screening Test***

This screening quiz is a symptoms' checklist for adult ADD. It is not a diagnostic test. To take the questionnaire, please click the radio button next to the selection which best reflects how each statement applies to you. The items below refer to how you have felt and conducted yourself *over the past 6 months*.

### ***Take the Quiz***

Please note: This test will only be scored correctly if you answer each one of the questions. Please check the one response to each item that best describes how you have felt and conducted yourself over the past 6 months.

**1. How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

**2. How often do you have difficulty getting things in order when you have to do a task that requires organization?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

**3. How often do you have problems remembering appointments or obligations?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

**4. When you have a task that requires a lot of thought, how often do you avoid or delay getting started?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

**5. How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

**6. How often do you feel overly active and compelled to do things, like you were driven by a motor?**

- Never
- Rarely
- Sometimes
- Often
- Very Often

## ***About Scoring this Psychological Questionnaire***

When your quiz is scored, one of 2 different information pages will appear to describe the results for scores in your range, along with further details of how your score was computed.

## ***Additional Information***

This is a screening examination for adult ADD. It is not a diagnostic test.

This quiz is adapted from the 18-question symptom checklist called the Adult ADHD Self-Report Scale (ASRS-v1.1), developed in conjunction with the World Health Organization and the Workgroup on Adult ADHD. On the original 18-question test, only 6 of the questions are directly scored to indicate ADHD symptoms, and it is these 6 questions which are included here. The original team of psychiatrists and researchers included:

- Lenard Adler, MD  
Associate Professor of Psychiatry and Neurology  
New York University Medical School
- Ronald C. Kessler, PhD  
Professor, Department of Health Care Policy  
Harvard Medical School
- Thomas Spencer, MD  
Associate Professor of Psychiatry  
Harvard Medical School

### References:

- I. Schweitzer JB, et al. *Med Clin North Am.* 2001;85(3):10-11, 757-777.
- II. Barkley RA. *Attention Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment.* 2nd ed. 1998.
- III. Biederman J, et al. *Am J Psychiatry.* 1993;150:1792-1798.
- IV. American Psychiatric Association: *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision.* Washington, DC, American Psychiatric Association. 2000: 85-93.

The following pages are images of a fax, please excuse the quality.

## Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist Instructions

*The questions on the back page are designed to stimulate dialogue between you and your patients and to help confirm if they may be suffering from the symptoms of attention-deficit/hyperactivity disorder (ADHD).*

Description: The Symptom Checklist is an instrument consisting of the eighteen DSM-IV-TR criteria. Six of the eighteen questions were found to be the most predictive of symptoms consistent with ADHD. These six questions are the basis for the ASRS v1.1 Screener and are also Part A of the Symptom Checklist. Part B of the Symptom Checklist contains the remaining twelve questions.

### Instructions:

#### Symptoms

1. Ask the patient to complete both Part A and Part B of the Symptom Checklist by marking an X in the box that most closely represents the frequency of occurrence of each of the symptoms.
2. Score Part A. If four or more marks appear in the darkly shaded boxes within Part A then the patient has symptoms highly consistent with ADHD in adults and further investigation is warranted.
3. The frequency scores on Part B provide additional cues and can serve as further probes into the patient's symptoms. Pay particular attention to marks appearing in the dark shaded boxes. The frequency-based response is more sensitive with certain questions. No total score or diagnostic likelihood is utilized for the twelve questions. It has been found that the six questions in Part A are the most predictive of the disorder and are best for use as a screening instrument.

#### Impairments

1. Review the entire Symptom Checklist with your patients and evaluate the level of impairment associated with the symptom.
2. Consider work/school, social and family settings.
3. Symptom frequency is often associated with symptom severity, therefore the Symptom Checklist may also aid in the assessment of impairments. If your patients have frequent symptoms, you may want to ask them to describe how these problems have affected the ability to work, take care of things at home, or get along with other people such as their spouse/significant other.

#### History

1. Assess the presence of these symptoms or similar symptoms in childhood. Adults who have ADHD need not have been formally diagnosed in childhood. In evaluating patient's history, look for evidence of early-appearing and long-standing problems with attention or self-control. Some significant symptoms should have been present in childhood, but full symptomology is not necessary.

## The Value of Screening for Adults With ADHD

Research suggests that the symptoms of ADHD can persist into adulthood, having a significant impact on the relationships, careers, and even the personal safety of your patients who may suffer from it.<sup>1-4</sup> Because this disorder is often misunderstood, many people who have it do not receive appropriate treatment and, as a result, may never reach their full potential. Part of the problem is that it can be difficult to diagnose, particularly in adults.

The Adult ADHD Self-Report Scale (ASRS v1.1) Symptom Checklist was developed in conjunction with the World Health Organization (WHO), and the Workgroup on Adult ADHD that included the following team of psychiatrists and researchers:

- **Lenard Adler, MD**  
Associate Professor of Psychiatry and Neurology  
New York University Medical School
- **Ronald C. Kessler, PhD**  
Professor, Department of Health Care Policy  
Harvard Medical School
- **Thomas Spencer, MD**  
Associate Professor of Psychiatry  
Harvard Medical School

As a healthcare professional, you can use the ASRS v1.1 as a tool to help screen for ADHD in adult patients. Insights gained through this screening may suggest the need for a more in-depth clinician interview. The questions in the ASRS v1.1 are consistent with DSM-IV criteria and address the manifestations of ADHD symptoms in adults. Content of the questionnaire also reflects the importance that DSM-IV places on symptoms, impairments, and history for a correct diagnosis.<sup>4</sup>

The checklist takes about 5 minutes to complete and can provide information that is critical to supplement the diagnostic process.

### References:

1. Schweitzer JB, et al. *Med Clin North Am*. 2001;85(3):10-11, 757-777.
2. Barkley RA. *Attention Deficit Hyperactivity Disorder: A Handbook for Diagnosis and Treatment*. 2nd ed. 1998.
3. Biederman J, et al. *Am J Psychiatry*. 1993;150:1792-1798.
4. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. Washington, DC, American Psychiatric Association, 2000: 85-93.

## Adult ADHD Self-Report Scale (ASRS v ) Symptom Checklist

Patient Name	Today's Date					
Please answer the questions below, rating yourself on each of the criteria shown using the scale on the right side of the page. As you answer each question, place an X in the box that best describes how you have felt and conducted yourself over the past 6 months. Please give this completed checklist to your healthcare professional to discuss during today's appointment.			Rarely	Sometimes	Often	Very Often
1. How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?						
2. How often do you have difficulty getting things in order when you have to do a task that requires organization?						
3. How often do you have problems remembering appointments or obligations?						
4. When you have a task that requires a lot of thought, how often do you avoid or delay getting started?						
5. How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?						
6. How often do you feel overly active and compelled to do things, like you were driven by a motor?						
<b>Part A</b>						
7. How often do you make careless mistakes when you have to work on a boring or difficult project?						
8. How often do you have difficulty keeping your attention when you are doing boring or repetitive work?						
9. How often do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?						
10. How often do you misplace or have difficulty finding things at home or at work?						
11. How often are you distracted by activity or noise around you?						
12. How often do you leave your seat in meetings or other situations in which you are expected to remain seated?						
13. How often do you feel restless or fidgety?						
14. How often do you have difficulty unwinding and relaxing when you have time to yourself?						
15. How often do you find yourself talking too much when you are in social situations?						
16. When you're in a conversation, how often do you find yourself finishing the sentences of the people you are talking to, before they can finish them themselves?						
17. How often do you have difficulty waiting your turn in situations when turn taking is required?						
18. How often do you interrupt others when they are busy?						
<b>Part B</b>						