

2005

Wisconsin
**Family Health
Survey**

*Bureau of Health Information and Policy
Division of Public Health
Wisconsin Department of Health and Family Services*

Wisconsin
Family Health Survey

2005

May 2007

*Bureau of Health Information and Policy
Division of Public Health
Wisconsin Department of Health and Family Services*

Foreword

This report on health status, health-related physical activity limitations, chronic conditions and use of health services in Wisconsin is based on information from the 2005 Wisconsin Family Health Survey. This report was compiled in the Wisconsin Department of Health and Family Services, Division of Public Health, Bureau of Health Information and Policy (BHIP).

Ann Spooner, Family Health Survey manager, created the final data set. Stephanie Ward was assisted in the production of this report by a BHIP team consisting of Eleanor Cautley, Ann Spooner and Kim González. Patricia Nametz edited the report. The report was prepared under the supervision of Christine Hill-Sampson, Section Chief, Population Health Information Section, and the overall direction of Patricia Guhleman, Interim Director, Bureau of Health Information and Policy.

Survey sampling and interviewing were conducted by the University of Wisconsin Survey Center. The Division of Health Care Financing, the Division of Disability and Elder Services and the Division of Public Health contributed funding for the Family Health Survey.

The Bureau of Health Information and Policy greatly appreciates the cooperation of the 2,414 survey respondents. We thank them for their contribution to making this information available.

This report is available on the Department of Health and Family Services Web site at the following address: <http://dhfs.wisconsin.gov/stats/familyhealthsurvey.htm>

Comments, suggestions and requests for further information about this report and the Family Health Survey may be addressed to Stephanie Ward at:

Division of Public Health
Bureau of Health Information and Policy
P.O. Box 2659
Madison, WI 53701-2659
608-267-0246
wardsL1@dhfs.state.wi.us

Suggested citation:

Wisconsin Department of Health and Family Services, Division of Public Health, Bureau of Health Information and Policy. **Wisconsin Family Health Survey, 2005** (PPH 5379-05). May 2007.

Contents

Foreword	ii
Introduction	1
Selected Findings	2
Perceived Health Status	3
Figure 1. Perceived Health Status	3
Table 1. Perceived Health Status	4
Figure 2. Fair or Poor Health by Insurance Coverage Over Past Year	6
Figure 3. Fair or Poor Health by Poverty Status	6
Health-Related Limitations Among Adults	7
Table 2. Adults Reported to Have Difficulty Doing at Least One of Five Physical Activities Due to a Health Problem, by Age	7
Figure 4. Adults Reported to Have Difficulty Doing at Least One of Five Physical Activities, by Age and Sex.....	7
Table 3. Adults with Difficulty Performing Physical Activities, by Age	8
Table 4. Health-Related Limitations Among Adults	9
Table 5. Adults with Selected Health-Related Limitations, by Age	10
Chronic Conditions Among Adults	11
Figure 5. Estimated Number of Adults (in Households) Ever Diagnosed with Various Chronic Conditions	11
Table 6. Chronic Conditions Ever Diagnosed Among Adults, by Age	12
Adults with Chronic Conditions and Physical Activity Limitations	14
Table 7. Adults with a Chronic Condition and at Least One of Five Physical Activity Limitations	14
Figure 6. Adults with a Chronic Condition and at Least One of Five Physical Activity Limitations	14
Use of Health Services	15
Doctor Visits:	
Figure 7. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Age and Sex	15
Figure 8. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Insurance Coverage During Year.....	16
Figure 9. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Perceived Health Status.....	16
Table 8. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year.....	17
Figure 10. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Poverty Status	18
Figure 11. Average Number of Visits in Past Year Among Those Who Received Care from a Doctor, Nurse Practitioner, or Physician Assistant, by Age	18

Physical Examinations and Dental Visits:	
Figure 12.	General Physical Examination in Past Year, by Age and Sex 19
Figure 13.	General Physical Examination in Past Year, by Poverty Status 19
Figure 14.	Saw Dentist in Past Year, by Insurance Coverage During Year..... 20
Figure 15.	Saw Dentist in Past Year, by Poverty Status 20
Table 9.	General Physical Examinations and Dental Visits in Past Year 21
Emergency Room Treatment:	
Figure 16.	Treated in Emergency Room in Past Year, by Age and Sex 22
Table 10.	Treated in Emergency Room in Past Year..... 23
Figure 17.	Treated in Emergency Room in Past Year, by Poverty Status 24
Household Population Characteristics	25
Figure 18.	Household Population by Age..... 25
Figure 19.	Household Population by Race and Ethnicity 26
Figure 20.	Metropolitan and Nonmetropolitan Wisconsin..... 27
Figure 21.	Household Population by Poverty Status 28
Table 11.	Household Population by Poverty Status and Age 28
Table 12.	Adult Household Population by Educational Attainment and Age 29
Table 13.	Household Population Aged 18-64 by Employment Status and Sex..... 29
Figure 22.	Household Population Aged 18-64 by Employment Status and Sex..... 30
Table 14.	Characteristics of Wisconsin’s Household Population 31
Technical Notes	32
Survey Design, Definitions of Terms, and Statistical Calculations and Tests..... 32	
Table 15.	Wisconsin Family Health Survey 2005 Sample 33
Table 16.	Wisconsin Family Health Survey Poverty Guidelines, 2004 35
Appendix.	Abbreviated Interview Schedule 38

Introduction

The Wisconsin Family Health Survey (FHS) collects information about health status, health-related physical activity limitations, chronic conditions, use of health care services and health insurance coverage among Wisconsin residents. This survey was started in 1989 and has been conducted annually since that time.

To monitor health status and health care utilization issues, survey questions ask about the current health status, chronic conditions and physical limitations of all household members, as well as visits to a doctor or other health professional, a dentist, and an emergency room in the past year. Background characteristics, such as age, race, poverty status and education, are also obtained for all persons in the household. Numerous survey questions ask about health insurance coverage; results from those questions have been reported in a separate publication, *Wisconsin Health Insurance Coverage, 2005*, available at <http://dhfs.wisconsin.gov/stats/healthinsurance.htm> on the Department of Health and Family Services Web site.

The survey results presented in this report are representative of Wisconsin household residents, who constitute approximately 97 percent of all persons residing in the state. (Non-household residents, including persons living in nursing homes, dormitories, prisons and other institutions, constitute the remaining 3 percent not represented by this survey.) Additional information about the survey design and the results presented here is included in the Technical Notes at the end of this document.

In the Family Health Survey, trained interviewers telephone a random sample of households and ask to speak with the household member most knowledgeable about the health of all household members. This respondent provides information for all people living in the household at the time of the interview. In 2005, the FHS interviewed respondents in 2,414 households; these households included 6,272 people.

The tables in this report show estimated percentages of Wisconsin residents based on survey responses. These estimates should not be treated as precise results because they are derived from a sample. A 95 percent confidence interval (\pm) is printed in a column next to each percentage estimate; this means that 95 percent of similar surveys would obtain an estimate within the confidence interval specified. Tables also include estimated numbers of the Wisconsin household population, based on the weighted sample. Confidence intervals, weighting procedures and statistical tests for significance are described in the Technical Notes at the end of this document, as are variables used in this report, such as insurance coverage, poverty status and metropolitan areas.

Selected Findings


- In 2005, an estimated 12 percent of the Wisconsin household population was reported to be in fair or poor health. Older adults, American Indians, blacks, those who live in the city of Milwaukee, those with less than a high school diploma, those with low incomes and those without health insurance for part or all of the past year were more likely to be reported in fair or poor health.
- An estimated 13 percent of the adult household population was reported to be unable, or found it very difficult, to perform at least one of five physical activities (such as walking three blocks). Those more likely to have health-related physical activity limitations were those aged 65 and over, blacks, those with less education, and those with low incomes.
- Eight percent of household residents were kept from going to school or working at a job or at home because of a health condition. Four percent of people aged 65-74 and 10 percent of those aged 75 and older had trouble eating, dressing, bathing or using the toilet.
- An estimated 44 percent of all adults living in households had ever been diagnosed with one or more of ten chronic health conditions. The most commonly reported of these conditions were high blood pressure and arthritis.
- Almost three-fourths of those who had congestive heart failure (70%) and more than one-half of those who had a stroke (60%) were reported to be unable, or found it very difficult, to perform at least one of five physical activities.
- An estimated 82 percent of household residents received care from a doctor, nurse practitioner or physician assistant at least once during the 12 months preceding the survey interview. Those with no health insurance coverage for the entire year were the least likely to have seen a doctor, nurse practitioner or physician assistant in the past 12 months (52%).
- Sixty-six percent of household residents had a general physical exam in the past year. Blacks were more likely than whites, Hispanics or American Indians to have had a general physical exam in the past year. Those with no health insurance coverage for the entire year were less likely to have had a general physical exam (29%) than those who had coverage part of the year (50%) or those who had insurance coverage all year (69%).
- Eighty percent of household residents (age one and older) were reported to have seen a dentist in the past 12 months. This proportion varied with poverty status: 62 percent of the poor, 68 percent of the near-poor and 86 percent of the non-poor were reported to have seen a dentist during the past year.
- An estimated 16 percent of household residents were treated in an emergency room in the 12 months prior to the survey interview. Adults aged 65 and over, those with less than a high school education, those with low incomes, blacks, those who live in the city of Milwaukee and those with health insurance coverage during part of the past year were more likely to have used an emergency room in the last 12 months.

Perceived Health Status

In 2005, only 12 percent of the Wisconsin household population was reported to be in fair or poor health (an estimated 662,000 people) (Figure 1). In general, reported health status worsens with advancing age. The majority of Wisconsin residents under aged 65 were reported to be in excellent or very good health. While only 4 percent of children were reported in fair or poor health in 2005, 29 percent of older adults (age 65+) were so reported. Those more likely to be reported in fair or poor health were people aged 45 and over, American Indians, blacks, those who live in the city of Milwaukee, those with less than a high school diploma, those with low incomes and those without health insurance for all or part of the past year (Table 1).

The household member with the most knowledge about the health of all household members is selected as the respondent for all people living in the household at the time of the Family Health Survey interview. Reported health status is the respondent's perception of his or her own health and that of other household members. Perceived health status is included in many health surveys because it is strongly related to individual health outcomes. For example, people in fair or poor health are more likely to have been hospitalized during the past year than are those in excellent to good health.

Figure 1. Perceived Health Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

According to the National Center for Health Statistics,¹ 9 percent of the U.S. household population was in fair or poor health in 2005. Thirty-six percent of this population was in excellent health, 31 percent were in very good health and 24 percent were in good health.

¹ National Center for Health Statistics, *Summary Health Statistics for the U.S. Population: National Health Interview Survey, 2005*, http://www.cdc.gov/nchs/about/major/nhis/reports_2005.htm

Perceived Health Status

Table 1. Perceived Health Status, Wisconsin 2005

	Excellent		Very Good		Good	
	Percent	(C.I.±)	Percent	(C.I.±)	Percent	(C.I.±)
Total	30%	(1%)	35%	(1%)	23%	(1%)
Age Groups						
0-17	47	(3)	33	(3)	17	(2)
18-44	33	(2)	37	(2)	21	(2)
45-64	19	(2)	39	(3)	25	(2)
65+	9	(2)	24	(3)	37	(3)
18-64	28	(2)	38	(2)	23	(1)
Sex and Age Groups						
Male	31	(2)	35	(2)	23	(2)
0-17	47	(4)	32	(4)	17	(3)
18-44	35	(3)	36	(3)	22	(3)
45-64	18	(3)	42	(4)	23	(3)
65+	9	(3)	22	(4)	39	(5)
Female	29	(2)	34	(2)	23	(2)
0-17	48	(4)	33	(4)	16	(3)
18-44	31	(3)	38	(3)	20	(3)
45-64	20	(3)	35	(4)	27	(3)
65+	9	(3)	26	(4)	36	(5)
Race/Ethnicity						
White, non-Hispanic	32	(1)	35	(1)	21	(1)
Black, non-Hispanic	15	(4)	30	(5)	35	(5)
Hispanic	28	(7)	25	(7)	31	(7)
American Indian, non-Hispanic	20	(7)	27	(9)	29	(9)
Residence						
City of Milwaukee	24	(4)	26	(3)	31	(4)
Other Metropolitan (excluding city of Milwaukee)	33	(2)	36	(2)	21	(1)
Non-metropolitan	26	(2)	35	(2)	25	(2)
Poverty Status						
Poor	14	(3)	21	(4)	36	(5)
Near-poor	19	(3)	30	(3)	31	(3)
Not poor	35	(2)	38	(2)	19	(1)
Educational Attainment (Ages 18+)						
Less than high school diploma	12	(4)	24	(5)	31	(5)
High school graduate	17	(2)	32	(3)	31	(3)
Education beyond high school	31	(2)	39	(2)	21	(2)
Employment						
Ages 0-17						
Live with employed adult(s)	48	(3)	34	(3)	15	(2)
Live with no employed adult(s)	33	(10)	19	(7)	37	(10)
Ages 18-64						
Employed full-time	29	(2)	40	(2)	23	(2)
Self-employed full-time	34	(6)	37	(6)	19	(5)
Employed part-time	30	(5)	40	(5)	20	(4)
Insurance Coverage over Past Year						
Covered all year	31	(1)	35	(1)	22	(1)
Covered part of year	20	(5)	34	(6)	27	(5)
Uninsured all year	20	(5)	25	(5)	35	(6)

Table 1. Perceived Health Status, Wisconsin 2005 (continued)	Fair or Poor			
	Percent	(C.I.±)	Number	(C.I.±)
Total	12%	(1%)	662,000	(50,000)
Age Groups				
0-17	4	(1)	45,000	(13,000)
18-44	9	(1)	185,000	(31,000)
45-64	17	(2)	226,000	(29,000)
65+	29	(3)	205,000	(27,000)
18-64	12	(1)	411,000	(42,000)
Sex and Age Groups				
Male	11	(1)	300,000	(35,000)
0-17	4	(2)	29,000	(11,000)
18-44	7	(2)	75,000	(20,000)
45-64	16	(3)	106,000	(20,000)
65+	31	(5)	90,000	(18,000)
Female	13	(1)	361,000	(38,000)
0-17	3	(1)	16,000	(8,000)
18-44	11	(2)	109,000	(23,000)
45-64	18	(3)	120,000	(21,000)
65+	29	(4)	116,000	(21,000)
Race/Ethnicity				
White, non-Hispanic	12	(1)	533,000	(45,000)
Black, non-Hispanic	19	(4)	57,000	(15,000)
Hispanic	16	(6)	30,000	(12,000)
American Indian, non-Hispanic	23	(8)	28,000	(11,000)
Residence				
City of Milwaukee	18	(3)	103,000	(19,000)
Other Metropolitan (excluding city of Milwaukee)	10	(1)	338,000	(38,000)
Nonmetropolitan	14	(2)	221,000	(28,000)
Poverty Status				
Poor	29	(4)	127,000	(23,000)
Near-poor	20	(3)	187,000	(29,000)
Not poor	8	(1)	303,000	(34,000)
Educational Attainment (Ages 18+)				
Less than high school diploma	33	(5)	113,000	(21,000)
High school graduate	19	(2)	274,000	(35,000)
Education beyond high school	9	(1)	220,000	(29,000)
Employment				
Ages 0-17				
Live with employed adult(s)	3	(1)	36,000	(12,000)
Live with no employed adult(s)	11	(6)	9,000	(5,000)
Ages 18-64				
Employed full-time	8	(1)	171,000	(28,000)
Self-employed full-time	11	(4)	25,000	(10,000)
Employed part-time	10	(3)	40,000	(14,000)
Insurance Coverage over Past Year				
Covered all year	12	(1)	553,000	(45,000)
Covered part of year	18	(5)	53,000	(15,000)
Uninsured all year	19	(5)	53,000	(17,000)


Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Note: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Perceived Health Status

Health status varies with health insurance coverage and poverty status. Those uninsured all year (19%) or part of the year (18%) were more likely to be reported as being in fair or poor health than those with health insurance coverage all year (12%) (Figure 2).


Figure 2. Fair or Poor Health by Insurance Coverage Over Past Year, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

The poor and near-poor were also more likely to be reported as being in fair or poor health (29% and 20%, respectively) than were the non-poor (8%) (Figure 3).

Figure 3. Fair or Poor Health by Poverty Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Health-Related Limitations Among Adults

Due to health conditions, about one-third (34%) of the Wisconsin adult household population in 2005 (an estimated 1.4 million people) had some level of difficulty doing one or more of five physical activities: walk three city blocks; walk up 10 steps without resting; stand or be on their feet about two hours; stoop, bend, or kneel; and/or reach up over their head. Thirteen percent of adults (an estimated 543,000 people) were unable, or found it very difficult, to do at least one of these five activities (Table 2 and Table 3). Those more likely to have some difficulty doing at least one of the five activities were people aged 65 and over, blacks, those with low incomes, and those with less education (Table 4).

Table 2. Adults Reported to Have Difficulty Doing at Least One of Five Physical Activities Due to a Health Problem, by Age, Wisconsin 2005


Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	5%	(1%)	99,000	(22,000)	13%	(2%)	263,000	(35,000)
45-64	14	(2)	191,000	(27,000)	25	(2)	333,000	(33,000)
65-74	28	(4)	109,000	(20,000)	35	(5)	139,000	(22,000)
75+	47	(5)	143,000	(22,000)	33	(5)	100,000	(19,000)
Total (ages 18+)	13	(1)	543,000	(43,000)	20	(1)	834,000	(53,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Note: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

In all age groups, the percentage of females having difficulty doing at least one of five physical activities was greater than the percentage of males (Figure 4).

Figure 4. Adults Reported to Have Difficulty Doing at Least One of Five Physical Activities, by Age and Sex, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Health-Related Limitations Among Adults

Table 3. Adults with Difficulty Performing Physical Activities, by Age, Wisconsin 2005

Walk About Three City Blocks								
Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	3%	(1%)	55,000	(17,000)	5%	(1%)	95,000	(21,000)
45-64	8	(1)	109,000	(20,000)	10	(2)	131,000	(21,000)
65-74	17	(4)	67,000	(15,000)	21	(4)	81,000	(17,000)
75+	33	(5)	100,000	(19,000)	24	(5)	74,000	(16,000)
Total (ages 18+)	8	(1)	332,000	(35,000)	9	(1)	380,000	(37,000)

Walk Up 10 Steps Without Resting								
Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	1%	(1%)	30,000	(12,000)	3%	(1%)	59,000	(17,000)
45-64	5	(1)	64,000	(16,000)	7	(1)	95,000	(19,000)
65-74	9	(3)	34,000	(11,000)	17	(3)	65,000	(15,000)
75+	22	(5)	68,000	(15,000)	20	(4)	62,000	(15,000)
Total (ages 18+)	5	(1)	196,000	(27,000)	7	(1)	281,000	(32,000)

Stand or Be on Feet About Two Hours								
Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	3%	(1%)	69,000	(19,000)	7%	(1%)	141,000	(26,000)
45-64	10	(2)	131,000	(22,000)	13	(2)	173,000	(25,000)
65-74	19	(4)	74,000	(16,000)	25	(4)	97,000	(19,000)
75+	37	(5)	113,000	(20,000)	24	(5)	73,000	(16,000)
Total (ages 18+)	9	(1)	387,000	(37,000)	12	(1)	485,000	(42,000)

Stoop, Bend, or Kneel								
Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	3%	(1%)	60,000	(17,000)	10%	(1%)	197,000	(31,000)
45-64	9	(2)	124,000	(22,000)	21	(2)	278,000	(31,000)
65-74	15	(3)	57,000	(14,000)	33	(5)	131,000	(21,000)
75+	30	(5)	91,000	(18,000)	37	(5)	112,000	(20,000)
Total (ages 18+)	8	(1)	332,000	(35,000)	17	(1)	718,000	(49,000)

Reach Up Over Head								
Age Group	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	1%	(--)	17,000	(10,000)	3%	(1%)	56,000	(16,000)
45-64	3	(1%)	36,000	(12,000)	6	(1)	79,000	(17,000)
65-74	4	(2)	14,000	(6,000)	13	(3)	52,000	(14,000)
75+	12	(4)	37,000	(11,000)	23	(5)	70,000	(16,000)
Total (ages 18+)	3	(--)	104,000	(20,000)	6	(1)	258,000	(31,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

These estimates were obtained by asking respondents about the degree of difficulty that adult household members had doing the five physical activities because of a physical, mental or emotional health problem. Five separate questions were asked: “By yourself and without any special equipment, how difficult is it for you to: walk a quarter of a mile, or about three city blocks; walk up 10 steps without resting; stand or be on your feet for about two hours; stoop, bend, or kneel; and/ or reach up over your head? Would you say: not difficult, somewhat difficult, very difficult, or are you unable to do this at all?”

Health-Related Limitations Among Adults

Table 4. Health-Related Limitations Among Adults, Wisconsin 2005

	Unable or Very Difficult to Do One or More Physical Activities *		Kept from Working at Job or Home or Going to School		Has Trouble with Eating, Dressing, Bathing, Using Toilet	
	Percent	(C.I.±)	Percent	(C.I.±)	Percent	(C.I.±)
Total (Ages 18+)	13%	(1%)	8%	(1%)	3%	(1%)
Age Groups						
18-44	5	(1)	6	(1)	1	(1)
45-64	14	(2)	9	(2)	3	(1)
65-74	28	(4)	10	(3)	4	(2)
75+	47	(5)	21	(4)	10	(3)
18-64	9	(1)	7	(1)	2	(1)
Sex and Age Groups						
Male (Ages 18+)	10	(1)	7	(1)	2	(1)
18-44	3	(1)	5	(2)	1	(1)
45-64	12	(3)	7	(2)	2	(1)
65-74	24	(6)	8	(4)	3	(2)
75+	43	(8)	21	(7)	10	(6)
Female (Ages 18+)	16	(2)	10	(1)	3	(1)
18-44	6	(2)	7	(2)	2	(1)
45-64	17	(3)	11	(2)	3	(1)
65-74	31	(6)	11	(4)	5	(3)
75+	50	(7)	22	(6)	10	(4)
Race/Ethnicity						
White, non-Hispanic	13	(1)	7	(1)	2	(1)
Black, non-Hispanic	22	(6)	20	(6)	7	(5)
Hispanic	11	(7)	12	(7)	3	(3)
Residence						
City of Milwaukee	20	(4)	17	(4)	7	(3)
Other Metropolitan (excluding city of Milwaukee)	12	(1)	7	(1)	2	(1)
Nonmetropolitan	14	(2)	8	(1)	2	(1)
Poverty Status						
Poor	33	(6)	27	(5)	6	(3)
Near-poor	22	(3)	17	(3)	6	(2)
Not poor	9	(1)	4	(1)	2	(--)
Educational Attainment (Ages 18+)						
Less than high school diploma	28	(5)	20	(4)	7	(3)
High school graduate	16	(2)	9	(2)	3	(1)
Education beyond high school	9	(1)	6	(1)	2	(1)
Employment – Ages 18-64						
Employed full-time	4	(1)	2	(1)	--	(--)
Self-employed full-time	6	(3)	3	(2)	--	(--)
Employed part-time	7	(3)	6	(3)	2	(2)
Insurance Coverage Over Past Year						
Covered all year	14	(1)	8	(1)	3	(1)
Covered part of year	12	(4)	10	(4)	1	(1)
Uninsured all year	8	(4)	8	(4)	2	(2)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

* Walking one block; climbing stairs; bending, lifting, or kneeling; standing two hours; and reaching over head.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Health-Related Limitations Among Adults

At any given time in 2005, an estimated 8 percent of adults in Wisconsin households (343,000 people) were reported to be kept from working at a job, doing work around the house, or going to school because of a health problem (Table 5). This estimate may include both people permanently unable to engage in these activities and those with temporary disabilities.

Three percent of adults in Wisconsin households (110,000 people) had trouble with eating, dressing, bathing, or using the toilet because of a health problem (Table 5).

Table 5. Adults with Selected Health-Related Limitations, by Age, Wisconsin 2005

Age Group	Kept from Working at Job or at Home or Going to School				Had Trouble Eating, Dressing, Bathing, or Using Toilet			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	6%	(1%)	119,000	(24,000)	1%	(1%)	27,000	(12,000)
45-64	9	(2)	120,000	(22,000)	3	(1)	36,000	(12,000)
65-74	10	(3)	38,000	(11,000)	4	(2)	16,000	(8,000)
75+	21	(4)	65,000	(15,000)	10	(3)	30,000	(11,000)
Total (age 18+)	8	(1)	343,000	(37,000)	3	(1)	110,000	(22,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, Pages 36-37.


Respondents were asked about limitations restricting the ability of adult household members to work or attend school: *“Does the physical or mental health of any of the adults in your household keep them from working at a job, doing work around the house, or going to school?”*

Respondents were also asked about limitations with activities of daily living: *“Do any of the adults in your household have trouble eating, dressing, bathing, or using the toilet because of a physical or mental health problem?”* This question is used to measure disability.

Chronic Conditions Among Adults

In 2005, 1.8 million adults living in Wisconsin households (44%) were reported to have one or more of 10 chronic conditions (see Figure 5). Approximately 23 percent of adults (932,000 people) had been diagnosed with high blood pressure, and 19 percent (783,000 people) had been told by a doctor that they have arthritis (Figure 5 and Table 6). Among 18-44-year-olds, 21 percent had at least one of these 10 conditions, compared to 57 percent of those aged 45-64, 83 percent of 65-74-year-olds, and 90 percent of those aged 75 and over. As with all other Family Health Survey results, these estimates do not include persons living in nursing homes and other institutional settings.

Figure 5. Estimated Number of Adults (in Households) Ever Diagnosed with Various Chronic Conditions, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

To obtain data on the prevalence of diagnosed heart disease, cancer, diabetes, stroke, high blood pressure, digestive tract disorders, osteoporosis, heart attack, arthritis, and congestive heart failure, respondents were asked: “*Has anyone in your household ever been told by a doctor or other health professional that they have (each of these 10 chronic diseases or conditions)?*” The population estimates in Figure 5 are underestimates of actual prevalence, since they do not include cases not yet diagnosed.

Chronic Conditions Among Adults

Table 6. Chronic Conditions Ever Diagnosed Among Adults, by Age, Wisconsin 2005

Age Group	Arthritis				Arthritis			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	5%	(2%)	48,000	(16,000)	9%	(2%)	87,000	(20,000)
45-64	19	(3)	126,000	(21,000)	29	(3)	193,000	(26,000)
65-74	37	(7)	62,000	(14,000)	46	(7)	102,000	(19,000)
75+	44	(8)	55,000	(13,000)	61	(7)	110,000	(20,000)
Total (ages 18+)	14	(2)	291,000	(32,000)	23	(2)	492,000	(41,000)

Age Group	Cancer				Cancer			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	1%	(1%)	14,000	(8,000)	3%	(1%)	27,000	(12,000)
45-64	6	(2)	43,000	(13,000)	9	(2)	63,000	(15,000)
65-74	18	(6)	31,000	(11,000)	16	(5)	36,000	(11,000)
75+	30	(8)	37,000	(12,000)	21	(6)	38,000	(12,000)
Total (ages 18+)	6	(1)	125,000	(22,000)	8	(1)	164,000	(25,000)

Age Group	Diabetes				Diabetes			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	2%	(1%)	18,000	(9,000)	3%	(1%)	34,000	(12,000)
45-64	10	(2)	65,000	(16,000)	9	(2)	63,000	(16,000)
65-74	20	(6)	33,000	(10,000)	17	(5)	37,000	(12,000)
75+	22	(7)	27,000	(10,000)	15	(5)	27,000	(10,000)
Total (ages 18+)	7	(1)	144,000	(23,000)	8	(1)	161,000	(25,000)

Age Group	Digestive Tract Disorder				Digestive Tract Disorder			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	7%	(2%)	72,000	(20,000)	10%	(2%)	98,000	(23,000)
45-64	13	(3)	85,000	(18,000)	18	(3)	119,000	(21,000)
65-74	18	(5)	29,000	(10,000)	20	(5)	44,000	(13,000)
75+	20	(6)	25,000	(9,000)	20	(6)	36,000	(12,000)
Total (ages 18+)	11	(1)	211,000	(30,000)	14	(2)	296,000	(34,000)

Age Group	Osteoporosis				Osteoporosis			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	--	(--)	--	(--)	1%	(1%)	12,000	(7,000)
45-64	1%	(1%)	8,000	(5,000)	9	(2)	58,000	(14,000)
65-74	1	(1)	2,000	(2,000)	20	(5)	46,000	(13,000)
75+	5	(4)	6,000	(5,000)	25	(6)	44,000	(12,000)
Total (ages 18+)	1	(--)	16,000	(7,000)	8	(1)	160,000	(24,000)

Age Group	Any of These Five Conditions				Any of These Five Conditions			
	Male		Female		Male		Female	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	13%	(2%)	135,000	(27,000)	19%	(3%)	194,000	(30,000)
45-64	36	(4)	243,000	(30,000)	49	(4)	329,000	(33,000)
65-74	64	(7)	107,000	(19,000)	69	(6)	156,000	(24,000)
75+	71	(8)	88,000	(17,000)	79	(6)	142,000	(23,000)
Total (ages 18+)	29	(2)	573,000	(45,000)	39	(2)	821,000	(51,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Table 6. Chronic Conditions Ever Diagnosed Among Adults, by Age, Wisconsin 2005

Age Group	Congestive Heart Failure							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	--	(--)	3,000	(3,000)	--	(--)	2,000	(2,000)
45-64	1%	(1%)	9,000	(5,000)	1%	(1%)	8,000	(5,000)
65-74	5	(3)	8,000	(5,000)	4	(2)	8,000	(5,000)
75+	11	(5)	14,000	(7,000)	8	(4)	15,000	(7,000)
Total (ages 18+)	2	(1)	34,000	(10,000)	2	(--)	33,000	(10,000)

Age Group	Heart Attack							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	1%	(--)	6,000	(5,000)	--	(--)	2,000	(2,000)
45-64	4	(1%)	25,000	(9,000)	2%	(1%)	10,000	(6,000)
65-74	14	(5)	24,000	(8,000)	5	(3)	12,000	(6,000)
75+	24	(7)	30,000	(10,000)	15	(5)	26,000	(10,000)
Total (ages 18+)	4	(1)	84,000	(17,000)	2	(1)	51,000	(14,000)

Age Group	Heart Disease							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	1%	(1%)	9,000	(6,000)	(--)	(--)	5,000	(4,000)
45-64	9	(2)	59,000	(14,000)	5%	(1%)	30,000	(10,000)
65-74	29	(6)	49,000	(12,000)	15	(5)	34,000	(11,000)
75+	48	(8)	60,000	(14,000)	26	(6)	47,000	(13,000)
Total (ages 18+)	9	(1)	176,000	(24,000)	6	(1)	117,000	(20,000)

Age Group	High Blood Pressure							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	10%	(2%)	102,000	(24,000)	6%	(2%)	65,000	(16,000)
45-64	29	(3)	192,000	(26,000)	30	(3)	200,000	(26,000)
65-74	48	(7)	80,000	(16,000)	55	(6)	123,000	(22,000)
75+	55	(8)	68,000	(15,000)	56	(7)	101,000	(19,000)
Total (ages 18+)	22	(2)	443,000	(40,000)	23	(2)	489,000	(41,000)

Age Group	Stroke							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	--	(--)	3,000	(3,000)	--	(--)	4,000	(4,000)
45-64	3	(1%)	18,000	(9,000)	2%	(1%)	13,000	(8,000)
65-74	7	(4)	12,000	(7,000)	6	(3)	13,000	(8,000)
75+	13	(6)	17,000	(7,000)	12	(4)	21,000	(8,000)
Total (ages 18+)	2	(1)	49,000	(13,000)	2	(1)	51,000	(14,000)

Age Group	Any of These Five Cardiovascular Conditions							
	Male				Female			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
18-44	10%	(2%)	109,000	(24,000)	7%	(2%)	70,000	(17,000)
45-64	33	(4)	220,000	(28,000)	32	(3)	216,000	(27,000)
65-74	62	(7)	104,000	(18,000)	60	(6)	136,000	(23,000)
75+	74	(7)	92,000	(17,000)	67	(7)	121,000	(20,000)
Total (ages 18+)	26	(2)	526,000	(43,000)	26	(2)	543,000	(42,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Adults with Chronic Conditions and Physical Activity Limitations


A majority of those with a chronic condition also had some health-related limitation in performing at least one of five physical activities. (See Table 3 for a list of the five activities.) For example, about 9 of every 10 Wisconsin household adults who had had a stroke (87%) and 8 of every 10 of those with osteoporosis (78%) were limited in performing at least one of the five physical activities (Table 7 and Figure 6).

Table 7. Adults with a Chronic Condition and at Least One of Five Physical Activity Limitations, Wisconsin 2005

Chronic Condition	One or More Physical Activity Limitations							
	Unable or Very Difficult to Do				Somewhat Difficult to Do			
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Number	(C.I.±)
Congestive Heart Failure	70%	(10%)	47,000	(12,000)	22%	(9%)	15,000	(7,000)
Stroke	60	(10)	61,000	(15,000)	27	(9)	27,000	(10,000)
Heart Attack	52	(8)	70,000	(16,000)	30	(8)	41,000	(12,000)
Osteoporosis	45	(7)	78,000	(17,000)	33	(7)	57,000	(14,000)
Heart Disease	47	(5)	138,000	(21,000)	30	(5)	87,000	(17,000)
Arthritis	40	(3)	313,000	(34,000)	37	(3)	290,000	(32,000)
Diabetes	43	(6)	132,000	(22,000)	29	(5)	90,000	(18,000)
Cancer	28	(5)	80,000	(17,000)	32	(5)	92,000	(18,000)
High Blood Pressure	27	(3)	256,000	(30,000)	32	(3)	298,000	(32,000)
Digestive Tract Disorder	27	(4)	139,000	(23,000)	32	(4)	163,000	(26,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Figure 6. Adults with a Chronic Condition and at Least One of Five Physical Activity Limitations, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Use of Health Services

Doctor Visits

Based on 2005 survey results, 82 percent of Wisconsin residents received care from a doctor, nurse practitioner, or physician assistant at least once during the past year (the 12 months preceding the survey interview). Among adults aged 18-64, women were more likely than men to have received care in the past year (Figure 7 and Table 8). This was especially true for women of childbearing age (18-44). Among adults aged 65 and over, however, there was no significant difference by sex for this measure.

Figure 7. Received Care From a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Age and Sex, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

These estimates were derived from the question: “*Approximately how many times have you received medical care from a medical doctor, a nurse practitioner or a physician assistant in the past 12 months?*” This includes a doctor visit for any reason: a checkup, injury, emergency room visit, etc.

People who were uninsured for the entire past year were less likely to have received care than were those with health insurance all or part of the past year (Figure 8).


Figure 8. Received Care From a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Insurance Coverage During Year, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

People reported to be in fair or poor health were the most likely to have seen a doctor in the past year (Figure 9).

Figure 9. Received Care From a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Perceived Health Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Table 8. Received Care from a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, Wisconsin 2005

	Received Medical Care in Past Year*				Received Medical Care 3 or More Times in Past Year*	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)
Total	82%	(1%)	4,433,000	(72,000)	42%	(1%)
Age Groups						
0-17	84	(2)	1,084,000	(59,000)	37	(3)
18-44	77	(2)	1,602,000	(79,000)	38	(2)
45-64	83	(2)	1,121,000	(58,000)	44	(3)
65+	90	(2)	625,000	(45,000)	59	(4)
18-64	80	(1)	2,723,000	(84,000)	40	(2)
Sex and Age Groups						
Male	77	(2)	2,060,000	(77,000)	37	(2)
0-17	85	(3)	559,000	(45,000)	38	(4)
18-44	67	(3)	704,000	(58,000)	28	(3)
45-64	81	(3)	538,000	(44,000)	41	(4)
65+	89	(4)	259,000	(29,000)	59	(5)
Female	87	(1)	2,373,000	(77,000)	47	(2)
0-17	83	(3)	525,000	(43,000)	36	(4)
18-44	88	(2)	898,000	(61,000)	48	(3)
45-64	86	(3)	583,000	(44,000)	47	(4)
65+	91	(3)	366,000	(36,000)	59	(5)
Race/Ethnicity						
White, non-Hispanic	82	(1)	3,786,000	(74,000)	41	(1)
Black, non-Hispanic	89	(3)	261,000	(29,000)	50	(6)
Hispanic	75	(7)	142,000	(26,000)	42	(8)
American Indian non-Hispanic	83	(7)	97,000	(21,000)	48	(10)
Residence						
City of Milwaukee	84	(3)	481,000	(38,000)	50	(4)
Other Metropolitan (excluding city of Milwaukee)	83	(1)	2,717,000	(74,000)	41	(2)
Nonmetropolitan	80	(2)	1,235,000	(53,000)	41	(2)
Poverty Status						
Poor	82	(4)	363,000	(36,000)	55	(5)
Near-poor	83	(3)	772,000	(53,000)	46	(3)
Not poor	82	(1)	3,179,000	(79,000)	39	(2)
Educational Attainment (Ages 18+)						
Less than high school diploma	83	(4)	286,000	(33,000)	47	(5)
High school graduate	81	(2)	1,145,000	(63,000)	43	(3)
Education beyond high school	82	(2)	1,901,000	(76,000)	43	(2)
Employment						
Ages 0-17						
Live with employed adult(s)	84	(2)	1,014,000	(57,000)	36	(3)
Live with no employed adult(s)	86	(8)	71,000	(15,000)	47	(10)
Ages 18-64						
Employed full-time	79	(2)	1,639,000	(75,000)	37	(2)
Self-employed full-time	73	(6)	169,000	(26,000)	34	(6)
Employed part-time	83	(4)	350,000	(39,000)	41	(5)
Insurance During Past Year						
Covered all year	85	(1)	4,059,000	(75,000)	44	(1)
Covered part of year	73	(5)	209,000	(30,000)	33	(6)
Uninsured all year	52	(6)	144,000	(25,000)	22	(5)


Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

* Received medical care from a medical doctor, a nurse practitioner, or a physician assistant.

Note: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

There was no significant difference by poverty level in the percentage who received care in the past year (Figure 10).

Figure 10. Received Care From a Doctor, Nurse Practitioner, or Physician Assistant in Past Year, by Poverty Status, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Forty-two percent of the Wisconsin household population received care from a doctor, nurse practitioner or physician assistant three or more times during the past year. Adults aged 65 and older were more likely to report three or more visits (59%) than those in younger age groups (Table 8).

Older adults also had the highest average number of visits (6.6) among all adults who saw a doctor, nurse practitioner or physician assistant in the past year (Figure 11).

Figure 11. Average Number of Visits in Past Year Among Those Who Received Care From a Doctor, Nurse Practitioner, or Physician Assistant, by Age, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Physical Examinations

Based on 2005 survey results, two-thirds (66%) of Wisconsin residents had a general physical examination during the past year (the 12 months preceding the survey interview) (Table 9). Overall, 72 percent of females and 60 percent of males had a physical exam in the past year. Men aged 18-44 were the least likely to have had a physical exam (42%) (Figure 12).


Survey data also indicate that blacks were more likely than whites and Hispanics to have had a general physical exam in the past year. People with no health insurance coverage for the entire year were less likely to have had a general physical exam (29%) than those who had coverage part of the year (50%) and those who had insurance coverage all year (69%) (Table 9). There was no significant difference in this measure by poverty status (Figure 13).

Figure 12. General Physical Examination in Past Year, by Age and Sex, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Figure 13. General Physical Examination in Past Year, by Poverty Status, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Data were obtained by asking: “Many adults and children get a general physical examination or checkup once in a while. About how long has it been since you visited a doctor for a routine physical exam?”

Dental Visits


Overall, 80 percent of Wisconsin residents one year of age and older were reported to have seen a dentist during the past year (Table 9). This proportion varied widely with health insurance coverage: 84 percent of those with coverage for the entire past year saw a dentist, compared with 51 percent of those uninsured all year (Figure 14). The proportion also varied with poverty status: 62 percent of the poor, 68 percent of the near-poor, and 86 percent of the non-poor were reported to have seen a dentist during the past year (Figure 15).

Figure 14. Saw Dentist in Past Year (Aged One Year and Older), by Insurance Coverage During Year, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Figure 15. Saw Dentist in Past Year (Aged One Year and Older), by Poverty Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Respondents were asked for each household member aged one and older: “How long ago did you last visit the dentist or other dental health care provider?” If respondents inquired, they were told to include any kind of dental health care provider, such as an endodontist, periodontist, or orthodontist.

Table 9. General Physical Examinations and Dental Visits in Past Year, Wisconsin 2005

	General Physical Exam in Past Year*				Dental Visit in Past Year**	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)
Total	66%	(1%)	3,568,000	(78,000)	80%	(1%)
Age Groups						
0-17	76	(2)	977,000	(56,000)	79	(2)
18-44	54	(2)	1,123,000	(68,000)	80	(2)
45-64	68	(2)	911,000	(53,000)	85	(2)
65+	80	(3)	557,000	(42,000)	74	(3)
18-64	60	(2)	2,035,000	(78,000)	82	(1)
Sex and Age Groups						
Male	60	(2)	1,599,000	(71,000)	79	(2)
0-17	76	(3)	505,000	(42,000)	81	(3)
18-44	42	(4)	438,000	(47,000)	76	(3)
45-64	63	(4)	420,000	(39,000)	83	(3)
65+	81	(4)	237,000	(28,000)	72	(5)
Female	72	(2)	1,969,000	(73,000)	82	(1)
0-17	75	(3)	472,000	(41,000)	76	(3)
18-44	67	(3)	685,000	(54,000)	85	(2)
45-64	73	(3)	492,000	(40,000)	86	(2)
65+	79	(4)	319,000	(34,000)	75	(4)
Race/Ethnicity						
White, non-Hispanic	66	(1)	3,019,000	(77,000)	81	(1)
Black, non-Hispanic	79	(4)	232,000	(27,000)	78	(5)
Hispanic	61	(8)	116,000	(23,000)	73	(7)
American Indian, non-Hispanic	75	(8)	88,000	(20,000)	77	(8)
Residence						
City of Milwaukee	74	(4)	426,000	(36,000)	75	(4)
Other Metropolitan (excluding city of Milwaukee)	67	(2)	2,183,000	(73,000)	83	(1)
Nonmetropolitan	62	(2)	959,000	(49,000)	76	(2)
Poverty Status						
Poor	67	(5)	298,000	(33,000)	62	(5)
Near-poor	66	(3)	613,000	(47,000)	68	(3)
Not poor	66	(2)	2,554,000	(27,000)	86	(1)
Educational Attainment (Ages 18+)						
Less than high school diploma	63	(5)	217,000	(29,000)	65	(5)
High school graduate	63	(3)	888,000	(56,000)	76	(2)
Education beyond high school	63	(2)	1,476,000	(69,000)	86	(2)
Employment						
Ages 0-17						
Live with employed adult(s)	76	(2)	919,000	(55,000)	78	(2)
Live with no employed adult(s)	70	(10)	58,000	(14,000)	86	(7)
Ages 18-64						
Employed full-time	58	(2)	1,201,000	(66,000)	83	(2)
Self-employed full-time	50	(7)	115,000	(21,000)	82	(5)
Employed part-time	64	(5)	269,000	(35,000)	86	(4)
Insurance During Past Year						
Covered all year	69	(1)	3,330,000	(78,000)	84	(1)
Covered part of year	50	(6)	145,000	(24,000)	59	(6)
Uninsured all year	29	(6)	81,000	(20,000)	51	(6)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

* Two percent did not answer this survey question.


** Aged one and older.

Note: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Emergency Room Treatment

An estimated 16 percent of Wisconsin household residents (839,000 people) were treated in an emergency room (ER) during the past year (see Table 10). Those with significantly higher rates of ER use included those aged 65 and over, those with less than a high school education, those with low incomes, blacks, those who live in the city of Milwaukee, and those with health insurance coverage for only part of the past year (Figure 16 and Table 10).

Figure 16. Treated in Emergency Room in Past Year, by Age and Sex, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Respondents were asked: “In the past 12 months, who in your household has been treated at a hospital emergency room?” “About how many times in the past 12 months has (name) been treated in an emergency room?”

Table 10. Treated in Emergency Room in Past Year, Wisconsin 2005

	Treated in ER in Past Year				Treated in ER 3+ Times in Past Year	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)
Total	16%	(1%)	839,000	(54,000)	3%	(--)
Age Groups						
0-17	16	(2)	213,000	(27,000)	2	(1)
18-44	14	(2)	297,000	(37,000)	3	(1)
45-64	14	(2)	182,000	(25,000)	2	(1)
65+	21	(3)	147,000	(23,000)	3	(1)
18-64	14	(1)	479,000	(44,000)	3	(1)
Sex and Age Groups						
Male	16	(1)	418,000	(40,000)	3	(1)
0-17	19	(3)	124,000	(21,000)	3	(1)
18-44	14	(2)	144,000	(27,000)	3	(1)
45-64	13	(2)	84,000	(18,000)	2	(1)
65+	23	(4)	66,000	(15,000)	2	(2)
Female	15	(1)	421,000	(39,000)	3	(1)
0-17	14	(3)	89,000	(18,000)	2	(1)
18-44	15	(2)	153,000	(25,000)	3	(1)
45-64	15	(3)	98,000	(18,000)	3	(1)
65+	20	(4)	81,000	(17,000)	4	(2)
Race/Ethnicity						
White, non-Hispanic	15	(1)	702,000	(50,000)	2	(--)
Black, non-Hispanic	21	(4)	62,000	(13,000)	8	(3)
Hispanic	14	(5)	26,000	(10,000)	3	(3)
American Indian, non-Hispanic	18	(7)	21,000	(10,000)	2	(2)
Residence						
City of Milwaukee	21	(3)	119,000	(21,000)	5	(2)
Other Metropolitan (excluding city of Milwaukee)	14	(1)	446,000	(42,000)	2	(1)
Nonmetropolitan	18	(2)	273,000	(30,000)	3	(1)
Poverty Status						
Poor	26	(4)	116,000	(21,000)	7	(3)
Near-poor	21	(3)	192,000	(28,000)	5	(2)
Not poor	13	(1)	502,000	(43,000)	1	(--)
Educational Attainment (Ages 18+)						
Less than high school diploma	23	(5)	78,000	(18,000)	5	(2)
High school graduate	16	(2)	226,000	(30,000)	3	(1)
Education beyond high school	14	(1)	316,000	(35,000)	2	(1)
Employment						
Ages 0-17						
Live with employed adult(s)	16	(2)	198,000	(27,000)	2	(1)
Live with no employed adult(s)	17	(7)	14,000	(6,000)	4	(4)
Ages 18-64						
Employed full-time	12	(1)	251,000	(32,000)	1	(--)
Self-employed full-time	11	(4)	26,000	(10,000)	--	(--)
Employed part-time	11	(3)	48,000	(14,000)	1	(1)
Insurance During Past Year						
Covered all year	16	(1)	744,000	(51,000)	3	(--)
Covered part of year	22	(5)	62,000	(15,000)	3	(2)
Uninsured all year	11	(4)	31,000	(11,000)	2	(2)


Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

As previously noted, emergency room treatment during the past year was more common among the poor and the near-poor than among those who were not poor. More than one in four poor residents and more than one in five near-poor residents were treated in emergency rooms, compared to one in eight who were not poor (Figure 17).

Figure 17. Treated in Emergency Room in Past Year, by Poverty Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Note: See Technical Notes (Table 16, page 35) for guidelines used to determine poverty status.

Blacks were more likely than American Indians, whites or Hispanics to have been treated in an emergency room during the year preceding the survey interview (Table 10). Twenty-one percent of black residents had been to an emergency room during the past year, compared with 18 percent of American Indian residents, 15 percent of white residents and 14 percent of Hispanic residents.


Household Population Characteristics

This section describes characteristics of the Wisconsin household population subgroups for whom health statistics are presented in this report. All of the characteristics described here are estimates from the 2005 Family Health Survey weighted data. The Family Health Survey is considered to be representative of all persons who live in Wisconsin households. Survey results can be used to describe household residents, keeping in mind that survey estimates are going to differ from results of a complete count, such as a census.

According to 2005 Family Health Survey results, approximately 63 percent of the household population is in the age bracket generally considered to be “working age” (ages 18-64) (Figure 18). Another 13 percent are adults aged 65 and older, while 24 percent of the household population are children.

The household population consists of males and females in roughly equal proportions (49% and 51%, respectively) (not shown in figure).

Figure 18. Household Population by Age, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Household Population Characteristics

The vast majority of the Wisconsin household population is non-Hispanic white (85%), according to estimates from the Family Health Survey. Five percent of the population is non-Hispanic black and 3 percent is Hispanic or Latino. Two percent of the population is non-Hispanic American Indian, 1 percent is non-Hispanic Asian, and 1 percent is composed of non-Hispanic members of two or more racial groups (Figure 19).

Among children (ages 0-17), 79 percent are non-Hispanic white, 8 percent are non-Hispanic black and 6 percent are Hispanic or Latino. Two percent of children are non-Hispanic American Indian and 2 percent are non-Hispanic members of two or more racial groups. One percent of children are non-Hispanic Asian.

Figure 19. Household Population by Race and Ethnicity, Wisconsin 2005




Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Race and ethnicity estimates are based on two survey questions. Respondents are first asked: “Are you Hispanic or Latino?” This is followed by: “Which one or more of the following is your race: American Indian, Asian, Black or African American, Pacific Islander, White, or something else?” These questions are then asked for each member of the household.

Based on 2005 Family Health Survey estimates, 11 percent of the state’s household population live in the city of Milwaukee, 61 percent live in the balance of Milwaukee County and the other 24 metropolitan counties, and 29 percent live in the 47 nonmetropolitan counties (Table 14).

Figure 20. Metropolitan and Nonmetropolitan Wisconsin



Source: U.S. Office of Management and Budget and U.S. Bureau of the Census.


Household Population Characteristics

According to 2005 Family Health Survey results, 8 percent of Wisconsin's household population lived in a poor household in 2004 (Figure 21).

Twelve percent of Wisconsin children lived in households considered poor, and another 19 percent lived in households considered near-poor (Table 11).

Poverty status was determined by asking respondents about total household income from all sources in 2004 and the number of people living in the household (see Technical Notes, pages 34-35).

Figure 21. Household Population by Poverty Status, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Table 11. Household Population by Poverty Status and Age, Wisconsin 2005

Age Group*	Poverty Status							
	Poor				Near-Poor		Not Poor	
	Percent	(C.I.±)	Number	(C.I.±)	Percent	(C.I.±)	Percent	(C.I.±)
0-17	12%	(2%)	160,000	(23,000)	19%	(2%)	67%	(2%)
18-44	8	(1)	166,000	(27,000)	17	(2)	74	(2)
45-64	5	(1)	62,000	(16,000)	11	(2)	81	(2)
Total (all ages)	8	(1)	445,000	(40,000)	17	(1)	72	(1)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

* Poverty status could not be estimated for persons aged 65 and older because the household income questions were not answered for 11 percent of this age group.

An estimated 91 percent of all Wisconsin adults living in households (3,738,000 people) have completed high school or more education (Table 12 and Table 14).

The proportion of “working-age” adults (ages 18-64) who have completed high school or more education (93%) is larger than the proportion among adults aged 65 and older (83%).

Table 12. Adult Household Population by Educational Attainment and Age, Wisconsin 2005

Age Groups	Education Completed					
	Less than high school		High school graduate		More than high school	
	Percent	(C.I.±)	Percent	(C.I.±)	Percent	(C.I.±)
18-44	9%	(1%)	30%	(2%)	61%	(2%)
45-64	4	(1)	36	(3)	59	(3)
65+	16	(3)	45	(4)	38	(3)
All Adults (18+)	8	(1)	34	(2)	57	(2)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.
The category “Less than high school” includes all those who did not graduate from high school and do not have a G.E.D. (General Educational Development certificate).

In 2005, an estimated 61 percent of adults ages 18-64 (2,078,000 people) were employed full-time, 7 percent (231,000) were self-employed full-time, and 12 percent (422,000 people) were employed part-time, making a total of 80 percent who were employed. Men and women differ considerably in the proportion employed full or part-time, with men more likely to be employed full-time (Table 13 and Figure 22).


Table 13. Household Population Aged 18-64 by Employment Status and Sex, Wisconsin 2005

	Employment					
	Employed Full-time		Self-employed Full-time		Employed Part-time	
	Percent	(C.I.±)	Percent	(C.I.±)	Percent	(C.I.±)
Males	68%	(2%)	9%	(1%)	7%	(1%)
Females	53	(3)	4	(1)	18	(2)
Total Aged 18-64	61	(2)	7	(1)	12	(1)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Note: C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Figure 22. Household Population Aged 18-64 by Employment Status and Sex, Wisconsin 2005



Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Table 14. Characteristics of Wisconsin's Household Population, 2005

	Percent	(C.I.±)	Number	(C.I.±)
Total	100%		5,398,000	
Age Groups				
0-17	24	(1%)	1,290,000	(62,000)
18-44	38	(1)	2,068,000	(86,000)
45-64	25	(1)	1,344,000	(62,000)
65+	13	(1)	696,000	(47,000)
Sex and Age Groups				
Male				
0-17	25	(2)	661,000	(48,000)
18-44	39	(2)	1,048,000	(69,000)
45-64	25	(2)	668,000	(48,000)
65+	11	(1)	292,000	(31,000)
Female				
0-17	23	(2)	630,000	(46,000)
18-44	37	(2)	1,020,000	(64,000)
45-64	25	(2)	675,000	(47,000)
65+	15	(1)	404,000	(38,000)
Race/Ethnicity				
White, non-Hispanic	85	(1)	4,603,000	(64,000)
Black, non-Hispanic	5	(1)	294,000	(30,000)
Hispanic	3	(1)	188,000	(30,000)
American Indian, non-Hispanic	2	(--)	117,000	(22,000)
Residence				
City of Milwaukee	11	(1)	574,000	(40,000)
Other Metropolitan (excluding city of Milwaukee)	61	(1)	3,279,000	(71,000)
Nonmetropolitan	29	(1)	1,544,000	(55,000)
Poverty Status				
Poor	8	(1)	445,000	(40,000)
Near-poor	17	(1)	933,000	(58,000)
Not poor	72	(1)	3,861,000	(75,000)
Educational Attainment				
Ages 18 and older				
Less than high school diploma	8	(1)	345,000	(37,000)
High school graduate	34	(2)	1,409,000	(68,000)
Education beyond high school	57	(2)	2,329,000	(80,000)
Employment				
Ages 0-17				
Live with employed adult(s)	94	(1)	1,207,000	(61,000)
Live with no employed adult(s)	6	(1)	83,000	(17,000)
Ages 18-64				
Employed full-time	61	(2)	2,078,000	(80,000)
Self-employed full-time	7	(1)	231,000	(30,000)
Employed part-time	12	(1)	422,000	(43,000)

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

Notes: A dash (--) indicates 0.5 percent or less.

C.I. = Confidence Interval (specifies a range within which the true value probably lies). See Technical Notes, pages 36-37.

Technical Notes

Wisconsin Family Health Survey Design

The Wisconsin Family Health Survey (FHS) is a telephone survey of Wisconsin households, designed to provide estimates of health care coverage, various health problems and use of health care services among people across the state.

The Family Health Survey sampling frame consists of all Wisconsin households with a working telephone. In 2005, the sample design for selecting telephone numbers for the survey divided the state into eight sample strata, five of which were defined geographically by grouping all 72 counties into five areas. Telephone area code/prefix combinations from these five strata were randomly sampled at rates proportionate to the population size of each stratum. A sixth sample stratum consisted of telephone prefixes within the city of Milwaukee that had previously been found to include at least 20 percent black respondents. This stratum was also randomly sampled.

The University of Wisconsin Survey Center, University of Wisconsin-Madison, the contracted survey laboratory, drew the samples and conducted all interviews for 2005. Trained interviewers called the sampled telephone numbers and conducted the survey using a computer-assisted telephone survey system (CASES). Each telephone number was called at least 10 times before being designated unanswered. The final overall response rate was 53 percent.

The questions asked in the FHS were designed in the Wisconsin Bureau of Health Information and Policy. Interviews were conducted from February through December of 2005. The final FHS sample consisted of 2,414 household interviews, representing a total of 6,272 Wisconsin household residents. A total of 439 households were interviewed from February through March; 672 from April through June; 754 from July through September; and 549 from October through December. The demographic characteristics of the 2005 sample are displayed in Table 15, which presents the unweighted frequencies. The results in this table are not representative of the Wisconsin population because they have not been weighted to correct for disproportionate sampling rates.

The adult in each household who knows the most about the health of all household members is selected to answer all survey questions during the telephone interview. This respondent answers survey questions for him/herself as well as for all other household members. Since each household member does not speak directly to the interviewer, survey answers are “reported” by the respondent. The reader will see the phrase . . .” *was reported to be . . .*” in this report. In places where this phrase is not used, the reader should keep in mind that all information here is reported by one respondent on behalf of all household members. In 2005, 70 percent of the respondents were women. Abbreviated versions of various survey questions appear with some of the tables in this report and in the Appendix. A copy of all questions asked in 2005 may be obtained from the Bureau of Health Information and Policy.

The data set for analysis of the 2005 Family Health Survey was constructed in the Bureau of Health Information and Policy, using the individual as the basic unit for analysis. Some missing data (i.e., respondent refused to answer or answered “don’t know”) on the age and sex variables were imputed, using interview transcripts and similar cases. About 8 percent of respondents did not answer questions needed to calculate poverty status. Through imputation from other income information, the final proportion of households with missing information on poverty status was reduced to 4 percent (unweighted for households).

Table 15. Wisconsin Family Health Survey 2005 Sample

Total	6,272		
Age Groups		Residence	
0-17	1,636	City of Milwaukee	1,004
18-44	2,035	Other Metropolitan (excluding city of Milwaukee)	3,434
45-64	1,695	Nonmetropolitan	1,834
65+	906		
Sex and Age Groups		Poverty Status	
Male		Poor	594
0-17	829	Near-poor	1,089
18-44	958	Not poor	4,368
45-64	825		
65+	395		
Female		Educational Attainment	
0-17	807	Ages 18 and older:	
18-44	1,077	Less than high school diploma	414
45-64	870	High school diploma	1,616
65+	511	More than high school	2,571
Ethnicity and Race		Employment	
Hispanic or Latino	193	Ages 0-17	
White, not Hispanic/Latino	5,196	Live with no employed adult(s)	118
Black or African American, not Hispanic/Latino	526	Live with employed adult(s)	1,518
American Indian or Alaska Native, not Hispanic/Latino	123	Ages 18-64	
Asian, not Hispanic/Latino	66	Employed full-time	2,223
Other, not Hispanic/Latino	19	Self-employed full-time	261
Two or more races, not Hispanic/Latino	99	Employed part-time	453

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services.

A weight was constructed for each person record in the data set, to adjust for the varying sampling rates, response rates by stratum and number of telephone numbers in each interviewed household. When these weights are applied to the data set, the results are considered to be representative of all Wisconsin household residents in 2005.

One additional component was included to construct the final weight: the total estimated household population in Wisconsin, tabulated for 40 separate subgroups. These subgroups were composed of the combinations of four age groups (0-17, 18-44, 45-64, 65+), by two sex groups, by five geographic regions. The sum total of the 40 subgroups is the estimated household population. Also, the black population within Milwaukee County was adjusted to match the percentage black in the 2000 Census. The population used to weight this data set was 5,397,901, the total estimated household population for Wisconsin on July 1, 2004. This “post-stratification” weight component is applied to each data set record along with the weight described above.

These data set weights were used in computing each percentage and number of people presented in this report. This is the best available method to produce reliable results from the survey data. All references to “weighted” data in this report refer to data that have been adjusted by using these weights so they are representative of the Wisconsin household population.

Definitions of Terms Used in This Report

Age and Sex. These characteristics are reported by the respondent for each household member. Individual years of age are classified into four groups for analysis: ages 0 through 17, 18 through 44, 45 through 64, and 65 and older.

Ethnicity and Race. FHS respondents were first asked if anyone in the household was Hispanic or Latino. Then they were asked to report each household member's race or races. Up to five races could be reported for each person.

In this report, all persons who were reported to be Hispanic or Latino are in the Hispanic/Latino category. All persons not reported as Hispanic/Latino, but for whom two or more races were reported, are in the "two or more races" category. All remaining persons are distributed in the "single-race, not Hispanic/Latino" categories. Some ethnic and racial groups are not included in the tables due to small sample sizes.

Metropolitan and Nonmetropolitan. Twenty-five Wisconsin counties are designated as metropolitan counties by the federal Office of Management and Budget, based on the 2000 U.S. Census standards. These counties are: Brown, Calumet, Chippewa, Columbia, Dane, Douglas, Eau Claire, Fond du Lac, Iowa, Kenosha, Kewaunee, La Crosse, Marathon, Milwaukee, Oconto, Outagamie, Ozaukee, Pierce, Racine, Rock, St. Croix, Sheboygan, Washington, Waukesha, and Winnebago. Counties are designated as metropolitan because they either 1) have a central city of at least 50,000 people or 2) are adjacent and economically linked to a "central city" county. For the tables in this report, results for the city of Milwaukee have been separated from the rest of the metropolitan counties. The "Other Metropolitan" category includes Milwaukee County outside the city plus the remaining 24 metropolitan counties. The other 47 counties are nonmetropolitan.

Poverty Status. The relationship between the number of people in a household and the annual income of that household determines the poverty status. The Family Health Survey asked several questions about total household income during the calendar year prior to the survey (2004), and used current household size to determine whether a household's income was below the federal poverty guideline. A household of four people was considered poor if the total income was below \$19,000. (This is an approximation of the 2004 federal guideline, which was \$18,850.) The "near-poor" category used in this report includes all people in households where the income was greater than the poverty guideline but less than twice the guideline. For a household of four, this was \$38,000 (Table 16).

Educational Attainment. Years of schooling completed are categorized in three groups for this report. Adults who finished 11 grades of school or less are in the first group, "less than high school diploma." Adults who completed 12 years of school or a G.E.D. are in the "high school graduate" group, and adults who attended college or technical school beyond high school are in the "education beyond high school" group.

Working-Age Adults (ages 18 to 64). People in this age range are classified by employment status. Those who were working full-time for an employer at the time of the survey interview are grouped together; some in this group also were self-employed. Among those not working full-time for an employer, those who were self-employed full-time are grouped together, as are those who were working part-time. The remaining adults ages 18-64 include homemakers, the retired, full-time students, persons laid off, the unemployed (either looking or not looking for work), and those disabled persons who are unable to work. These adults were not grouped together, as they are too disparate.

Table 16. Wisconsin Family Health Survey Poverty Guidelines, 2004

Household Size	Poor	Near-Poor
1	\$9,000	\$19,000
2	\$12,000	\$25,000
3	\$16,000	\$31,000
4	\$19,000	\$38,000
5	\$22,000	\$44,000
6	\$25,000	\$50,000

Source: 2005 Family Health Survey, Bureau of Health Information and Policy, Division of Public Health, Wisconsin Department of Health and Family Services. Guidelines derived from *Federal Register*, February 13, 2004, and rounded to nearest \$1,000.

Note: All members of a household were considered “poor” if total household income was less than the poverty guideline shown for a household of that size. Household members were considered “near-poor” if total household income fell between the poor and near-poor guidelines shown for a household of that size.

Children Under Age 18. All children under age 18 are classified by the employment status of the adults in their household. If at least one adult was employed either part-time or full-time, then the child was classified as living with an employed adult. If no adult in the child’s household was employed at the time of the interview, then the child was classified as living with no employed adults.

Health Insurance. As used in this report, “health insurance” includes any kind of private or public coverage for health care costs, including Medicare, Wisconsin Medicaid (or BadgerCare) and other government-funded insurance. The FHS does not obtain detailed information about the extent of services covered by insurance, nor information about costs of premiums, deductibles and co-payments.

Questions about health insurance coverage inquire about specific types of insurance in this sequence: Medicare, employer-sponsored, Medicare supplement or Medigap, private (insurance bought directly from an agent or company), coverage from someone not living in the household, military health care (TRICARE, CHAMPUS, CHAMP-VA, VA), Medicaid (including Title 19, BadgerCare and Healthy Start), and other types of coverage (HIRSP and GAMP are specifically mentioned). For each type of insurance, the respondent is asked whether any household members are currently enrolled and, for each enrolled person, whether that person has been enrolled for less than or more than 12 months.

At the end of this set of questions, the respondent is asked about each person who was not reported to be covered by any type of insurance. This verification question locates another small group of people who otherwise would mistakenly be considered uninsured.

People with Indian Health Service medical care and no other coverage are considered uninsured in this report.

Health Insurance Coverage Over the Past Year. This estimates three groups: the percentage of residents who were covered by any type of insurance over the entire 12 months preceding the telephone interview, the percentage who had coverage during part of the 12 months and had no insurance part of the time, and the percentage who had no health insurance at all during the preceding 12 months.

Because FHS interviews were conducted throughout the year, the “preceding 12-month” period is variable. For example, respondents interviewed in May 2005 were asked to report their health insurance coverage for the 12-month period between May 2004 and May 2005.

A comparison between 2004 and 2005 estimates of the percent without health insurance for all of the past year shows no statistically significant difference (5% each year).

Insured and Uninsured. The “current” estimate of health insurance coverage is the percentage (or number) who had health insurance coverage at the time of the interview. It is a “snapshot” estimate, a cross-section of the Wisconsin household population at one point in time. Any type of public or private insurance coverage at the time of the interview classifies a person as having health insurance. Those with no insurance at the time of the interview are considered uninsured.

There is no statistically significant increase in the estimates of the currently uninsured for 2004 and 2005 (7% each year).

Information About Statistical Calculations and Tests

With the exception of Tables 15 and 16, all information presented in the tables and figures in this report, including the estimates of Wisconsin’s household population characteristics, was produced from the weighted 2005 Family Health Survey.

The tables include estimated percentages, 95 percent confidence intervals, and estimated numbers of people. Results are referred to as “estimated” percentages and numbers because all of the results are derived from a sample survey. The weighted survey data provide reliable estimates of characteristics of Wisconsin’s population. The percentage estimates, as well as the percentage confidence intervals, are rounded to whole percentages to avoid the impression of greater precision than is warranted from a sample survey. The estimated numbers of people, which are estimates of the Wisconsin household population, are rounded to the nearest 1,000 for the same reason.

The Family Health Survey conducts interviews with randomly selected households, a sample of all Wisconsin households. The random sample is used to represent the actual Wisconsin population, but the sample will have some small amount of variation from the actual population. Statistical procedures, such as constructing confidence intervals, are a guide to the amount of precision attributed to the survey results. Starting with this year’s report, a new statistical procedure was used to calculate confidence intervals. The new procedure, based on SAS software, yields results that are very similar to the old procedure based on formulas created in Excel software.

In most tables presented in this report, the 95 percent confidence interval (for both the estimated percents and number of people) is in parentheses. Add the confidence interval value to the estimated percent to find the high boundary of the 95 percent confidence interval, and subtract it from the percent to find the low boundary. For example, on the top line of Table 1 (page 5), 12 percent of Wisconsin household residents were reported to be in fair or poor health at the time of the survey interview. Adding and subtracting the 1 percent value yields a 95 percent confidence interval of 11 to 13 percent. This means that 95 out of 100 random surveys would estimate that 11 to 13 percent of Wisconsin household residents in 2005 were in fair or poor health at a given point in time.

The same procedure applies to the estimated number of people: adding and subtracting 50,000 from 662,000 yields a 95 percent confidence interval of 612,000 to 712,000 persons who were in fair or poor health.

A statistical test was used each time a difference between two estimates is identified in the text. For example, the phrase “those more likely to be uninsured ” means that the difference between the identified groups was tested and found to be a statistically significant difference, not due to random variation. Only those differences that are statistically significant at the 0.05 level are mentioned in this report. A t-test of the differences between percents was used to determine statistical significance.

In some tables the percentage estimates would be expected to sum to 100 percent, but they do not. This is due to two factors: rounding to whole numbers and the omission of “no answer” categories. The “no answer” category includes refusals to answer and answers of “don’t know.” Information about the “no answer” or missing data category is presented in tables when it is a notable percentage.

United States and Wisconsin Comparisons

In this report, some Wisconsin household statistics are compared with those of households across the United States. The U.S. statistical report can be viewed on the Web at:
http://www.cdc.gov/nchs/about/major/nhis/reports_2005.htm

Appendix

Abbreviated Interview Schedule 2005 Family Health Survey

The questions are presented here as if they were asked only of the respondent, but in fact most questions were asked about each person living in the respondent's household. The respondent answered all questions on behalf of the other household members. The complicated skip patterns built into the interview schedule are not shown here (nor are the response categories); skip patterns are based on the answers to prior questions. This is a simplified version presented for ease of understanding.

After the interviewer asks who is the most knowledgeable person in the household (in matters related to the health of other household members), that person is selected to be the respondent and answers questions on behalf of everyone in the household. At the start of the interview, the respondent is asked to list all persons living in the household and to give their first name, their relationship to the respondent, and their age and sex.

FAMILY HEALTH INTERVIEW SCHEDULE

- Overall, how satisfied are you with the health care available to you and your family? Would you say you are satisfied or dissatisfied?
- Are you and your family worried about the cost of health care or not worried about the cost of health care?
- Would you say the health care available to you and your family is better, worse or about the same as the health care you had a year ago?
- How many persons live in your household counting all adults and children and including yourself?
- What is the age, sex and relationship to you of each person in this household?
- I would like you to rate the general health of each person in your household. In general, would you say your health is excellent, very good, good, fair, or poor?

(The following questions are asked about adults aged 18 and older.)

- The next questions ask about difficulties that persons 18 and over in your household may have doing certain activities because of a health problem. By "health problem" we mean any physical, mental or emotional problem or illness (not including pregnancy).
- By yourself and without any special equipment, is it not difficult, somewhat difficult, very difficult, or are you unable to do this:
 - Walk a quarter of a mile, or about 3 city blocks?
 - Walk up 10 steps without resting?
 - Stand or be on your feet for about 2 hours?
 - Stoop, bend or kneel?
 - Reach up over your head?
- Are any of the adults in your household limited in any way because of difficulty remembering or because of difficulties due to periods of memory loss or confusion?
- Are any of the adults in your household limited in any way because of an emotional or mental health condition?
- Does the physical or mental health of any of the adults in your household keep them from working at a job, doing work around the house, or going to school?
- Have you had this trouble for less than one year or more than one year?

-
- Do any of the adults in your household have trouble eating, dressing, bathing, or using the toilet because of a physical or mental health problem?
 - Have you had this trouble for less than one year or more than one year?

(End of adult-only section.)

- Do any of the children in your household have any physical, learning or mental health condition that limits their participation in the usual kinds of activities done by most children their age?
- Is there one particular clinic, health center, doctor's office, or other place that you usually go to if you are sick or need advice about your health?
- Would you say there is more than one place or that there is no place that you usually go to?
- Which one of the following places do you go to most often: a community or neighborhood health center, a doctor's office or clinic, a hospital emergency room, urgent care, a VA hospital or clinic or some other kind of place?
- Approximately how many times have you received medical care from a medical doctor, a nurse practitioner or a physician assistant during the past 12 months?
- Many adults and children get a general physical examination or check-up once in a while. About how long has it been since you visited a doctor for a routine physical exam?
- Who in your household has stayed overnight in a hospital during the past 12 months, that is, since *(date)*? This includes staying overnight in the hospital after a birth or for childbirth.
- Who in your household has been treated at a hospital emergency room during the past 12 months, that is, since *(date)*?
- About how many times in the past 12 months have you been treated in an emergency room?
- Have you been treated in an emergency room for a tooth or dental problem in the last 12 months?
- Sometimes people have problems getting medical care or surgery when they need it. During the last 12 months, was there any time when anyone in your household needed medical care or surgery but did not get it?
- What was the main reason you didn't get the care you needed?
- Now I'd like you to think about dental care. How long ago did you last visit the dentist or other dental health care provider?
- What was the main reason you have not visited the dentist in the past 12 months?
- What was the reason you went to the dentist or other dental health care provider at that time? Was it for cleaning, a check-up or examination, treatment of a condition that the dentist discovered at an earlier visit or something was wrong, bothering or hurting you?
- Do you have any kind of insurance coverage that pays for some or all of your routine dental care, including dental insurance, prepaid plans such as HMOs or government plans such as Medicaid or BadgerCare?
- Now I would like you to think about mental health. In the past 12 months, has anyone in your household received any mental health counseling or advice from their doctor, psychologist, psychiatrist, other mental health professional, social worker or case worker?
- Do you have any insurance that covers mental health counseling or advice?

(The following questions are asked about adults aged 65 or over.)

Appendix

- Next I have a few questions about medicines your doctor might have prescribed not counting things like vitamins, calcium tablets or aspirin that anyone can buy at a store.
- Are you currently using any medicine prescribed by a doctor, not counting things like vitamins or aspirin?
- Did you have to pay for all, most, some, or none of the costs of the prescription drugs you currently use?

(End of prescription drug questions.)

- Now I'm going to ask you about certain health conditions. For each one, please tell me whether you or anyone in your household has ever been told by a doctor or other health professional that they had it.
 - First, has anyone in your household ever been told by a doctor that they had diabetes?
 - Heart disease?
 - Stroke?
 - Cancer?
 - Arthritis?
 - Osteoporosis?
 - Digestive tract disorder?
 - High blood pressure?
 - Heart attack?
 - Congestive heart failure?
- Now I have some questions about insurance coverage. At this time, is anyone in your household enrolled in:
 - Medicare, the health insurance for people 65 and older and people with certain disabilities?
 - Health insurance provided through a current or former employer or union?
 - A Medicare supplement or Medigap insurance plan?
 - Health insurance plan bought directly from an insurance agent or insurance company?
 - Health insurance plan of someone who does not live in this household?
 - Military health care or the Indian Health Service? (This includes services like TRICARE, CHAMPUS, CHAMP-VA, or VA.)
 - There are a number of government programs that pay for health care for low-income people and working families. At this time, is anyone in your household enrolled in Medicaid, Title 19, T-19, Medical Assistance, BadgerCare or Healthy Start?
 - HIRSP, the Health Insurance Risk Sharing Plan, GAMP, the General Assistance Medical Program, or any other insurance?

(The next questions were asked as needed for each type of insurance coverage.)

- Have you been enrolled in this health insurance plan for less than 12 months or for more than 12 months?
- Whose employer or union provides this plan?

- At this time, in addition to the policyholder, who else is covered by this plan?
- Does this health plan cover all, most, some or none of the costs of general check-ups and other preventive services?
- When you are sick or injured, does this health plan cover all, most, some or none of the costs of health care at a doctor's office or health care clinic?
- For overnight hospital stays, does this health plan cover all, most, some or none of the costs?
- Is this plan an HMO, that is, a Health Maintenance Organization?

(The next questions were asked for those who were insured for less than 12 months.)

- Were you uninsured at some time during the past 12 months?
- For how many months were you uninsured?

(The next questions were asked of those who did not have health insurance coverage at this time.)

- According to the information I have so far, you do not have health care coverage at this time. Is that correct?
 - Were you covered by health insurance at any time during the last 12 months?
 - For how many of the past 12 months were you uninsured?
 - What kind of health insurance did you have during the time you were insured?
 - What was the main reason your health insurance coverage stopped?

(The next questions were asked about household members of working-age, 18-64.)

- Last week, did you do any work, either full-time or part-time for pay or profit?
- Do you have a job from which you were temporarily absent last week?
- What was the main reason you did not have a job last week?
- Are you going to school full-time?
- Last week, did you have a second job or business, in addition to your main job or business?
- Let's talk about your main job – the job where you worked the most hours last week. Were you working for an employer, self-employed, or both?
- Was your employer the government, a privately-owned company or business, a non-profit organization, or something else?
- How long have you been working for this employer?
- Were you working on a farm?
- Do you work at a place that has more than 50 employees?
- How many hours per week do you usually work on this job?
- Do you consider your job temporary? Why?
- You said that you have health insurance coverage from a current or former employer or union and that other household members are covered through that policy. Is that insurance through this job?
- Do you pay all, most, some, or none of the costs of premiums for this health insurance?
- You said that you did not have health insurance through your place of employment last week. Is that correct?
- Has the employer or the union offered you health insurance?

Appendix

- Does your employer or union offer health insurance to any other employees?
- Would the health insurance offered by your employer or union cover anyone in your household besides you?

(The following questions were asked about all household members 18-64 who are self-employed.)

- How many hours per week do you usually work at your business?
- How many employees, if any, do you have?
- How long have you had this business?
- Is this business farming?

(The next two questions were asked about all household members aged 18 and older.)

- What is the highest grade or level in school or college you have completed?
- Are you now married, widowed, divorced, separated, or never married?
- Are you Hispanic or Latino?
- What is your Hispanic or Latino origin? Is it Mexican-American, Puerto Rican, Cuban, or something else?
- Which one or more of the following is your race? American Indian, Asian, Black or African American, Pacific Islander, White, or something else?
- In what county is this residence located?
(Asked if residence is in Milwaukee County.)
 - Is this residence in the city of Milwaukee?
- What is your Zip code?
- Do you live on a farm?

(The following question was asked if there were children aged 0-5 or a female aged 17-40 in the household.)

- Are you enrolled in the WIC nutrition program which provides nutritious foods to women, infants and children?
- In the past month, did you or anyone in your household get any food or meals from a food pantry; Meals on Wheels; a free or reduced fee school meal program; a senior center; a meal provided by a church or synagogue, or any other kind of food assistance program?
- In the past month, have you or anyone in your household received or used Food Stamps?
- Are you receiving SSI payments?
- Are you receiving SSDI payments?
- Are you receiving Social Security payments?

(The next series of questions was about annual household income. Respondents were asked three income questions, depending on their household size. Answers to these questions were used to compute poverty status. Because this is a complex section of the interview, only one example is given here, based on a household of four.)

- Thinking of the total income for everyone in your household from all sources, before taxes, in 2004, was that income less than \$19,000, between \$19,000 and \$38,000, or greater than \$38,000?

(If the respondent answers “greater than \$38,000,” the following question is asked.)

- Was your total household income in 2004 less than \$57,000 or greater than \$57,000?

(If the respondent answers “greater than \$57,000,” the following question is asked.)

- Was your total household income in 2004 less than \$75,000 or greater than \$75,000?

(End of household income questions.)

- Approximately how long have you had this phone number?
- Is this number listed in the white pages of the telephone directory?
- Do you have more than one telephone number in your household?
- How many residential telephone numbers do you have, not counting cellular numbers, fax lines, or separate computer lines?
- During the last 12 months, was the household you lived in without telephone service at any time, for any reason?
- During the last 12 months, for how long, in total, was your household without phone service?
- Finally, is there something I haven’t asked about your household’s health or health insurance that you think is important for the Wisconsin Department of Health and Family Services to know?

That’s my last question. Thank you very much for your time and cooperation.

