

## Cervical cancer screening, women 18+

Health Priority A: Access to Primary and Preventive Health Services

Objective A2b: Increase provider screening for chronic diseases and other health risks.

2010 Target: No target established

Indicator: Cervical cancer screening, women 18+

### Percent of Wisconsin Women Aged 18+ Who Had a Pap Smear in the Past Three Years

| Year<br>(N) | Total   |     |
|-------------|---------|-----|
|             | Percent | +/- |
| 2000        | 87%     | 2%  |
| (N)         | 1,195   |     |
| 2001        | 90%     | 1%  |
| (N)         | 1,574   |     |
| 2002        | 88%     | 1%  |
| (N)         | 1,991   |     |
| 2004        | 86%     | 2%  |
| (N)         | 2,098   |     |
| 2006        | 86%     | 2%  |
| (N)         | 2,088   |     |
| 2008        | 83%     | 2%  |
| (N)         | 2,902   |     |
| 2010        | 84%     | 2%  |
| (N)         | 1,927   |     |

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Excludes women who reported having had a hysterectomy.

N        Number in sample. See data documentation.

+/-      Confidence interval. See data documentation.

**Continued on next page**

## Cervical cancer screening, women 18+, continued

### Percent of Wisconsin Women Aged 18+ Who Had a Pap Smear in the Past Three Years, by Race/Ethnicity

| Years<br>(N)               | Total   |     | African<br>American* |     | American<br>Indian* |     | Hispanic |     | White*  |     |
|----------------------------|---------|-----|----------------------|-----|---------------------|-----|----------|-----|---------|-----|
|                            | Percent | +/- | Percent              | +/- | Percent             | +/- | Percent  | +/- | Percent | +/- |
| 2000-2002<br>(N)           | 88%     | 1%  | 92%                  | 3%  | **                  |     | 87%      | 5%  | 88%     | 1%  |
|                            | 4,760   |     | 449                  |     |                     |     | 198      |     | 3,927   |     |
| 2001-2002,<br>2004<br>(N)  | 88%     | 1%  | 90%                  | 6%  | **                  |     | 84%      | 8%  | 88%     | 1%  |
|                            | 5,663   |     | 573                  |     |                     |     | 216      |     | 4,654   |     |
| 2002, 2004,<br>2006<br>(N) | 86%     | 1%  | 90%                  | 4%  | **                  |     | 80%      | 16% | 87%     | 1%  |
|                            | 6,177   |     | 626                  |     |                     |     | 120      |     | 5,212   |     |
| 2004, 2006,<br>2008<br>(N) | 85%     | 1%  | 87%                  | 5%  | 80%                 | 16% | 76%      | 15% | 85%     | 1%  |
|                            | 7,057   |     | 683                  |     | 110                 |     | 129      |     | 5,965   |     |
| 2006, 2008,<br>2010<br>(N) | 84%     | 1%  | 88%                  | 4%  | 74%                 | 17% | 74%      | 17% | 85%     | 1%  |
|                            | 6,799   |     | 665                  |     | 112                 |     | 114      |     | 5,767   |     |

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Excludes women who reported having had a hysterectomy.

- N Number in sample. See data documentation.
- +/- Confidence interval. See data documentation.
- \* Non-Hispanic
- \*\* Fewer than 100 cases