

Mesothelioma deaths

Health Priority D: Environmental and Occupational Health Hazards

Objective D2c: By 2010, reduce occupational mesothelioma illness and death by 30 percent below the 2000 baseline.

2010 Target: Death rate of 0.7 deaths per 100,000 population

Indicator: Mesothelioma deaths

Wisconsin Deaths Due to Mesothelioma, by Sex and Age

Year	Total	Deaths per 100,000 Population	Males	Females	Under Age 65	Ages 65-74	Age 75+
2000	55	1.0	50	5	7	18	30
2001	65	1.2	51	14	17	22	26
2002	65	1.1	53	12	11	21	33
2003	67	1.2	54	13	22	19	26
2004	64	1.1	45	19	17	17	30
2005	59	1.0	44	15	12	16	31
2006	78	1.3	65	13	16	23	39
2007	62	1.0	37	25	12	21	29
2008	64	1.1	49	15	10	26	28
2009	69	1.1	53	16	10	27	32

Source: Resident death certificates, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Rates (deaths per 100,000 population) have been age-adjusted to the 2000 U.S. standard population. (See data documentation.)

Continued on next page

Mesothelioma deaths, continued

Wisconsin Deaths Due to Mesothelioma, by Race/Ethnicity

Year	Total	African American*	American Indian*	Asian*	Hispanic	White*
2000	55	0	0	0	0	55
2001	65	2	0	0	0	63
2002	65	0	1	0	0	64
2003	67	0	0	0	0	67
2004	64	0	0	0	0	64
2005	59	2	0	0	0	57
2006	78	2	1	0	1	74
2007	62	0	0	0	1	61
2008	64	1	0	0	1	62
2009	69	1	0	0	1	67

Source: Resident death certificates, Office of Health Informatics, Division of Public Health, Department of Health Services.

* Non-Hispanic