

**Table 3F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
16th Street	1,710	9.1	6.8	78.4	0.2	5.1	0.4	0.0
Aurora	862	5.9	82.3	4.9	0.5	5.9	0.6	0.0
Barron Cty	306	89.5	4.2	5.2	0.0	0.3	0.7	0.0
Bay Area	215	87.0	0.0	1.9	7.0	0.9	3.3	0.0
Buffalo	106	90.6	0.0	5.7	0.0	0.0	3.8	0.0
Burnett Cty	82	*	*	*	*	*	*	*
Calumet Cty	156	73.7	1.3	16.7	0.0	7.7	0.6	0.0
Chippewa	363	96.4	0.3	1.1	0.3	1.7	0.3	0.0
City Mad	1,891	39.9	21.5	26.7	0.4	9.0	2.5	0.0
Clark Cty	183	76.5	0.0	20.2	1.6	0.0	1.6	0.0
Columbia Cty	245	81.6	0.8	15.1	0.0	0.4	2.0	0.0
Cudahy	744	77.8	2.7	14.8	1.1	2.0	1.6	0.0
Dane Cty	256	53.1	16.0	20.7	0.4	6.6	3.1	0.0
Dodge Cty	363	78.2	1.1	18.7	0.3	0.8	0.8	0.0
Door Cty	105	86.7	0.0	12.4	0.0	1.0	0.0	0.0
Dunn Cty	237	82.7	0.8	4.2	0.4	9.7	2.1	0.0
Eau Claire	618	79.3	1.1	3.4	1.0	12.1	3.1	0.0
Family Health	317	74.1	0.3	24.3	0.3	0.0	0.9	0.0
Family Plan	1,003	74.8	1.0	5.1	0.3	18.0	0.8	0.0
Florence Cty	28	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Fond du Lac	557	75.4	2.0	17.6	0.0	4.3	0.7	0.0
Forest Cty	61	*	*	*	*	*	*	*
GLITC	304	12.5	0.0	4.3	78.0	0.0	5.3	0.0
Grant Cty	283	91.9	0.0	6.4	0.0	1.1	0.7	0.0
Green Cty	174	87.4	0.6	11.5	0.0	0.0	0.6	0.0
Health	983	42.3	23.7	31.2	0.5	0.5	1.7	0.0
Jackson	152	88.2	0.0	5.3	6.6	0.0	0.0	0.0
Jefferson Cty	464	64.2	0.0	33.8	0.2	0.9	0.9	0.0
Juneau	333	89.5	0.0	4.2	0.6	0.6	5.1	0.0
Kenosha Cty	1,141	48.5	17.4	31.6	0.2	1.1	1.3	0.0
Kewaunee Cty	81	*	*	*	*	*	*	*
La Crosse Cty	666	77.3	3.9	3.9	0.3	12.6	2.0	0.0
Marinette	253	94.5	0.0	4.0	0.4	0.0	1.2	0.0
Menominee	150	4.0	0.0	4.7	42.7	0.0	48.7	0.0
Mil. City	2,450	14.9	60.7	18.2	0.4	4.1	1.7	0.0
MLK	997	6.0	84.3	5.4	0.5	1.7	2.0	0.1
Monitowoc	434	75.3	1.8	11.8	0.7	8.3	2.1	0.0
Monroe Cty	311	84.9	1.0	12.2	1.0	1.0	0.0	0.0
NEW Comm	1,590	56.2	4.5	27.3	3.2	8.0	0.8	0.0
NHA	1,160	68.3	11.0	18.2	0.2	1.0	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
NWCSA	319	87.5	1.9	3.8	1.6	0.6	4.7	0.0
Oconto Cty	163	97.5	0.0	1.2	1.2	0.0	0.0	0.0
Oneida	173	9.8	0.0	11.0	74.6	0.6	4.0	0.0
Oneida Cty	187	95.7	0.0	2.1	0.5	1.1	0.5	0.0
Outagamie	721	68.2	3.6	14.4	0.8	12.1	0.8	0.0
Pierce Cty	189	89.9	1.6	5.3	2.6	0.0	0.5	0.0
Polk Cty	253	95.3	0.0	3.2	0.4	0.4	0.8	0.0
Portage Cty	328	77.7	0.6	10.7	0.0	10.7	0.3	0.0
Price Cty	109	95.4	0.0	0.9	0.9	2.8	0.0	0.0
Racine	492	27.2	39.4	30.7	0.4	0.8	1.4	0.0
Rusk Cty	123	95.9	0.0	0.8	0.0	0.0	3.3	0.0
Sauk Cty	368	81.3	1.1	16.3	1.1	0.0	0.3	0.0
Sawyer Cty	115	79.1	0.0	4.3	15.7	0.0	0.9	0.0
Seeds	923	27.8	4.6	62.1	1.5	1.6	2.4	0.0
Shawano	211	84.4	0.9	6.2	4.3	0.9	3.3	0.0
Sheboygan	604	58.8	2.8	20.0	0.3	17.7	0.3	0.0
St. Croix Cty	314	88.5	1.0	4.5	1.3	3.5	1.3	0.0
SWCAP	341	93.0	0.0	6.5	0.3	0.0	0.3	0.0
Taylor Cty	134	88.8	0.0	9.7	0.0	0.7	0.7	0.0
Tremp	215	69.8	0.5	27.9	0.5	0.5	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Vernon Cty	165	94.5	0.0	5.5	0.0	0.0	0.0	0.0
Vilas	82	*	*	*	*	*	*	*
Walworth Cty	542	57.0	1.8	39.1	0.6	0.6	0.9	0.0
Wash/Oz	557	86.9	1.1	9.0	0.7	1.1	1.3	0.0
Washburn Cty	123	90.2	1.6	4.9	0.0	0.8	2.4	0.0
Waukesha Cty	699	67.5	3.9	25.3	0.4	1.4	1.4	0.0
Waupaca Cty	244	89.3	0.0	9.0	0.0	0.4	1.2	0.0
Wee Care	2,342	6.0	85.1	2.4	0.2	4.3	1.9	0.0
West Allis	405	72.6	7.4	13.3	1.0	4.2	1.5	0.0
Winnebago	744	76.2	3.4	12.8	0.1	6.6	0.9	0.0
Wood Cty	415	83.6	0.7	8.2	0.2	5.8	1.4	0.0
<b>Wisconsin</b>	<b>34,934</b>	<b>53.6</b>	<b>19.3</b>	<b>19.1</b>	<b>2.0</b>	<b>4.5</b>	<b>1.7</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,153,475</b>	<b>41.5</b>	<b>22.7</b>	<b>28.7</b>	<b>1.1</b>	<b>3.1</b>	<b>0.9</b>	<b>2.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**

Comparison of Age Distribution <sup>(2)</sup>								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
16th Street	1,710	0.6	8.0	11.4	58.4	19.5	2.0	0.0
Aurora	862	0.7	9.0	13.9	63.2	12.8	0.3	0.0
Barron Cty	306	0.0	3.9	14.1	65.7	15.0	1.3	0.0
Bay Area	215	0.0	4.2	8.8	61.4	23.3	2.3	0.0
Buffalo	106	0.0	2.8	16.0	67.0	13.2	0.9	0.0
Burnett Cty	82	*	*	*	*	*	*	*
Calumet Cty	156	0.0	4.5	13.5	66.7	14.1	1.3	0.0
Chippewa	363	0.0	5.0	9.6	69.7	14.6	1.1	0.0
City Mad	1,891	0.4	6.0	12.2	61.8	18.6	0.9	0.0
Clark Cty	183	0.0	3.8	18.6	62.8	14.8	0.0	0.0
Columbia Cty	245	0.0	5.7	13.5	60.4	18.0	2.4	0.0
Cudahy	744	0.1	3.2	11.3	67.2	17.5	0.7	0.0
Dane Cty	256	0.4	4.3	11.3	64.8	17.6	1.6	0.0
Dodge Cty	363	0.3	5.8	12.4	63.6	16.5	1.1	0.3
Door Cty	105	0.0	4.8	8.6	61.9	24.8	0.0	0.0
Dunn Cty	237	0.0	6.8	16.9	60.8	13.9	1.7	0.0
Eau Claire	618	0.2	4.5	14.4	65.5	15.0	0.3	0.0
Family Health	317	0.0	5.4	14.5	61.8	16.7	1.3	0.3
Family Plan	1,003	0.2	5.0	14.0	63.8	16.2	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**

Comparison of Age Distribution <sup>(2)</sup>								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Florence Cty	28	*	*	*	*	*	*	*
Fond du Lac	557	0.0	5.7	14.0	62.7	16.2	1.4	0.0
Forest Cty	61	*	*	*	*	*	*	*
GLITC	304	1.0	7.6	19.7	55.6	15.5	0.3	0.3
Grant Cty	283	0.0	3.9	13.8	65.7	15.2	1.4	0.0
Green Cty	174	0.6	4.0	14.9	64.9	14.9	0.6	0.0
Health	983	0.2	7.4	15.6	61.2	15.0	0.6	0.0
Jackson	152	0.0	4.6	17.1	61.8	15.8	0.7	0.0
Jefferson Cty	464	0.2	5.0	12.1	61.6	19.0	2.2	0.0
Juneau	333	0.3	10.5	13.2	62.2	12.6	1.2	0.0
Kenosha Cty	1,141	0.4	7.7	14.9	59.6	16.0	1.3	0.1
Kewaunee Cty	81	*	*	*	*	*	*	*
La Crosse Cty	666	0.2	5.0	13.4	62.9	17.4	1.2	0.0
Marinette	253	0.0	3.6	12.6	71.5	11.5	0.8	0.0
Menominee	150	0.7	14.0	13.3	55.3	14.7	2.0	0.0
Mil. City	2,450	0.4	9.1	14.7	57.3	17.2	1.1	0.1
MLK	997	1.4	11.8	13.6	58.4	13.9	0.8	0.0
Monitowoc	434	0.2	6.5	15.9	60.6	15.4	1.4	0.0
Monroe Cty	311	0.3	6.1	14.8	65.3	12.9	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**

Comparison of Age Distribution <sup>(2)</sup>								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
NEW Comm	1,590	0.4	6.5	13.0	60.0	19.1	1.0	0.1
NHA	1,160	0.3	7.4	15.8	62.3	13.8	0.4	0.0
NWCSA	319	0.0	6.3	11.6	68.3	12.5	1.3	0.0
Oconto Cty	163	0.6	3.7	19.6	62.0	13.5	0.6	0.0
Oneida	173	0.0	6.4	15.0	64.7	11.0	2.3	0.6
Oneida Cty	187	0.0	2.1	13.4	65.8	16.6	2.1	0.0
Outagamie	721	0.3	5.7	13.6	64.9	14.1	1.4	0.0
Pierce Cty	189	0.5	2.6	12.7	65.1	16.4	2.6	0.0
Polk Cty	253	0.0	7.5	14.2	61.3	15.0	2.0	0.0
Portage Cty	328	0.0	6.4	17.1	61.0	13.7	1.8	0.0
Price Cty	109	0.0	0.9	8.3	69.7	19.3	1.8	0.0
Racine	492	0.4	9.8	15.2	59.1	14.0	1.4	0.0
Rusk Cty	123	0.0	1.6	14.6	62.6	18.7	1.6	0.8
Sauk Cty	368	0.0	5.7	14.7	61.4	17.1	1.1	0.0
Sawyer Cty	115	0.0	5.2	14.8	67.0	13.0	0.0	0.0
Seeds	923	0.4	8.9	15.0	57.1	17.2	1.4	0.0
Shawano	211	0.0	3.8	11.4	63.5	19.0	1.9	0.5
Sheboygan	604	0.2	6.5	13.6	62.3	15.9	1.3	0.3
St. Croix Cty	314	0.3	5.4	11.8	63.4	17.8	1.3	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
SWCAP	341	0.0	2.9	12.6	66.6	15.8	2.1	0.0
Taylor Cty	134	0.7	4.5	9.7	65.7	19.4	0.0	0.0
Tremp	215	0.0	2.3	15.8	66.5	15.3	0.0	0.0
Vernon Cty	165	0.0	1.8	11.5	66.7	18.8	1.2	0.0
Vilas	82	*	*	*	*	*	*	*
Walworth Cty	542	0.4	4.6	14.6	62.0	17.7	0.6	0.2
Wash/Oz	557	0.0	3.6	11.3	69.8	14.7	0.4	0.2
Washburn Cty	123	0.0	4.9	8.9	71.5	14.6	0.0	0.0
Waukesha Cty	699	0.3	4.0	12.0	63.2	19.0	1.4	0.0
Waupaca Cty	244	0.0	5.3	14.8	63.1	14.8	2.0	0.0
Wee Care	2,342	0.7	9.5	17.8	55.4	15.0	1.5	0.0
West Allis	405	0.2	3.5	9.4	64.7	20.2	2.0	0.0
Winnebago	744	0.0	5.1	13.0	67.3	13.6	0.9	0.0
Wood Cty	415	0.0	5.5	11.8	67.0	15.2	0.5	0.0
<b>Wisconsin</b>	<b>34,934</b>	<b>0.3</b>	<b>6.6</b>	<b>13.8</b>	<b>61.8</b>	<b>16.2</b>	<b>1.2</b>	<b>0.1</b>
<b>Nation (Prior Year)</b>	<b>1,153,475</b>	<b>0.4</b>	<b>6.5</b>	<b>12.6</b>	<b>60.0</b>	<b>19.0</b>	<b>1.4</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
16th Street	1,679	5.1 (2)	50.6 (55)	1,494	30.3 (55)	34.9 (1)	198	29.8 (11)	1,494	27.2 (48)	1,521	6.0 (56)	1,521	6.1 (42)
Aurora	858	7.3 (7)	55.2 (65)	36	*	*	140	72.1 (19)	722	57.2 (63)	737	2.4 (10)	737	5.7 (34)
Barron Cty	299	8.4 (12)	48.8 (51)	258	21.7 (19)	51.6 (48)	38	*	261	23.0 (41)	270	4.8 (44)	270	4.8 (28)
Bay Area	212	6.1 (5)	50.9 (56)	171	22.2 (22)	44.4 (18)	25	*	183	15.8 (15)	191	5.2 (47)	191	2.1 (8)
Buffalo	105	8.6 (16)	48.6 (50)	95	*	*	18	*	92	*	93	*	93	*
Burnett Cty	82	*	*	56	*	*	15	*	71	*	75	*	75	*
Calumet Cty	153	8.5 (14)	40.5 (9)	132	28.0 (50)	43.2 (11)	24	*	134	14.9 (12)	147	6.1 (58)	147	6.8 (46)
Chippewa	360	9.2 (23)	45.6 (40)	328	27.7 (48)	41.2 (6)	57	*	322	20.8 (31)	332	3.6 (21)	332	6.3 (44)
City Mad	1,870	10.7 (43)	42.7 (17)	1,182	27.7 (48)	43.0 (10)	358	28.2 (10)	1,305	20.9 (32)	1,729	4.5 (36)	1,729	4.6 (24)
Clark Cty	183	10.4 (37)	44.3 (28)	168	29.8 (54)	43.5 (13)	31	*	168	39.9 (60)	175	2.3 (9)	175	2.9 (12)
Columbia Cty	244	10.7 (43)	40.2 (5)	190	22.1 (20)	53.2 (54)	32	*	204	13.7 (9)	221	4.5 (36)	221	10.0 (59)
Cudahy	742	9.0 (20)	45.3 (37)	639	20.5 (14)	49.6 (41)	152	26.3 (6)	645	22.5 (39)	674	5.2 (47)	674	6.2 (43)
Dane Cty	252	8.7 (17)	41.3 (11)	107	22.4 (24)	52.3 (50)	46	*	136	21.3 (34)	173	3.5 (20)	173	9.8 (58)
Dodge Cty	354	11.9 (56)	42.1 (14)	298	19.1 (8)	43.6 (14)	65	*	316	25.3 (44)	331	3.9 (27)	331	5.7 (34)
Door Cty	105	12.4 (58)	31.4 (1)	97	*	*	23	*	95	*	98	*	98	*
Dunn Cty	236	9.7 (29)	49.2 (52)	107	16.8 (2)	45.8 (24)	40	*	178	11.2 (3)	209	5.7 (55)	209	7.2 (47)
Eau Claire	618	10.8 (47)	42.4 (15)	30	*	*	107	25.2 (5)	523	26.0 (45)	556	2.9 (14)	556	3.2 (15)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Local Agency	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)
Family Health	302	8.9 (19)	48.0 (47)	266	23.3 (33)	50.0 (44)	48	* *	240	19.2(26)	270	5.2 (47)	270	4.4(21)
Family Plan	989	10.5 (40)	43.7 (23)	878	25.7 (42)	40.8 (5)	169	16.6 (1)	886	11.2 (3)	920	3.4 (19)	920	4.7(26)
Florence Cty	28	* *	* *	22	* *	* *	6	* *	20	* *	23	* *	23	* *
Fond du Lac	548	11.3 (50)	43.4 (20)	488	29.3 (53)	39.1 (3)	78	* *	469	27.1(47)	497	4.0 (29)	497	5.0(30)
Forest Cty	61	* *	* *	56	* *	* *	6	* *	50	* *	58	* *	58	* *
GLITC	303	7.6 (9)	51.2 (59)	210	24.8 (39)	47.1 (35)	50	* *	210	41.0(61)	277	3.2 (16)	277	3.2(15)
Grant Cty	279	9.7 (29)	48.0 (47)	230	18.7 (4)	46.5 (30)	56	* *	247	19.4(28)	259	4.2 (32)	259	5.8(37)
Green Cty	174	9.8 (31)	44.3 (28)	153	22.9 (30)	46.4 (29)	28	* *	151	27.2(48)	152	3.3 (18)	152	1.3(4)
Health	974	9.0 (20)	50.9 (56)	855	25.1 (41)	46.9 (34)	153	30.1 (12)	842	22.4(38)	863	5.3 (50)	863	8.5(56)
Jackson	151	9.9 (33)	44.4 (30)	140	22.1 (20)	45.7 (23)	22	* *	136	22.1(36)	135	0.7 (1)	135	0.7(3)
Jefferson Cty	454	9.9 (33)	45.4 (39)	250	22.4 (24)	42.8 (8)	81	* *	393	13.2 (8)	415	3.9 (27)	415	5.5(33)
Juneau	333	8.4 (12)	51.7 (60)	142	24.6 (37)	51.4 (46)	48	* *	266	16.5(17)	298	3.7 (24)	298	7.7(50)
Kenosha Cty	1,111	9.4 (25)	47.3 (45)	1,021	23.8 (36)	45.2 (21)	209	35.4 (14)	977	32.1(56)	985	2.7 (12)	985	5.7(34)
Kewaunee Cty	81	* *	* *	70	* *	* *	10	* *	67	* *	69	* *	69	* *
La Crosse Cty	665	12.3 (57)	41.7 (12)	334	23.7 (35)	46.7 (33)	80	* *	542	20.1(30)	600	2.7 (12)	600	0.5(2)
Marinette	246	17.5 (66)	37.0 (2)	211	19.0 (6)	48.8 (39)	36	* *	167	22.2(37)	226	6.2 (59)	226	4.4(21)
Menominee	149	7.4 (8)	47.7 (46)	96	* *	* *	15	* *	129	21.7(35)	134	8.2 (63)	134	8.2(55)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mil. City	2,426	9.5 (27)	49.7 (53)	1,890	26.8 (45)	44.3 (17)	433	46.0 (16)	2,024	35.9 (58)	2,206	4.0 (29)	2,206	7.7 (50)
MLK	985	8.3 (10)	53.8 (63)	506	26.9 (46)	48.6 (38)	184	52.2 (18)	853	45.1 (62)	876	3.2 (16)	876	10.2 (60)
Monitowoc	428	13.8 (63)	40.7 (10)	132	20.5 (14)	51.5 (47)	50	* *	352	24.1 (43)	394	3.6 (21)	394	5.8 (37)
Monroe Cty	305	9.8 (31)	44.9 (32)	0	* *	* *	64	* *	278	28.4 (54)	288	4.5 (36)	288	2.4 (9)
NEW Comm	1,568	11.3 (50)	40.4 (8)	969	28.7 (52)	45.2 (21)	242	38.4 (15)	1,418	28.1 (52)	1,448	4.7 (42)	1,448	5.9 (40)
NHA	1,131	10.8 (47)	48.5 (49)	322	22.7 (28)	37.6 (2)	201	31.3 (13)	983	27.3 (50)	1,026	2.0 (7)	1,026	2.9 (12)
NWCSA	316	10.8 (47)	46.2 (41)	252	19.0 (6)	54.0 (55)	48	* *	262	32.8 (57)	267	7.1 (61)	267	7.9 (53)
Oconto Cty	159	15.7 (65)	45.3 (37)	149	18.8 (5)	52.3 (50)	20	* *	123	17.1 (20)	148	5.4 (52)	148	8.1 (54)
Oneida	173	5.8 (3)	57.8 (66)	144	22.9 (30)	46.5 (30)	30	* *	151	23.2 (42)	159	7.5 (62)	159	4.4 (21)
Oneida Cty	186	11.8 (55)	43.5 (21)	29	* *	* *	30	* *	149	10.7 (2)	162	4.3 (33)	162	1.9 (7)
Outagamie	710	10.4 (37)	43.7 (23)	576	22.6 (26)	46.2 (27)	110	27.3 (9)	615	26.8 (46)	636	4.6 (40)	636	5.3 (32)
Pierce Cty	188	9.6 (28)	42.6 (16)	128	19.5 (9)	46.1 (25)	22	* *	154	17.5 (21)	169	2.4 (10)	169	4.7 (26)
Polk Cty	249	13.3 (61)	47.0 (44)	155	22.6 (26)	43.2 (11)	36	* *	213	18.3 (23)	223	4.5 (36)	223	10.3 (61)
Portage Cty	326	9.2 (23)	45.1 (36)	266	22.9 (30)	45.1 (20)	46	* *	278	15.5 (13)	291	2.1 (8)	291	2.7 (11)
Price Cty	109	6.4 (6)	44.0 (25)	101	14.9 (1)	60.4 (56)	17	* *	100	12.0 (5)	101	1.0 (2)	101	5.0 (30)
Racine	485	10.1 (36)	53.8 (63)	424	26.4 (44)	42.9 (9)	75	* *	423	30.0 (55)	391	3.8 (25)	391	3.6 (17)
Rusk Cty	123	11.4 (52)	43.1 (19)	88	* *	* *	17	* *	108	13.0 (7)	117	4.3 (33)	117	0.0 (1)

- (1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sauk Cty	359	10.6 (42)	52.6 (61)	303	19.5 (9)	49.5 (40)	77	* *	324	16.7 (19)	326	5.5 (54)	326	5.8 (37)
Sawyer Cty	115	8.7 (17)	37.4 (3)	104	20.2 (12)	51.9 (49)	13	* *	91	* *	96	* *	96	* *
Seeds	916	9.4 (25)	46.8 (43)	824	24.8 (39)	44.2 (16)	145	22.1 (2)	841	19.9 (29)	850	6.0 (56)	850	7.2 (47)
Shawano	209	4.3 (1)	45.0 (34)	160	20.0 (11)	52.5 (52)	31	* *	121	9.1 (1)	193	3.6 (21)	193	6.7 (45)
Sheboygan	604	9.1 (22)	44.0 (25)	454	24.7 (38)	42.1 (7)	124	23.4 (4)	535	16.6 (18)	546	4.6 (40)	546	3.1 (14)
St. Croix Cty	305	11.5 (53)	43.6 (22)	274	18.6 (3)	49.6 (41)	38	* *	243	12.3 (6)	263	5.3 (50)	263	10.6 (62)
SWCAP	336	6.0 (4)	50.9 (56)	254	23.6 (34)	52.8 (53)	52	* *	249	20.9 (32)	301	4.7 (42)	301	9.0 (57)
Taylor Cty	133	12.8 (60)	44.4 (30)	121	28.1 (51)	43.8 (15)	19	* *	126	19.0 (25)	124	1.6 (4)	124	2.4 (9)
Trempealeau	210	12.4 (58)	38.1 (4)	58	* *	* *	41	* *	187	22.5 (39)	195	5.1 (46)	195	4.6 (24)
Vernon Cty	164	10.4 (37)	53.0 (62)	130	20.8 (17)	50.0 (44)	19	* *	140	15.7 (14)	149	4.0 (29)	149	4.0 (19)
Vilas	82	* *	* *	27	* *	* *	15	* *	59	* *	67	* *	67	* *
Walworth Cty	532	8.3 (10)	40.2 (5)	489	26.0 (43)	46.2 (27)	81	* *	462	14.1 (11)	472	1.5 (3)	472	1.3 (4)
Wash/Ozaukee	548	10.0 (35)	44.9 (32)	470	21.3 (18)	48.1 (37)	102	26.5 (7)	476	16.4 (16)	482	5.4 (52)	482	7.7 (50)
Washburn Cty	122	10.7 (43)	41.8 (13)	96	* *	* *	22	* *	102	17.6 (22)	105	1.9 (5)	105	3.8 (18)
Waukesha Cty	691	13.5 (62)	40.2 (5)	618	22.3 (23)	47.4 (36)	120	22.5 (3)	553	19.2 (26)	629	4.8 (44)	629	4.9 (29)
Waupaca Cty	240	13.8 (63)	45.0 (34)	220	30.5 (56)	40.5 (4)	31	* *	189	13.8 (10)	210	3.8 (25)	210	7.6 (49)
Wee Care	2,326	8.5 (14)	49.8 (54)	1,369	27.0 (47)	44.5 (19)	431	50.6 (17)	1,986	38.2 (59)	2,134	3.0 (15)	2,134	6.0 (41)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
West Allis	401	10.7 (43)	44.1 (27)	343	22.7 (28)	49.6 (41)	87	* *	356	28.1 (52)	363	6.6 (60)	363	10.7 (63)
Winnebago	733	11.6 (54)	42.8 (18)	557	20.5 (14)	46.1 (25)	180	27.2 (8)	644	18.3 (23)	667	4.3 (33)	667	4.3 (20)
Wood Cty	411	10.5 (40)	46.7 (42)	352	20.2 (12)	46.6 (32)	93	* *	329	27.7 (51)	374	1.9 (5)	374	1.3 (4)
<b>Wisconsin</b>	<b>34,504</b>	<b>9.7</b>	<b>46.4</b>	<b>24,164</b>	<b>24.7</b>	<b>45.1</b>	<b>5,810</b>	<b>33.1</b>	<b>29,138</b>	<b>25.8</b>	<b>31,161</b>	<b>4.1</b>	<b>31,161</b>	<b>5.6</b>
<b>Nation (Prior Year)</b>	<b>1,081,550</b>	<b>11.2</b>	<b>43.7</b>	<b>872,976</b>	<b>24.8</b>	<b>43.1</b>	<b>161,665</b>	<b>32.2</b>	<b>359,221</b>	<b>29.2</b>	<b>150,176</b>	<b>3.6</b>	<b>164,043</b>	<b>3.0</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

**Comparison of Maternal Behavioral Indicators by Local Agency**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>		
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
16th Street	1,496	76.3 (62)	2.7 (45)	1,672	46.7 (6)	1,427	12.7 (53)	1,307	71.8 (60)
Aurora	765	77.1 (59)	4.7 (58)	822	32.8 (41)	742	7.5 (61)	662	82.6 (48)
Barron Cty	279	92.1 (2)	0.7 (8)	300	43.7 (8)	253	24.9 (18)	227	89.9 (17)
Bay Area	193	88.6 (21)	1.6 (34)	211	43.1 (10)	183	28.4 (9)	168	89.3 (20)
Buffalo	98	* *	* *	102	35.3 (29)	89	* *	80	* *
Burnett Cty	74	* *	* *	81	* *	71	* *	66	* *
Calumet Cty	141	83.0 (44)	3.5 (51)	151	27.2 (57)	142	21.8 (24)	106	84.9 (39)
Chippewa	336	92.0 (3)	0.9 (15)	361	33.5 (37)	307	29.0 (7)	286	88.5 (24)
City Mad	1,710	83.3 (42)	1.9 (38)	1,809	24.5 (60)	1,706	18.1 (37)	1,358	83.2 (46)
Clark Cty	174	90.8 (7)	0.0 (1)	173	24.3 (62)	158	31.6 (5)	131	94.7 (3)
Columbia Cty	221	83.3 (42)	2.3 (41)	239	32.6 (43)	210	39.5 (2)	178	91.6 (7)
Cudahy	679	83.8 (39)	1.0 (20)	722	25.8 (59)	651	18.3 (36)	496	87.3 (31)
Dane Cty	179	89.4 (13)	0.6 (6)	249	32.9 (39)	135	23.0 (22)	179	93.9 (4)
Dodge Cty	325	88.6 (21)	1.2 (27)	356	32.3 (45)	314	16.6 (41)	250	87.2 (32)
Door Cty	97	* *	* *	104	32.7 (42)	93	* *	84	* *
Dunn Cty	213	87.3 (27)	0.9 (15)	231	41.1 (15)	209	26.3 (15)	191	93.7 (5)
Eau Claire	546	89.2 (14)	0.0 (1)	607	39.5 (18)	526	13.5 (52)	505	84.2 (41)
Family Health	283	86.6 (31)	1.1 (23)	310	27.4 (56)	269	28.6 (8)	211	90.0 (15)
Family Plan	919	82.0 (46)	5.4 (59)	981	29.3 (55)	838	19.5 (32)	763	88.5 (24)
Florence Cty	23	* *	* *	27	* *	21	* *	23	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Fond du Lac	500	85.2 (37)	4.2 (57)	541	38.3 (22)	478	19.2 (33)	411	83.5 (44)
Forest Cty	57	* *	* *	57	* *	53	* *	49	* *
GLITC	269	76.6 (60)	1.5 (31)	284	37.7 (25)	271	9.6 (57)	244	84.0 (43)
Grant Cty	258	91.5 (5)	0.8 (11)	277	31.4 (47)	246	26.4 (14)	209	90.4 (13)
Green Cty	152	88.8 (20)	2.6 (43)	169	39.6 (17)	150	20.0 (28)	139	89.2 (21)
Health	875	81.3 (49)	1.6 (34)	953	38.6 (21)	864	12.2 (54)	741	80.8 (52)
Jackson	139	87.1 (30)	1.4 (29)	148	41.9 (12)	127	26.8 (12)	120	75.8 (57)
Jefferson Cty	392	80.6 (52)	6.6 (63)	447	33.3 (38)	305	20.3 (27)	353	85.3 (37)
Juneau	303	85.8 (34)	5.9 (61)	317	38.2 (23)	283	27.9 (10)	255	86.3 (34)
Kenosha Cty	1,034	80.5 (53)	3.5 (51)	1,107	29.8 (54)	952	16.1 (46)	795	85.3 (37)
Kewaunee Cty	69	* *	* *	79	* *	71	* *	59	* *
La Crosse Cty	586	90.3 (9)	0.9 (15)	636	41.5 (14)	531	21.1 (25)	479	63.3 (61)
Marinette	229	87.3 (27)	3.9 (55)	243	41.6 (13)	208	24.5 (20)	200	88.0 (28)
Menominee	132	72.7 (63)	3.8 (54)	144	52.1 (3)	133	8.3 (59)	123	95.1 (2)
Mil. City	2,226	77.8 (58)	2.8 (46)	2,314	32.9 (39)	2,175	14.2 (51)	1,921	80.8 (52)
MLK	875	79.0 (57)	1.1 (23)	946	31.4 (47)	887	11.0 (55)	757	82.7 (47)
Monitowoc	381	84.5 (38)	3.4 (50)	425	48.9 (4)	367	17.7 (38)	356	84.6 (40)
Monroe Cty	290	86.2 (33)	0.0 (1)	303	27.1 (58)	267	19.9 (30)	222	80.2 (54)
NEW Comm	1,418	79.1 (56)	2.6 (43)	1,528	36.8 (26)	1,366	16.5 (42)	1,247	76.5 (55)
NHA	1,053	88.5 (24)	5.8 (60)	1,135	30.1 (52)	1,033	14.5 (50)	885	87.6 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**

**Comparison of Maternal Behavioral Indicators by Local Agency**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>		<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NWCSA	280	87.5 (26)	1.1 (23)	306	42.8 (11)	265	25.3 (17)	230	89.1 (22)
Oconto Cty	151	90.1 (10)	0.7 (8)	153	37.9 (24)	147	22.4 (23)	123	93.5 (6)
Oneida	158	76.6 (60)	1.9 (38)	168	36.3 (27)	155	6.5 (62)	143	90.2 (14)
Oneida Cty	164	85.4 (36)	3.0 (47)	180	33.9 (34)	138	34.1 (4)	127	91.3 (9)
Outagamie	623	83.5 (41)	1.3 (28)	713	34.2 (32)	618	24.6 (19)	544	80.9 (50)
Pierce Cty	163	89.0 (17)	2.5 (42)	182	32.4 (44)	147	16.3 (45)	132	87.9 (29)
Polk Cty	226	87.2 (29)	3.1 (48)	251	35.5 (28)	210	26.2 (16)	181	90.6 (11)
Portage Cty	300	82.7 (45)	1.7 (37)	318	43.4 (9)	149	16.1 (46)	239	85.8 (35)
Price Cty	97	* *	* *	107	53.3 (2)	93	* *	86	* *
Racine	417	80.8 (50)	0.7 (8)	477	40.3 (16)	390	7.9 (60)	387	73.1 (58)
Rusk Cty	118	89.0 (17)	0.8 (11)	116	46.6 (7)	112	38.4 (3)	95	* *
Sauk Cty	334	88.6 (21)	1.5 (31)	363	30.6 (50)	294	15.3 (48)	180	90.0 (15)
Sawyer Cty	102	89.2 (14)	5.9 (61)	113	54.9 (1)	92	* *	84	* *
Seeds	847	81.6 (48)	0.1 (4)	905	39.0 (20)	857	19.1 (34)	766	83.4 (45)
Shawano	191	88.5 (24)	3.7 (53)	204	33.8 (36)	175	30.9 (6)	157	95.5 (1)
Sheboygan	549	80.7 (51)	1.6 (34)	592	22.5 (64)	561	16.4 (43)	454	87.0 (33)
St. Croix Cty	276	89.9 (12)	0.4 (5)	306	34.0 (33)	257	24.1 (21)	215	91.6 (7)
SWCAP	298	90.6 (8)	1.0 (20)	333	31.8 (46)	286	19.6 (31)	239	89.1 (22)
Taylor Cty	129	89.1 (16)	0.8 (11)	129	34.9 (30)	117	26.5 (13)	106	89.6 (19)
Tremp	198	86.4 (32)	1.5 (31)	203	31.0 (49)	189	16.4 (43)	158	72.8 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>											
<b>Local Agency</b>	<b>Medical Care</b>				<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>		<b>1st Trimester</b>			<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Vernon Cty	145	94.5 (1)	1.4 (29)		163	39.3 (19)		126	42.9 (1)	126	76.2 (56)
Vilas	68	* *	* *		81	* *		62	* *	63	* *
Walworth Cty	490	81.8 (47)	1.0 (20)		531	29.9 (53)		440	20.0 (28)	386	81.9 (49)
Wash/Oz	491	90.0 (11)	0.6 (6)		539	34.3 (31)		477	19.1 (34)	417	88.2 (26)
Washburn Cty	106	91.5 (5)	0.9 (15)		118	48.3 (5)		105	21.0 (26)	101	91.1 (10)
Waukesha Cty	626	85.6 (35)	1.1 (23)		682	24.5 (60)		609	17.6 (39)	473	90.5 (12)
Waupaca Cty	218	91.7 (4)	0.9 (15)		236	33.9 (34)		201	8.5 (58)	170	88.2 (26)
Wee Care	2,124	79.8 (55)	3.3 (49)		2,250	30.2 (51)		2,004	9.8 (56)	1,768	80.9 (50)
West Allis	362	80.1 (54)	3.9 (55)		384	22.9 (63)		361	27.4 (11)	275	89.8 (18)
Winnebago	675	83.7 (40)	2.1 (40)		727	22.4 (65)		680	14.6 (49)	575	84.2 (41)
Wood Cty	370	88.9 (19)	0.8 (11)		384	20.8 (66)		382	17.3 (40)	312	85.6 (36)
<b>Wisconsin</b>	<b>31,285</b>	<b>83.3</b>	<b>2.3</b>		<b>33,775</b>	<b>33.7</b>		<b>29,810</b>	<b>17.6</b>	<b>26,148</b>	<b>83.6</b>
<b>Nation (Prior Year)</b>	<b>579,816</b>	<b>78.5</b>	<b>5.6</b>		<b>944,971</b>	<b>31.2</b>		<b>38,592</b>	<b>15.7</b>	<b>206,967</b>	<b>74.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
16th Street	1,565	16.2 (1)	1,526	8.4 (1)	192	38.5 (3)	1,460	13.6 (1)	1,524	6.2 (23)
Aurora	813	21.5 (3)	753	13.1 (3)	117	29.9 (15)	698	25.4 (39)	754	14.7 (59)
Barron Cty	289	46.4 (48)	281	28.8 (42)	116	23.3 (26)	254	18.9 (16)	281	4.6 (9)
Bay Area	203	53.2 (61)	192	30.2 (46)	86	* *	179	18.4 (14)	192	4.7 (10)
Buffalo	99	* *	96	* *	39	* *	85	* *	98	* *
Burnett Cty	79	* *	75	* *	34	* *	69	* *	75	* *
Calumet Cty	149	39.6 (26)	147	31.3 (50)	44	* *	114	19.3 (20)	147	6.1 (20)
Chippewa	354	52.0 (58)	338	31.7 (52)	152	26.3 (20)	300	23.0 (31)	337	7.1 (32)
City Mad	1,793	29.4 (8)	1,733	17.1 (7)	354	29.4 (17)	1,452	16.0 (5)	1,727	4.9 (12)
Clark Cty	180	36.7 (20)	175	25.7 (31)	51	* *	141	17.7 (12)	175	8.6 (41)
Columbia Cty	236	41.5 (31)	221	24.4 (27)	66	* *	192	25.5 (40)	218	5.5 (16)
Cudahy	709	45.3 (43)	686	24.8 (30)	216	34.7 (6)	538	35.1 (58)	686	25.5 (64)
Dane Cty	239	28.9 (7)	181	19.3 (12)	34	* *	204	17.6 (10)	179	3.9 (4)
Dodge Cty	343	39.7 (27)	330	24.5 (28)	99	* *	273	20.5 (23)	330	5.2 (14)
Door Cty	104	43.3 (38)	98	* *	38	* *	85	* *	98	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Dunn Cty	234	43.6 (39)	213	20.7 (13)	85	* *	198	17.2 (9)	213	4.2 (8)
Eau Claire	600	41.8 (32)	554	27.3 (35)	195	24.6 (23)	528	18.4 (14)	555	4.0 (5)
Family Health	283	36.7 (20)	272	31.6 (51)	72	* *	224	26.8 (42)	272	10.7 (51)
Family Plan	966	38.3 (23)	933	27.3 (35)	283	20.1 (29)	783	21.6 (28)	934	7.2 (34)
Florence Cty	26	* *	23	* *	6	* *	23	* *	23	* *
Fond du Lac	530	47.4 (50)	513	27.3 (35)	196	31.6 (13)	442	21.5 (27)	509	7.1 (32)
Forest Cty	60	* *	58	* *	15	* *	52	* *	58	* *
GLITC	293	57.3 (63)	279	35.1 (61)	140	25.0 (22)	263	36.9 (60)	277	11.6 (53)
Grant Cty	270	36.3 (18)	262	22.5 (20)	79	* *	221	19.0 (17)	264	6.1 (20)
Green Cty	168	50.0 (55)	153	28.8 (42)	67	* *	151	37.7 (61)	153	11.8 (54)
Health	950	34.5 (15)	884	17.9 (10)	236	30.5 (14)	796	21.9 (29)	883	8.0 (37)
Jackson	143	44.1 (41)	141	34.0 (56)	53	* *	124	28.2 (46)	141	6.4 (25)
Jefferson Cty	433	36.7 (20)	415	23.6 (24)	124	25.8 (21)	370	21.1 (25)	414	10.1 (49)
Juneau	319	41.4 (30)	305	32.5 (53)	105	13.3 (34)	282	35.1 (58)	303	20.5 (62)
Kenosha Cty	1,037	33.6 (14)	1,030	21.6 (16)	258	33.7 (10)	833	29.7 (50)	1,030	9.3 (42)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee Cty	77	* *	73	* *	24	* *	63	* *	73	* *
La Crosse Cty	639	32.9 (13)	597	20.8 (14)	157	17.8 (32)	545	14.5 (3)	596	3.4 (3)
Marinette	238	50.4 (57)	226	32.7 (54)	97	* *	206	25.7 (41)	226	9.3 (42)
Menominee	145	67.6 (65)	136	38.2 (64)	83	* *	133	28.6 (47)	135	5.2 (14)
Mil. City	2,335	22.4 (4)	2,235	13.2 (4)	394	29.7 (16)	2,026	23.3 (32)	2,233	11.2 (52)
MLK	948	30.6 (10)	884	17.3 (8)	209	28.2 (18)	821	34.6 (57)	882	16.1 (60)
Monitowoc	416	47.1 (49)	394	30.7 (47)	163	19.6 (30)	362	27.9 (45)	396	6.6 (26)
Monroe Cty	289	50.2 (56)	290	34.5 (59)	114	19.3 (31)	240	24.2 (35)	288	10.4 (50)
NEW Comm	1,517	36.3 (18)	1,453	21.7 (18)	437	31.8 (12)	1,306	18.3 (13)	1,451	6.8 (29)
NHA	1,099	43.1 (37)	1,045	27.6 (39)	370	24.6 (23)	935	29.8 (51)	1,041	9.4 (45)
NWCSA	300	54.3 (62)	275	34.5 (59)	128	20.3 (28)	266	28.6 (47)	274	6.2 (23)
Oconto Cty	153	48.4 (53)	151	35.8 (62)	61	* *	127	30.7 (53)	151	11.9 (55)
Oneida	163	46.0 (47)	158	22.2 (19)	63	* *	147	34.0 (56)	156	12.2 (56)
Oneida Cty	178	61.2 (64)	160	34.4 (58)	75	* *	139	27.3 (43)	160	8.1 (39)
Outagamie	666	44.0 (40)	639	23.6 (24)	219	34.2 (7)	570	16.8 (8)	640	4.8 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Pierce Cty	177	52.5 (59)	169	36.1 (63)	72	* *	144	33.3 (55)	168	9.5 (47)
Polk Cty	236	47.9 (52)	230	27.4 (38)	88	* *	188	27.7 (44)	229	5.7 (18)
Portage Cty	315	35.6 (17)	298	18.5 (11)	90	* *	278	16.5 (7)	299	5.0 (13)
Price Cty	107	42.1 (34)	101	26.7 (32)	39	* *	97	* *	101	6.9 (31)
Racine	452	27.0 (5)	440	17.5 (9)	104	24.0 (25)	410	22.0 (30)	440	6.1 (20)
Rusk Cty	119	40.3 (28)	117	30.8 (48)	40	* *	100	25.0 (37)	117	6.8 (29)
Sauk Cty	351	39.0 (24)	335	29.3 (45)	102	17.6 (33)	289	38.1 (62)	335	20.9 (63)
Sawyer Cty	110	52.7 (60)	105	22.9 (22)	44	* *	96	* *	104	5.8 (19)
Seeds	894	27.1 (6)	852	13.3 (5)	200	40.5 (2)	814	17.6 (10)	850	8.0 (37)
Shawano	201	41.8 (32)	195	26.7 (32)	67	* *	162	24.7 (36)	194	9.3 (42)
Sheboygan	588	35.0 (16)	553	21.3 (15)	137	21.9 (27)	479	20.0 (21)	552	7.4 (35)
St. Croix Cty	290	44.1 (41)	273	26.7 (32)	93	* *	232	19.0 (17)	273	9.9 (48)
SWCAP	314	45.9 (46)	299	28.8 (42)	94	* *	250	19.2 (19)	300	6.7 (28)
Taylor Cty	130	40.8 (29)	130	30.8 (48)	44	* *	111	30.6 (52)	130	12.3 (57)
Tremp	208	30.8 (11)	194	21.6 (16)	48	* *	174	14.4 (2)	195	3.1 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Vernon Cty	161	30.4 (9)	149	22.8 (21)	39	* *	135	16.3 (6)	150	3.3 (2)
Vilas	79	* *	67	* *	27	* *	70	* *	67	* *
Walworth Cty	506	32.2 (12)	493	16.6 (6)	115	42.6 (1)	402	14.7 (4)	492	4.1 (6)
Wash/Oz	525	47.8 (51)	490	23.3 (23)	193	34.2 (7)	436	21.1 (25)	490	4.1 (6)
Washburn Cty	117	45.3 (43)	106	34.0 (56)	49	* *	108	20.4 (22)	106	7.5 (36)
Waukesha Cty	676	42.9 (36)	632	23.9 (26)	183	33.9 (9)	503	20.9 (24)	632	5.5 (16)
Waupaca Cty	237	48.5 (54)	225	33.8 (55)	85	* *	180	31.1 (54)	223	17.5 (61)
Wee Care	2,186	20.4 (2)	2,139	10.8 (2)	334	36.2 (4)	1,855	23.8 (34)	2,135	12.7 (58)
West Allis	375	45.3 (43)	363	24.5 (28)	124	35.5 (5)	308	28.9 (49)	362	9.4 (45)
Winnebago	711	42.1 (34)	661	28.4 (40)	211	27.0 (19)	593	23.6 (33)	663	6.6 (26)
Wood Cty	399	39.1 (25)	376	28.7 (41)	114	32.5 (11)	322	25.2 (38)	375	8.5 (40)
<b>Wisconsin</b>	<b>33,094</b>	<b>35.6</b>	<b>31,575</b>	<b>21.6</b>	<b>8,882</b>	<b>28.0</b>	<b>27,956</b>	<b>22.7</b>	<b>31,534</b>	<b>8.8</b>
<b>Nation (Prior Year)</b>	<b>847,411</b>	<b>27.2</b>	<b>692,984</b>	<b>16.8</b>	<b>156,923</b>	<b>35.0</b>	<b>514,582</b>	<b>22.5</b>	<b>443,951</b>	<b>24.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
16th Street	1,489	6.6 (22)	8.7 (33)	1,510	9.9 (33)	1,315	3.1 (36)	1,508	85.9 (1)
Aurora	680	12.4 (62)	4.3 (2)	715	12.6 (54)	567	6.2 (61)	695	50.5 (61)
Barron Cty	259	7.3 (40)	8.9 (38)	280	10.0 (35)	228	3.1 (36)	272	70.2 (24)
Bay Area	191	4.7 (7)	14.7 (62)	192	8.3 (12)	173	2.9 (31)	194	68.6 (30)
Buffalo	93	* * *	* *	96	* *	86	* *	97	* *
Burnett Cty	71	* * *	* *	74	* *	64	* *	71	* *
Calumet Cty	145	9.0 (56)	8.3 (27)	142	9.9 (33)	127	2.4 (18)	146	60.3 (50)
Chippewa	333	7.2 (38)	5.4 (6)	339	9.1 (20)	303	2.3 (14)	338	70.7 (21)
City Mad	1,658	7.7 (46)	8.6 (32)	1,670	9.0 (18)	1,438	2.9 (31)	1,695	77.7 (8)
Clark Cty	171	9.4 (57)	7.6 (21)	164	9.8 (29)	146	2.7 (25)	171	72.5 (16)
Columbia Cty	210	6.7 (26)	11.9 (57)	213	6.1 (1)	193	4.1 (55)	214	67.8 (32)
Cudahy	628	7.6 (44)	8.8 (36)	665	12.0 (50)	542	2.4 (18)	646	67.8 (32)
Dane Cty	176	7.4 (42)	10.2 (47)	174	6.3 (2)	161	3.7 (49)	178	78.1 (4)
Dodge Cty	314	7.3 (40)	8.0 (23)	326	11.7 (49)	273	2.6 (23)	323	77.7 (8)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Door Cty	93	* * * *			95	* *	86	* *	95	* *
Dunn Cty	205	4.9 (9)	11.7	(55)	207	8.2 (9)	183	2.2 (11)	206	66.0 (38)
Eau Claire	545	6.6 (22)	6.4	(11)	544	9.7 (25)	484	3.3 (43)	548	72.1 (18)
Family Health	262	6.9 (30)	9.9	(42)	275	9.8 (29)	234	2.6 (23)	272	55.1 (56)
Family Plan	901	7.4 (42)	7.2	(19)	920	7.9 (8)	810	2.8 (29)	912	59.8 (51)
Florence Cty	22	* * * *			22	* *	19	* *	22	* *
Fond du Lac	488	6.6 (22)	6.6	(13)	502	10.2 (37)	429	2.3 (14)	496	67.9 (31)
Forest Cty	56	* * * *			54	* *	45	* *	58	* *
GLITC	260	4.2 (3)	12.7	(60)	256	9.8 (29)	222	2.3 (14)	260	53.8 (57)
Grant Cty	253	6.7 (26)	6.7	(14)	258	7.4 (6)	228	2.2 (11)	258	67.1 (34)
Green Cty	150	4.0 (1)	6.0	(9)	148	10.8 (42)	130	0.8 (2)	148	62.2 (45)
Health	859	8.7 (55)	8.4	(29)	861	11.3 (45)	738	2.7 (25)	867	63.2 (41)
Jackson	138	7.2 (38)	10.9	(51)	138	10.9 (43)	120	2.5 (21)	140	77.9 (5)
Jefferson Cty	399	4.5 (5)	10.0	(44)	404	9.7 (25)	345	1.4 (5)	402	72.6 (15)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Juneau	287	4.5 (5)	9.4 (40)		286	8.4 (15)	257	1.9 (8)	289	68.9 (29)
Kenosha Cty	1,023	8.1 (50)	5.5 (7)		1,025	10.5 (39)	891	3.9 (52)	1,028	61.0 (48)
Kewaunee Cty	68	* * * *			71	* *	59	* *	70	* *
La Crosse Cty	579	6.2 (17)	11.9 (57)		568	11.1 (44)	490	2.2 (11)	584	72.3 (17)
Marinette	221	8.1 (50)	10.0 (44)		224	11.6 (47)	188	3.2 (41)	224	49.1 (63)
Menominee	128	8.6 (54)	12.5 (59)		127	12.6 (54)	107	3.7 (49)	129	69.0 (27)
Mil. City	2,092	10.0 (58)	4.4 (3)		2,103	12.1 (51)	1,743	4.2 (56)	2,132	53.1 (58)
MLK	811	12.8 (63)	3.9 (1)		844	12.9 (57)	674	5.3 (59)	828	57.1 (54)
Monitowoc	392	6.4 (20)	8.4 (29)		397	9.1 (20)	350	3.1 (36)	399	56.9 (55)
Monroe Cty	276	5.4 (12)	8.3 (27)		282	8.9 (16)	246	1.6 (7)	281	66.5 (35)
NEW Comm	1,403	7.6 (44)	8.1 (24)		1,405	12.1 (51)	1,191	3.1 (36)	1,422	63.9 (40)
NHA	1,015	7.1 (36)	9.4 (40)		1,036	9.7 (25)	901	2.8 (29)	1,033	50.1 (62)
NWCSA	273	4.8 (8)	9.9 (42)		271	10.0 (35)	238	1.3 (4)	273	73.3 (12)
Oconto Cty	148	6.8 (29)	8.1 (24)		143	7.0 (4)	129	4.7 (58)	148	50.7 (60)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Oneida	149	4.0 (1)	18.8 (63)		151	10.6 (41)	128	0.0 (1)	154	61.7 (46)
Oneida Cty	154	5.2 (11)	7.1 (17)		158	9.5 (23)	140	2.1 (9)	156	62.8 (42)
Outagamie	598	8.4 (53)	6.4 (11)		632	8.2 (9)	540	3.7 (49)	615	62.4 (43)
Pierce Cty	167	4.2 (3)	10.8 (50)		164	7.3 (5)	149	2.7 (25)	169	66.3 (37)
Polk Cty	221	6.3 (19)	10.4 (49)		227	6.6 (3)	205	3.4 (45)	226	77.9 (5)
Portage Cty	285	7.7 (46)	11.2 (53)		290	12.1 (51)	243	2.5 (21)	291	72.9 (14)
Price Cty	96	* * *	* *		99	* *	88	* *	98	* *
Racine	434	11.5 (60)	4.4 (3)		433	14.8 (63)	358	3.9 (52)	433	51.0 (59)
Rusk Cty	114	6.1 (15)	11.4 (54)		110	8.2 (9)	100	4.0 (54)	114	58.8 (53)
Sauk Cty	325	4.9 (9)	11.7 (55)		336	8.3 (12)	298	2.3 (14)	333	77.8 (7)
Sawyer Cty	100	6.0 (14)	6.0 (9)		102	9.8 (29)	89	* *	100	72.0 (19)
Seeds	832	7.0 (34)	8.7 (33)		836	7.8 (7)	754	2.9 (31)	835	73.3 (12)
Shawano	184	7.1 (36)	8.7 (33)		190	13.7 (60)	155	3.2 (41)	190	69.5 (25)
Sheboygan	533	6.4 (20)	6.9 (15)		536	9.7 (25)	473	3.0 (34)	537	65.5 (39)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
St. Croix Cty	275	6.9 (30)	10.9	(51)	273	9.2 (22)	243	3.3 (43)	280	73.9 (11)
SWCAP	291	6.2 (17)	10.0	(44)	295	10.2 (37)	256	3.1 (36)	295	69.2 (26)
Taylor Cty	122	6.6 (22)	8.2	(26)	124	12.9 (57)	104	1.0 (3)	123	70.7 (21)
Tremp	179	5.6 (13)	8.4	(29)	182	13.7 (60)	146	2.1 (9)	184	76.6 (10)
Vernon Cty	147	6.1 (15)	14.3	(61)	148	9.5 (23)	131	1.5 (6)	149	80.5 (2)
Vilas	65	*	*	*	66	*	*	60	*	*
Walworth Cty	488	7.0 (34)	10.2	(47)	494	8.3 (12)	440	3.4 (45)	496	78.4 (3)
Wash/Oz	466	6.7 (26)	7.7	(22)	478	10.5 (39)	409	3.4 (45)	478	69.0 (27)
Washburn Cty	103	7.8 (48)	5.8	(8)	101	13.9 (62)	85	*	104	61.5 (47)
Waukesha Cty	611	6.9 (30)	8.8	(36)	620	9.0 (18)	542	3.0 (34)	622	70.6 (23)
Waupaca Cty	198	12.1 (61)	9.1	(39)	219	13.2 (59)	166	5.4 (60)	207	62.3 (44)
Wee Care	2,012	11.0 (59)	5.0	(5)	2,055	12.7 (56)	1,686	4.3 (57)	2,050	59.2 (52)
West Allis	352	8.2 (52)	7.1	(17)	344	11.6 (47)	296	2.7 (25)	356	71.9 (20)
Winnebago	642	7.8 (48)	7.3	(20)	661	11.3 (45)	561	3.4 (45)	659	66.5 (35)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wood Cty	347	6.9 (30)	6.9 (15)	347	8.9 (16)	288	2.4 (18)	356	60.4 (49)
<b>Wisconsin</b>	<b>30,255</b>	<b>7.8</b>	<b>7.8</b>	<b>30,697</b>	<b>10.4</b>	<b>26,286</b>	<b>3.2</b>	<b>30,747</b>	<b>65.8</b>
<b>Nation (Prior Year)</b>	<b>972,241</b>	<b>8.3</b>	<b>6.8</b>	<b>873,261</b>	<b>12.1</b>	<b>802,123</b>	<b>3.7</b>	<b>897,379</b>	<b>64.2</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaskan Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
16th Street	4,892	8.5	5.8	79.7	0.3	4.9	0.5	0.3
Aurora	2,345	6.8	80.6	4.1	0.3	5.6	0.5	2.1
Barron Cty	877	87.3	3.9	4.4	1.1	0.8	0.3	2.1
Bay Area	638	87.5	0.0	1.9	5.8	1.4	2.0	1.4
Buffalo	285	91.9	0.0	4.6	0.0	0.4	1.8	1.4
Burnett Cty	280	93.6	0.0	1.4	2.5	0.4	1.4	0.7
Calumet Cty	431	73.8	2.3	13.9	0.2	7.9	0.2	1.6
Chippewa	1,040	95.0	0.3	1.2	0.3	1.3	0.6	1.4
City Mad	3,983	36.0	24.3	26.5	0.4	9.8	2.0	1.2
Clark Cty	539	80.0	0.2	16.5	0.9	0.0	1.3	1.1
Columbia Cty	657	83.7	1.1	12.3	0.2	0.5	1.5	0.8
Cudahy	1,844	80.5	2.3	12.1	1.0	1.5	1.6	1.0
Dane Cty	1,723	49.6	12.3	29.4	0.4	4.8	1.8	1.6
Dodge Cty	1,039	78.2	1.2	18.4	0.2	0.5	0.6	1.1
Door Cty	316	85.8	0.6	10.4	0.0	0.9	0.6	1.6
Dunn Cty	717	84.2	1.0	2.8	0.4	8.5	1.4	1.7
Eau Claire	1,690	79.9	1.6	3.3	0.9	10.7	2.2	1.4
Family Health	812	72.2	0.4	24.6	0.1	0.6	0.6	1.5
Family Plan	2,647	74.3	0.9	4.7	0.6	18.0	0.6	0.8
Florence Cty	73	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Fond du Lac	1,505	73.5	2.3	17.9	0.5	3.9	1.2	0.8
Forest Cty	179	93.9	0.0	1.7	2.8	0.6	1.1	0.0
GLITC	857	13.1	0.2	3.0	78.9	0.0	3.5	1.3
Grant Cty	801	92.0	0.4	4.9	0.0	0.7	1.0	1.0
Green Cty	504	88.9	0.4	9.1	0.0	0.0	0.2	1.4
Health	2,921	43.9	23.5	29.6	0.2	0.5	1.3	1.0
Jackson	420	88.3	0.0	3.8	6.4	0.0	0.0	1.4
Jefferson Cty	1,251	63.3	1.0	33.5	0.1	0.7	0.6	0.9
Juneau	786	91.2	0.3	3.9	0.4	0.6	2.9	0.6
Kenosha Cty	3,052	50.1	17.8	29.5	0.2	0.9	1.1	0.5
Kewaunee Cty	215	81.4	0.5	14.9	1.9	0.0	0.0	1.4
La Crosse Cty	1,873	76.4	3.7	3.3	0.8	13.1	1.2	1.4
Marinette	788	94.8	0.0	2.2	0.5	0.1	0.8	1.6
Menominee	386	4.4	0.0	4.1	68.9	0.3	20.7	1.6
Mil. City	6,854	15.3	60.1	17.5	0.5	4.0	1.5	1.1
MLK	2,636	6.6	83.6	5.3	0.4	1.3	1.4	1.3
Monitowoc	1,187	75.1	1.6	11.6	0.5	8.7	1.2	1.3
Monroe Cty	837	83.6	1.1	13.3	0.4	0.5	0.6	0.6
NEW Comm	4,266	56.7	4.2	27.0	3.0	7.4	0.7	1.0
NHA	3,244	67.0	10.8	18.1	0.1	1.2	1.1	1.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 3F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaskan Native %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
NWCSA	879	89.2	0.9	1.9	1.5	0.8	3.4	2.3
Oconto Cty	438	95.2	0.5	2.1	0.5	0.5	0.0	1.4
Oneida	421	8.1	0.0	6.7	79.6	0.2	3.8	1.7
Oneida Cty	481	93.3	0.4	2.7	0.4	0.8	0.8	1.5
Outagamie	1,995	68.1	2.5	14.6	1.2	11.8	0.7	1.2
Pierce Cty	498	89.6	1.2	3.6	1.2	0.8	1.0	2.6
Polk Cty	729	93.0	0.4	1.9	1.1	0.1	1.0	2.5
Portage Cty	904	76.7	0.8	8.6	0.3	11.8	0.3	1.4
Price Cty	271	95.2	0.0	1.5	0.4	2.6	0.0	0.4
Racine	1,139	26.4	38.6	31.4	0.4	0.5	1.1	1.6
Rusk Cty	317	93.4	0.6	0.9	0.0	0.6	2.2	2.2
Sauk Cty	863	80.9	0.7	16.1	0.8	0.3	0.1	1.0
Sawyer Cty	299	79.3	0.3	2.7	13.4	0.3	2.0	2.0
Seeds	2,466	32.4	4.5	57.7	1.1	1.7	1.8	0.8
Shawano	592	85.0	1.2	6.8	3.9	0.7	1.7	0.8
Sheboygan	1,673	59.2	2.5	19.9	0.2	17.2	0.4	0.6
St. Croix Cty	833	87.0	1.0	5.2	1.0	3.4	0.8	1.7
SWCAP	1,017	91.8	0.2	5.5	0.2	0.2	0.5	1.6
Taylor Cty	296	88.9	0.3	8.8	0.0	0.3	1.4	0.3
Trempealeau	526	75.1	0.2	21.7	0.4	0.4	1.1	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaskan Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
Vernon Cty	435	91.5	0.2	5.7	0.2	0.2	0.5	1.6
Vilas	206	90.8	0.0	4.4	1.5	0.0	0.5	2.9
Walworth Cty	1,552	55.2	1.4	40.3	0.6	0.5	0.8	1.2
Wash/Oz	1,461	85.8	1.4	9.2	0.4	0.7	1.1	1.4
Washburn Cty	311	92.3	0.6	4.2	0.3	0.3	1.3	1.0
Waukesha Cty	1,978	66.1	3.7	26.3	0.6	1.4	1.1	0.9
Waupaca Cty	633	89.6	0.2	7.7	0.0	0.3	0.8	1.4
Wee Care	6,550	5.9	84.5	2.2	0.2	4.6	1.3	1.3
West Allis	1,058	75.5	5.6	11.4	0.9	2.9	1.4	2.2
Winnebago	2,018	74.6	3.6	12.2	0.3	7.3	1.1	0.7
Wood Cty	1,133	83.8	1.0	7.1	0.5	5.1	1.0	1.5
Unknown Local Agencies	3	*	*	*	*	*	*	*
<b>Wisconsin</b>	<b>95,365</b>	<b>53.4</b>	<b>19.1</b>	<b>18.6</b>	<b>2.0</b>	<b>4.3</b>	<b>1.2</b>	<b>1.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
16th Street	4,892	0.7	7.4	10.9	59.6	19.7	1.6	0.1
Aurora	2,345	0.7	9.3	14.5	62.1	12.4	0.9	0.0
Barron Cty	877	0.1	3.3	17.1	65.5	13.3	0.7	0.0
Bay Area	638	0.0	3.3	9.7	64.4	20.1	2.5	0.0
Buffalo	285	0.4	3.5	13.0	63.9	18.9	0.4	0.0
Burnett Cty	280	0.4	5.0	15.4	63.6	14.3	1.4	0.0
Calumet Cty	431	0.0	5.3	14.6	60.8	18.1	1.2	0.0
Chippewa	1,040	0.2	4.6	11.6	67.8	15.1	0.7	0.0
City Mad	3,983	0.4	5.8	12.1	61.3	19.1	1.4	0.1
Clark Cty	539	0.2	5.2	16.3	63.3	13.9	0.9	0.2
Columbia Cty	657	0.2	6.2	14.9	61.3	16.0	1.4	0.0
Cudahy	1,844	0.2	3.4	11.9	66.9	16.6	1.0	0.1
Dane Cty	1,723	0.2	5.9	11.5	63.8	17.2	1.2	0.1
Dodge Cty	1,039	0.2	5.5	14.5	62.8	16.0	0.9	0.1
Door Cty	316	0.0	5.4	12.0	63.9	17.1	1.6	0.0
Dunn Cty	717	0.0	5.4	14.8	62.9	15.5	1.4	0.0
Eau Claire	1,690	0.4	4.9	13.7	66.0	14.0	0.9	0.1
Family Health	812	0.1	6.0	14.5	62.7	15.4	1.1	0.1
Family Plan	2,647	0.1	5.2	15.1	63.8	14.8	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
Florence Cty	73	*	*	*	*	*	*	*
Fond du Lac	1,505	0.2	5.7	15.9	63.5	13.5	1.2	0.1
Forest Cty	179	0.0	1.7	15.6	66.5	16.2	0.0	0.0
GLITC	857	0.5	8.1	16.2	59.6	14.9	0.6	0.1
Grant Cty	801	0.0	4.5	12.1	65.0	16.7	1.6	0.0
Green Cty	504	0.2	5.6	15.5	63.7	14.3	0.8	0.0
Health	2,921	0.6	7.8	15.3	61.1	14.3	0.9	0.0
Jackson	420	0.5	3.6	16.9	63.1	15.0	1.0	0.0
Jefferson Cty	1,251	0.3	5.0	13.1	62.7	17.3	1.6	0.0
Juneau	786	0.5	8.7	14.5	63.1	12.0	1.3	0.0
Kenosha Cty	3,052	0.3	6.9	14.0	62.5	15.4	0.8	0.0
Kewaunee Cty	215	0.0	4.2	14.4	60.9	18.6	1.4	0.5
La Crosse Cty	1,873	0.2	5.9	13.5	63.9	15.3	1.3	0.0
Marinette	788	0.0	4.2	15.1	64.8	15.1	0.8	0.0
Menominee	386	0.8	11.9	15.5	56.2	14.5	1.0	0.0
Mil. City	6,854	0.8	9.0	14.9	58.0	16.0	1.2	0.1
MLK	2,636	1.3	12.6	14.4	57.7	13.0	0.9	0.0
Monitowoc	1,187	0.3	6.4	15.1	62.1	14.9	1.2	0.1
Monroe Cty	837	0.4	4.8	15.4	63.9	14.6	1.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

Comparison of Age Distribution <sup>(2)</sup>								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
NEW Comm	4,266	0.3	6.6	13.2	61.9	16.7	1.1	0.1
NHA	3,244	0.4	7.2	15.9	61.3	14.4	0.8	0.1
NWCSA	879	0.2	6.5	12.5	67.7	12.1	1.0	0.0
Oconto Cty	438	0.2	3.9	17.6	63.0	13.9	1.1	0.2
Oneida	421	0.0	6.7	14.0	63.7	14.3	1.2	0.2
Oneida Cty	481	0.0	2.1	12.1	65.7	18.3	1.9	0.0
Outagamie	1,995	0.4	5.3	14.0	64.3	14.9	1.2	0.1
Pierce Cty	498	0.4	3.4	16.3	61.0	17.5	1.4	0.0
Polk Cty	729	0.0	6.6	15.0	62.8	14.1	1.4	0.1
Portage Cty	904	0.0	6.0	15.6	65.3	11.5	1.5	0.1
Price Cty	271	0.4	2.2	10.0	66.1	19.9	1.5	0.0
Racine	1,139	0.5	9.0	13.3	60.4	15.0	1.7	0.1
Rusk Cty	317	0.3	3.5	14.8	58.0	21.5	1.6	0.3
Sauk Cty	863	0.1	5.1	15.8	62.9	15.3	0.8	0.0
Sawyer Cty	299	0.0	5.4	15.4	63.9	14.4	1.0	0.0
Seeds	2,466	0.4	9.0	13.0	58.9	17.2	1.6	0.0
Shawano	592	0.3	4.1	12.5	64.2	17.7	1.0	0.2
Sheboygan	1,673	0.2	6.6	15.1	60.4	16.3	1.2	0.2
St. Croix Cty	833	0.5	4.8	11.3	65.1	17.3	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 4F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
SWCAP	1,017	0.0	3.5	12.4	67.0	15.2	1.9	0.0
Taylor Cty	296	0.3	5.1	12.5	65.2	16.6	0.3	0.0
Tremp	526	0.0	3.6	14.3	65.0	16.7	0.4	0.0
Vernon Cty	435	0.0	2.3	14.5	64.4	17.0	1.8	0.0
Vilas	206	0.0	4.4	10.7	59.7	21.8	3.4	0.0
Walworth Cty	1,552	0.5	5.4	13.1	62.8	17.3	0.8	0.1
Wash/Oz	1,461	0.0	3.4	13.1	66.2	16.2	1.1	0.1
Washburn Cty	311	0.0	4.5	10.3	68.8	15.8	0.6	0.0
Waukesha Cty	1,978	0.2	4.4	13.5	62.8	17.9	1.1	0.1
Waupaca Cty	633	0.0	5.5	16.4	63.5	13.0	1.4	0.2
Wee Care	6,550	0.7	10.8	16.2	56.3	14.9	1.1	0.1
West Allis	1,058	0.5	4.4	11.5	64.7	16.9	2.0	0.0
Winnebago	2,018	0.0	5.0	13.8	65.3	14.6	1.2	0.1
Wood Cty	1,133	0.2	4.1	13.9	65.3	15.8	0.6	0.1
Unknown Local Agencies	3	*	*	*	*	*	*	*
<b>Wisconsin</b>	<b>95,365</b>	<b>0.4</b>	<b>6.6</b>	<b>14.0</b>	<b>62.0</b>	<b>15.8</b>	<b>1.2</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
16th Street	4,614	6.0 (2)	48.7 (59)	3,873	30.3 (67)	35.3 (1)	558	24.0 (13)	4,211	24.3 (52)	2,937	6.0 (53)	2,932	6.2 (33)
Aurora	2,237	7.9 (7)	53.2 (67)	772	33.8 (69)	37.7 (3)	323	63.2 (41)	1,700	55.2 (70)	1,129	3.5 (20)	1,142	6.8 (45)
Barron Cty	822	10.0 (25)	46.7 (50)	707	22.3 (19)	51.1 (68)	98	* *	600	21.7 (41)	398	6.3 (57)	397	6.0 (31)
Bay Area	586	9.2 (14)	44.9 (39)	509	22.8 (23)	46.8 (51)	73	* *	500	16.0 (15)	325	4.9 (38)	327	4.9 (25)
Buffalo	259	10.4 (35)	47.1 (53)	226	26.5 (53)	47.3 (54)	50	* *	210	18.1 (23)	117	5.1 (44)	115	7.0 (48)
Burnett Cty	265	10.9 (43)	41.5 (13)	202	18.8 (2)	49.5 (66)	36	* *	182	12.6 (5)	155	1.3 (1)	155	1.3 (4)
Calumet Cty	406	8.6 (9)	37.9 (3)	344	26.2 (50)	41.0 (9)	63	* *	337	16.9 (19)	264	7.2 (65)	267	6.7 (44)
Chippewa	1,013	11.6 (52)	44.6 (34)	938	25.9 (47)	41.5 (12)	140	20.7 (4)	794	14.6 (8)	616	6.3 (57)	609	6.9 (47)
City Mad	3,640	11.3 (48)	42.7 (22)	2,370	31.2 (68)	40.1 (6)	743	30.6 (27)	2,277	22.3 (43)	2,536	4.9 (38)	2,534	4.7 (20)
Clark Cty	526	14.1 (67)	45.2 (40)	439	25.5 (43)	41.0 (9)	96	* *	424	29.2 (63)	349	2.9 (11)	353	4.8 (23)
Columbia Cty	602	14.1 (67)	39.7 (7)	487	22.2 (17)	46.4 (48)	110	23.6 (11)	516	14.9 (12)	389	3.9 (27)	392	7.7 (57)
Cudahy	1,787	9.6 (21)	42.6 (21)	1,538	19.9 (6)	48.8 (62)	352	24.7 (15)	1,554	21.4 (40)	961	5.3 (45)	967	6.5 (39)
Dane Cty	1,577	11.0 (45)	40.3 (9)	1,006	24.1 (29)	42.4 (18)	329	29.8 (26)	1,244	21.1 (38)	673	4.8 (35)	684	8.8 (62)
Dodge Cty	1,000	12.3 (58)	41.1 (11)	708	25.1 (41)	41.5 (12)	175	26.9 (22)	874	22.9 (47)	663	3.6 (22)	667	5.2 (27)
Door Cty	313	11.5 (50)	38.3 (4)	285	20.0 (8)	43.2 (26)	49	* *	265	14.7 (10)	157	4.5 (32)	164	10.4 (66)
Dunn Cty	656	9.5 (20)	47.7 (56)	457	24.1 (29)	44.4 (34)	84	* *	498	14.7 (10)	310	5.8 (50)	315	7.3 (56)
Eau Claire	1,645	11.6 (52)	42.1 (16)	422	28.2 (62)	41.9 (15)	229	21.8 (7)	1,378	24.9 (56)	1,066	4.5 (32)	1,058	3.7 (11)

- (1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>%</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Family Health	764	10.1 (28)	44.6 (34)	658	24.6 (36)	43.8 (29)	149	24.8 (16)	490	20.4 (33)	374	4.5 (32)	377	4.8 (23)
Family Plan	2,550	10.4 (35)	42.9 (24)	2,246	26.2 (50)	40.1 (6)	433	17.6 (2)	2,023	9.9 (1)	1,651	3.6 (22)	1,647	4.1 (13)
Florence Cty	71	* *	* *	60	* *	* *	10	* *	48	* *	43	* *	45	* *
Fond du Lac	1,445	11.5 (50)	42.8 (23)	1,275	28.2 (62)	40.3 (8)	190	35.3 (33)	1,124	24.6 (54)	794	6.9 (63)	791	7.0 (48)
Forest Cty	177	6.2 (3)	53.7 (68)	160	30.0 (66)	47.5 (55)	16	* *	120	18.3 (25)	103	6.8 (62)	106	8.5 (60)
GLITC	830	5.8 (1)	55.1 (69)	583	24.9 (38)	45.8 (41)	132	37.9 (36)	533	34.9 (66)	420	5.5 (47)	423	6.4 (38)
Grant Cty	761	10.1 (28)	46.5 (47)	670	21.3 (13)	47.9 (57)	118	24.6 (14)	664	17.5 (21)	447	3.8 (25)	457	6.6 (41)
Green Cty	501	12.4 (60)	42.9 (24)	450	20.9 (12)	48.4 (60)	92	* *	418	27.0 (60)	302	2.3 (4)	305	2.3 (7)
Health	2,662	10.7 (40)	47.0 (52)	2,409	41.2 (70)	37.0 (2)	446	27.1 (23)	2,220	18.6 (29)	1,253	7.6 (68)	1,282	11.3 (69)
Jackson	388	9.8 (23)	45.9 (44)	353	24.1 (29)	41.1 (11)	57	* *	299	18.4 (28)	158	1.3 (1)	157	0.6 (1)
Jefferson Cty	1,209	11.7 (54)	42.9 (24)	771	27.4 (59)	39.2 (5)	219	21.5 (6)	1,020	14.6 (8)	726	4.8 (35)	726	5.9 (30)
Juneau	739	9.2 (14)	46.8 (51)	419	23.2 (26)	46.5 (49)	98	* *	600	17.0 (20)	340	4.4 (31)	338	7.7 (57)
Kenosha Cty	2,828	10.3 (31)	45.8 (42)	2,375	25.5 (43)	44.2 (33)	539	35.6 (34)	2,286	29.4 (64)	1,341	3.4 (18)	1,351	6.8 (45)
Kewaunee Cty	212	9.0 (13)	46.2 (46)	192	22.9 (24)	49.0 (63)	24	* *	174	25.9 (59)	111	2.7 (7)	110	0.9 (2)
La Crosse Cty	1,785	12.7 (62)	40.1 (8)	1,292	24.1 (29)	45.4 (40)	209	29.2 (25)	1,303	22.3 (43)	1,235	3.3 (17)	1,223	0.9 (2)
Marinette	741	15.1 (70)	39.3 (5)	627	19.1 (4)	46.3 (46)	113	15.9 (1)	433	18.2 (24)	421	5.0 (42)	429	7.0 (48)
Menominee	381	8.7 (10)	48.8 (60)	279	29.4 (64)	42.7 (20)	24	* *	324	20.7 (36)	235	8.9 (70)	238	10.9 (68)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)			(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Mil. City	6,695	9.3 (16)	50.2 (62)	5,346	27.2 (58)	42.9 (22)	1,147	44.2 (38)	5,250	37.0 (67)	3,800	4.1 (28)	3,837	7.2 (53)
MLK	2,566	10.3 (31)	51.1 (64)	1,857	28.1 (61)	45.0 (37)	435	51.5 (40)	2,004	43.4 (69)	1,544	3.1 (14)	1,575	10.0 (64)
Monitowoc	1,123	12.1 (57)	40.7 (10)	526	25.9 (47)	46.0 (42)	158	25.3 (19)	919	24.8 (55)	716	4.2 (29)	720	5.8 (28)
Monroe Cty	793	10.1 (28)	47.4 (54)	156	26.9 (55)	42.3 (17)	151	32.5 (30)	671	25.6 (57)	461	5.4 (46)	458	3.1 (9)
NEW Comm	4,088	12.0 (56)	42.4 (18)	2,609	26.1 (49)	43.8 (29)	642	36.1 (35)	3,656	28.3 (61)	2,521	5.0 (42)	2,540	7.0 (48)
NHA	3,085	10.7 (40)	46.5 (47)	1,212	21.9 (16)	42.1 (16)	550	30.9 (28)	2,389	25.8 (58)	1,152	3.0 (13)	1,155	4.1 (13)
NWCSA	828	10.4 (35)	44.8 (37)	606	19.0 (3)	52.6 (69)	122	32.8 (31)	711	28.6 (62)	428	7.5 (67)	429	8.4 (59)
Oconto Cty	426	14.8 (69)	46.0 (45)	402	22.6 (22)	49.0 (63)	74	* *	336	15.8 (13)	264	6.1 (56)	260	6.5 (39)
Oneida	417	6.5 (4)	59.7 (70)	345	23.8 (27)	48.1 (59)	55	* *	342	20.5 (34)	272	8.1 (69)	272	6.3 (35)
Oneida Cty	470	10.0 (25)	44.3 (32)	227	26.9 (55)	44.1 (32)	68	* *	339	11.5 (3)	274	6.6 (61)	277	6.1 (32)
Outagamie	1,907	12.3 (58)	42.5 (19)	1,519	25.0 (40)	43.3 (27)	328	26.2 (21)	1,621	22.8 (46)	1,225	5.7 (48)	1,217	5.8 (28)
Pierce Cty	478	11.1 (46)	41.8 (15)	335	15.8 (1)	47.8 (56)	55	* *	350	23.4 (50)	313	3.2 (16)	315	6.3 (35)
Polk Cty	673	12.9 (63)	43.1 (27)	474	20.0 (8)	46.6 (50)	90	* *	581	21.9 (42)	381	6.0 (53)	376	8.5 (60)
Portage Cty	874	10.6 (39)	42.1 (16)	746	24.8 (37)	42.8 (21)	127	25.2 (18)	588	16.2 (17)	485	2.9 (11)	493	4.7 (20)
Price Cty	265	8.7 (10)	43.8 (29)	251	19.9 (6)	54.6 (70)	32	* *	219	16.0 (15)	164	2.4 (5)	164	4.9 (25)
Racine	1,094	9.4 (19)	52.9 (66)	907	26.8 (54)	43.0 (23)	187	40.6 (37)	954	30.0 (65)	431	3.5 (20)	432	3.5 (10)
Rusk Cty	283	7.8 (6)	48.1 (58)	218	29.4 (64)	45.0 (37)	34	* *	231	13.9 (6)	164	3.7 (24)	166	1.8 (6)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	% (Rank)	%	Number	% (Rank)	% (Rank)	Number	% (Rank)
Sauk Cty	827	11.4 (49)	47.9 (57)	696	22.4 (20)	46.3 (46)	158	25.9 (20)	696	17.7 (22)	480	6.3 (57)	482	6.2 (33)
Sawyer Cty	280	10.4 (35)	42.5 (19)	257	22.2 (17)	49.4 (65)	35	* *	206	22.3 (43)	110	2.7 (7)	113	2.7 (8)
Seeds	2,384	9.3 (16)	46.5 (47)	2,089	24.1 (29)	43.4 (28)	420	25.0 (17)	2,151	20.5 (34)	1,645	5.7 (48)	1,647	6.3 (35)
Shawano	535	9.3 (16)	44.3 (32)	408	20.6 (10)	48.0 (58)	88	* *	291	10.3 (2)	329	4.3 (30)	331	6.6 (41)
Sheboygan	1,604	9.9 (24)	41.2 (12)	1,216	28.0 (60)	41.6 (14)	310	22.9 (8)	1,436	19.4 (32)	858	6.4 (60)	854	4.2 (15)
St. Croix Cty	755	10.3 (31)	44.8 (37)	667	19.8 (5)	51.0 (67)	99	* *	509	13.9 (6)	388	7.2 (65)	383	10.2 (65)
SWCAP	959	7.5 (5)	50.4 (63)	797	25.6 (45)	44.5 (35)	163	33.7 (32)	661	21.2 (39)	587	4.8 (35)	609	10.8 (67)
Taylor Cty	292	9.6 (21)	45.2 (40)	271	24.0 (28)	46.1 (44)	42	* *	227	23.3 (48)	153	2.6 (6)	154	4.5 (18)
Trempealeau	493	13.0 (64)	37.7 (2)	201	26.9 (55)	37.8 (4)	105	23.8 (12)	391	24.0 (51)	250	6.0 (53)	251	7.2 (53)
Vernon Cty	419	10.7 (40)	49.6 (61)	349	24.9 (38)	45.3 (39)	59	* *	289	18.3 (25)	261	3.4 (18)	262	4.2 (15)
Vilas	204	13.2 (65)	35.3 (1)	109	22.9 (24)	48.6 (61)	29	* *	126	11.9 (4)	126	1.6 (3)	130	4.6 (19)
Walworth Cty	1,465	10.0 (25)	43.1 (27)	1,320	25.8 (46)	42.4 (18)	236	21.2 (5)	1,198	18.3 (25)	727	2.8 (9)	720	1.4 (5)
Wash/Ozaukee	1,418	10.9 (43)	44.0 (31)	1,211	20.6 (10)	46.1 (44)	237	23.2 (9)	1,175	15.9 (14)	851	4.9 (38)	856	7.0 (48)
Washburn Cty	303	8.9 (12)	47.5 (55)	250	21.6 (15)	46.8 (51)	39	* *	237	18.6 (29)	132	3.8 (25)	130	3.8 (12)
Waukesha Cty	1,869	11.1 (46)	39.3 (5)	1,662	24.1 (29)	44.9 (36)	371	19.9 (3)	1,330	19.2 (31)	1,086	4.9 (38)	1,105	7.2 (53)
Waupaca Cty	617	13.9 (66)	44.7 (36)	557	25.1 (41)	43.1 (24)	98	* *	436	16.3 (18)	255	5.9 (51)	254	9.4 (63)
Wee Care	6,366	8.5 (8)	51.7 (65)	4,316	26.4 (52)	44.0 (31)	1,087	49.2 (39)	4,887	40.4 (68)	4,072	3.1 (14)	4,143	6.6 (41)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 5F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>%</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
West Allis	1,017	12.4 (60)	43.9 (30)	859	22.4 (20)	46.0 (42)	207	27.1 (23)	855	24.3 (52)	496	7.1 (64)	494	11.5 (70)
Winnebago	1,904	11.8 (55)	41.5 (13)	1,547	24.2 (35)	43.1 (24)	408	23.3 (10)	1,718	20.8 (37)	1,114	5.9 (51)	1,107	4.7 (20)
Wood Cty	1,099	10.3 (31)	45.8 (42)	922	21.4 (14)	47.1 (53)	248	32.3 (29)	873	23.3 (48)	633	2.8 (9)	643	4.4 (17)
Unknown Local Agencies	3	* *	* *	3	* *	* *	2	* *	1	* *	0	* *	0	* *
<b>Wisconsin</b>	<b>90,870</b>	<b>10.2</b>	<b>45.7</b>	<b>67,545</b>	<b>26.0</b>	<b>43.3</b>	<b>15,073</b>	<b>31.1</b>	<b>72,497</b>	<b>25.3</b>	<b>52,147</b>	<b>4.7</b>	<b>52,437</b>	<b>6.2</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>			<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	
16th Street	3,042	68.9 (70)	12.1 (69)	4,582	44.3 (6)	1,432	12.6 (53)	1,307	71.8 (60)	
Aurora	1,785	76.7 (62)	4.0 (32)	2,182	30.2 (48)	756	7.7 (61)	664	82.7 (47)	
Barron Cty	574	89.9 (5)	1.2 (7)	814	42.0 (13)	255	24.7 (18)	227	89.9 (17)	
Bay Area	432	85.4 (28)	5.6 (47)	594	44.1 (7)	184	28.3 (9)	168	89.3 (20)	
Buffalo	194	81.4 (48)	5.7 (48)	261	31.8 (43)	90	* *	80	* *	
Burnett Cty	191	92.7 (3)	0.0 (1)	265	42.6 (10)	72	* *	66	* *	
Calumet Cty	310	83.2 (35)	4.2 (34)	405	25.7 (60)	146	21.9 (24)	106	84.9 (39)	
Chippewa	709	89.3 (7)	1.4 (12)	989	27.8 (56)	309	29.1 (7)	286	88.5 (24)	
City Mad	2,876	79.5 (54)	5.8 (50)	3,699	24.4 (63)	1,720	18.0 (37)	1,361	83.2 (46)	
Clark Cty	392	87.8 (18)	1.8 (16)	479	25.3 (62)	162	32.1 (5)	132	94.7 (3)	
Columbia Cty	500	75.0 (67)	7.2 (59)	627	31.1 (45)	216	39.8 (2)	179	91.6 (7)	
Cudahy	1,360	81.4 (48)	3.2 (26)	1,749	22.5 (68)	656	18.6 (36)	498	87.3 (31)	
Dane Cty	960	82.0 (42)	4.4 (38)	1,606	24.0 (64)	138	23.2 (22)	182	93.4 (6)	
Dodge Cty	703	85.6 (26)	4.4 (38)	994	25.5 (61)	319	16.3 (43)	251	87.3 (31)	
Door Cty	203	87.7 (20)	1.0 (5)	303	26.1 (58)	93	* *	84	* *	
Dunn Cty	476	83.0 (37)	5.7 (48)	663	41.5 (14)	209	26.3 (15)	191	93.7 (4)	
Eau Claire	1,087	81.9 (44)	7.2 (59)	1,552	38.7 (18)	531	13.4 (52)	507	84.2 (41)	
Family Health	611	89.2 (8)	1.0 (5)	780	26.2 (57)	270	28.9 (8)	211	90.0 (15)	
Family Plan	2,006	80.5 (51)	6.7 (57)	2,466	30.3 (47)	846	19.5 (31)	765	88.5 (24)	
Florence Cty	49	* *	* *	70	* *	21	* *	23	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Fond du Lac	1,024	85.1 (29)	5.4 (46)	1,418	37.2 (21)	482	19.5 (31)	411	83.5 (44)
Forest Cty	129	86.0 (24)	1.6 (13)	171	48.5 (3)	55	* *	49	* *
GLITC	691	73.8 (68)	2.2 (18)	781	34.7 (32)	275	10.2 (56)	246	84.1 (42)
Grant Cty	551	89.1 (10)	2.2 (18)	747	31.2 (44)	246	26.4 (13)	209	90.4 (13)
Green Cty	323	88.9 (12)	1.9 (17)	479	32.6 (40)	150	20.0 (30)	139	89.2 (22)
Health	1,942	71.4 (69)	12.8 (70)	2,703	34.4 (34)	877	12.5 (54)	743	80.8 (51)
Jackson	272	81.6 (46)	5.9 (51)	393	36.4 (24)	127	26.8 (12)	120	75.8 (57)
Jefferson Cty	831	80.6 (50)	8.5 (66)	1,176	32.7 (39)	309	20.7 (27)	354	85.3 (37)
Juneau	637	82.3 (41)	10.5 (68)	727	40.3 (16)	284	28.2 (10)	256	86.3 (34)
Kenosha Cty	2,017	79.9 (52)	6.0 (52)	2,858	29.4 (51)	959	16.0 (46)	796	85.2 (38)
Kewaunee Cty	147	87.1 (22)	0.0 (1)	205	36.1 (25)	72	* *	59	* *
La Crosse Cty	1,449	89.1 (10)	1.3 (8)	1,751	40.8 (15)	539	21.0 (25)	482	63.3 (61)
Marinette	583	87.8 (18)	6.2 (54)	727	42.1 (11)	209	24.4 (20)	202	87.6 (29)
Menominee	271	75.3 (66)	3.3 (28)	357	55.5 (1)	133	8.3 (59)	123	95.1 (2)
Mil. City	4,679	75.9 (64)	5.0 (44)	6,331	32.0 (42)	2,199	14.2 (51)	1,928	80.8 (51)
MLK	1,902	75.6 (65)	3.0 (24)	2,433	30.7 (46)	903	11.0 (55)	759	82.7 (47)
Monitowoc	763	78.2 (58)	6.3 (55)	1,116	43.4 (8)	369	17.6 (39)	356	84.6 (40)
Monroe Cty	547	84.8 (31)	3.1 (25)	769	26.0 (59)	271	20.3 (28)	222	80.2 (54)
NEW Comm	2,876	76.0 (63)	7.4 (62)	3,980	36.9 (22)	1,379	16.9 (41)	1,249	76.5 (55)
NHA	2,815	84.8 (31)	8.3 (65)	3,077	30.2 (48)	1,065	14.8 (49)	885	87.6 (29)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
NWCSA	550	85.5 (27)	3.5 (29)	822	38.1 (20)	266	25.2 (17)	231	89.2 (22)
Oconto Cty	306	87.6 (21)	1.3 (8)	409	35.7 (28)	149	22.8 (23)	123	93.5 (5)
Oneida	297	79.5 (54)	1.7 (15)	397	42.1 (11)	156	6.4 (62)	143	90.2 (14)
Oneida Cty	342	81.6 (46)	7.3 (61)	456	36.0 (26)	139	34.5 (4)	127	91.3 (9)
Outagamie	1,271	78.4 (56)	6.7 (57)	1,902	32.4 (41)	629	24.6 (19)	544	80.9 (50)
Pierce Cty	329	83.9 (33)	6.1 (53)	468	35.5 (29)	148	16.2 (45)	132	87.9 (28)
Polk Cty	469	82.5 (39)	8.1 (63)	681	38.6 (19)	212	26.4 (13)	181	90.6 (11)
Portage Cty	692	85.0 (30)	1.6 (13)	856	36.9 (22)	151	15.9 (47)	240	85.4 (36)
Price Cty	192	82.8 (38)	6.3 (55)	258	50.0 (2)	93	* *	87	* *
Racine	633	77.4 (60)	5.2 (45)	1,052	35.4 (30)	396	7.8 (60)	389	73.0 (58)
Rusk Cty	208	88.0 (17)	3.8 (31)	292	45.9 (5)	112	38.4 (3)	96	* *
Sauk Cty	724	88.8 (13)	2.2 (18)	830	28.3 (55)	298	15.1 (48)	180	90.0 (15)
Sawyer Cty	249	89.2 (8)	2.8 (23)	280	47.5 (4)	93	* *	84	* *
Seeds	1,611	78.0 (59)	4.8 (43)	2,339	34.0 (35)	876	19.2 (34)	766	83.4 (45)
Shawano	401	83.3 (34)	10.0 (67)	544	33.6 (37)	177	31.6 (6)	157	95.5 (1)
Sheboygan	1,058	78.4 (56)	4.2 (34)	1,583	21.7 (69)	573	16.6 (42)	454	87.0 (33)
St. Croix Cty	545	82.0 (42)	8.1 (63)	766	36.0 (26)	258	24.4 (20)	215	91.6 (7)
SWCAP	674	88.1 (16)	2.4 (21)	949	29.4 (51)	290	19.3 (33)	242	89.3 (20)
Taylor Cty	257	89.5 (6)	0.8 (4)	286	33.9 (36)	119	26.1 (16)	106	89.6 (19)
Tremp	408	88.5 (14)	2.5 (22)	483	35.2 (31)	190	16.3 (43)	158	72.8 (59)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 6F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Vernon Cty	315	93.0 (1)	4.4	(38)	405	39.0 (17)	128	42.2 (1)	126	76.2 (56)
Vilas	151	91.4 (4)	1.3	(8)	196	34.7 (32)	63	* *	63	* *
Walworth Cty	970	79.7 (53)	4.6	(41)	1,422	29.0 (54)	446	20.2 (29)	389	81.7 (49)
Wash/Oz	971	88.2 (15)	4.3	(36)	1,386	32.9 (38)	481	19.1 (35)	417	88.2 (26)
Washburn Cty	212	92.9 (2)	0.5	(3)	296	42.9 (9)	105	21.0 (25)	101	91.1 (10)
Waukesha Cty	1,394	82.4 (40)	4.1	(33)	1,864	22.6 (66)	615	17.7 (38)	477	90.6 (11)
Waupaca Cty	437	87.0 (23)	3.2	(26)	607	29.3 (53)	203	8.9 (58)	170	88.2 (26)
Wee Care	4,430	77.1 (61)	4.6	(41)	6,079	30.2 (48)	2,017	9.8 (57)	1,771	80.8 (51)
West Allis	691	81.9 (44)	3.6	(30)	981	22.7 (65)	373	27.6 (11)	277	89.9 (17)
Winnebago	1,324	83.1 (36)	4.3	(36)	1,886	22.6 (66)	690	14.8 (49)	576	84.0 (43)
Wood Cty	787	86.0 (24)	1.3	(8)	1,046	17.9 (70)	387	17.6 (39)	315	85.7 (35)
Unknown Local Agencies	3	* *	* *		3	* *	0	* *	0	* *
<b>Wisconsin</b>	<b>65,810</b>	<b>80.5</b>	<b>5.4</b>		<b>89,103</b>	<b>32.4</b>	<b>30,161</b>	<b>17.7</b>	<b>26,213</b>	<b>83.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
16th Street	3,714	15.1 (1)	4,114	8.3 (1)	448	38.8 (1)	3,287	16.3 (1)	4,062	11.3 (4)
Aurora	1,739	20.5 (2)	1,925	14.5 (4)	263	24.7 (40)	1,455	31.1 (39)	1,855	23.1 (57)
Barron Cty	663	49.0 (54)	770	32.5 (60)	286	23.4 (45)	568	31.0 (37)	728	17.9 (31)
Bay Area	500	48.0 (51)	556	30.2 (49)	208	31.3 (16)	437	27.7 (24)	541	18.3 (34)
Buffalo	212	44.8 (40)	245	28.6 (41)	83	* *	185	17.3 (2)	238	8.8 (1)
Burnett Cty	220	56.8 (68)	259	35.5 (67)	110	21.8 (51)	186	44.1 (68)	231	26.8 (68)
Calumet Cty	320	37.2 (18)	377	27.6 (33)	97	* *	247	26.3 (21)	349	16.9 (19)
Chippewa	777	53.7 (65)	938	32.4 (59)	368	28.0 (29)	660	32.4 (43)	901	17.2 (24)
City Mad	3,091	27.3 (6)	3,361	15.9 (6)	580	31.9 (13)	2,511	19.9 (3)	3,259	11.1 (3)
Clark Cty	432	38.9 (24)	496	26.8 (29)	143	18.2 (58)	353	28.0 (25)	465	17.8 (30)
Columbia Cty	508	46.9 (47)	556	29.7 (45)	184	28.3 (27)	418	35.4 (51)	531	18.5 (35)
Cudahy	1,383	43.0 (36)	1,645	25.1 (24)	461	31.5 (14)	1,100	34.5 (49)	1,567	24.9 (64)
Dane Cty	1,149	29.2 (7)	1,362	20.4 (10)	259	34.0 (10)	966	24.0 (11)	1,281	17.0 (20)
Dodge Cty	758	43.3 (38)	893	28.4 (39)	267	24.0 (43)	628	31.8 (42)	883	19.5 (41)
Door Cty	225	51.1 (61)	277	29.2 (42)	102	27.5 (33)	184	30.4 (33)	265	18.9 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dunn Cty	538	44.8 (40)	598	25.8 (26)	199	28.6 (25)	466	30.0 (28)	588	15.3 (15)
Eau Claire	1,286	45.4 (44)	1,489	28.1 (35)	504	29.8 (21)	1,085	31.0 (37)	1,405	17.4 (25)
Family Health	585	38.3 (23)	686	29.2 (42)	180	18.3 (56)	486	37.2 (58)	666	23.9 (60)
Family Plan	2,082	41.0 (32)	2,375	28.5 (40)	717	19.8 (54)	1,707	30.3 (32)	2,248	19.3 (39)
Florence Cty	53	* *	56	* *	19	* *	42	* *	56	* *
Fond du Lac	1,152	47.5 (50)	1,315	28.2 (38)	464	28.4 (26)	966	23.9 (10)	1,262	11.9 (5)
Forest Cty	145	40.7 (26)	160	28.1 (35)	47	* *	124	29.8 (27)	155	20.6 (49)
GLITC	683	60.0 (69)	735	40.1 (69)	348	24.4 (41)	584	45.5 (70)	698	26.1 (66)
Grant Cty	575	37.4 (20)	715	25.7 (25)	186	25.8 (38)	480	24.0 (11)	695	13.5 (10)
Green Cty	382	53.1 (63)	430	30.5 (51)	173	35.3 (5)	330	41.5 (66)	407	23.8 (59)
Health	2,226	33.9 (11)	2,415	21.9 (13)	591	28.1 (28)	1,861	27.3 (23)	2,409	16.6 (18)
Jackson	330	49.4 (55)	365	33.7 (65)	141	15.6 (59)	284	40.1 (64)	353	21.2 (51)
Jefferson Cty	946	37.4 (20)	1,078	24.6 (22)	296	29.4 (23)	799	25.3 (16)	1,026	17.6 (29)
Juneau	634	40.7 (26)	700	30.9 (53)	218	12.8 (60)	537	39.7 (63)	655	27.0 (69)
Kenosha Cty	2,343	34.7 (16)	2,698	22.5 (15)	673	29.9 (20)	1,928	33.5 (46)	2,504	20.3 (45)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

Comparison of Smoking Indicators (2, 3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee Cty	170	34.1 (13)	187	22.5 (15)	50	* *	143	30.1 (31)	182	17.0 (20)
La Crosse Cty	1,419	37.3 (19)	1,586	23.1 (19)	430	20.7 (53)	1,191	23.6 (9)	1,525	13.7 (12)
Marinette	593	53.1 (63)	680	33.4 (64)	268	26.9 (36)	495	35.6 (52)	627	21.9 (54)
Menominee	309	68.3 (70)	341	42.5 (70)	184	31.0 (17)	264	41.3 (65)	319	20.7 (50)
Mil. City	5,156	22.6 (4)	5,835	14.3 (3)	956	27.8 (30)	4,395	26.9 (22)	5,697	18.1 (33)
MLK	2,024	30.0 (8)	2,194	19.1 (9)	471	27.6 (31)	1,695	36.8 (54)	2,123	24.7 (62)
Monitowoc	917	47.4 (49)	1,031	32.8 (61)	375	23.7 (44)	774	37.0 (56)	986	24.4 (61)
Monroe Cty	641	47.1 (48)	756	31.9 (56)	257	24.9 (39)	540	32.8 (44)	698	19.9 (44)
NEW Comm	3,344	37.6 (22)	3,670	22.2 (14)	1,050	31.5 (14)	2,821	23.4 (8)	3,545	14.4 (13)
NHA	2,392	41.5 (33)	2,780	27.5 (32)	830	22.0 (50)	1,990	33.5 (46)	2,637	19.5 (41)
NWCSA	670	55.4 (66)	745	33.0 (62)	315	27.6 (31)	587	39.4 (62)	721	20.4 (48)
Oconto Cty	325	50.8 (59)	395	38.2 (68)	147	19.0 (55)	278	39.2 (61)	384	22.1 (56)
Oneida	322	49.7 (57)	369	27.1 (31)	137	30.7 (18)	281	43.4 (67)	348	24.7 (62)
Oneida Cty	381	56.4 (67)	414	34.1 (66)	168	29.8 (21)	310	36.5 (53)	381	21.5 (52)
Outagamie	1,501	42.0 (34)	1,707	23.4 (20)	511	33.9 (11)	1,260	26.2 (19)	1,637	15.9 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Pierce Cty	372	51.9 (62)	421	32.1 (58)	157	24.2 (42)	308	37.0 (56)	403	19.4 (40)
Polk Cty	526	48.9 (53)	629	29.9 (47)	221	22.6 (48)	426	37.8 (59)	605	17.9 (31)
Portage Cty	694	36.0 (17)	768	19.0 (7)	214	34.1 (8)	581	24.3 (13)	745	12.8 (8)
Price Cty	226	40.7 (26)	241	27.0 (30)	82	* *	202	30.7 (35)	238	23.5 (58)
Racine	816	30.3 (9)	942	19.0 (7)	210	27.1 (34)	690	25.5 (17)	905	10.1 (2)
Rusk Cty	259	40.9 (30)	271	29.9 (47)	90	* *	229	31.4 (41)	268	19.8 (43)
Sauk Cty	691	44.9 (42)	756	31.6 (55)	254	22.8 (47)	549	45.2 (69)	683	36.6 (70)
Sawyer Cty	241	49.8 (58)	268	24.6 (22)	97	* *	200	30.5 (34)	244	21.7 (53)
Seeds	2,006	26.8 (5)	2,146	14.8 (5)	457	38.7 (2)	1,736	21.4 (4)	2,080	13.2 (9)
Shawano	455	46.4 (46)	526	29.5 (44)	182	23.1 (46)	383	34.7 (50)	507	21.9 (54)
Sheboygan	1,270	34.1 (13)	1,431	22.6 (17)	323	26.6 (37)	1,058	26.2 (19)	1,385	17.1 (22)
St. Croix Cty	620	44.0 (39)	696	26.3 (27)	220	32.7 (12)	504	25.0 (14)	667	17.1 (22)
SWCAP	745	45.8 (45)	861	31.9 (56)	266	22.6 (48)	624	30.9 (36)	836	17.5 (27)
Taylor Cty	258	40.7 (26)	282	30.9 (53)	93	* *	217	36.9 (55)	262	26.7 (67)
Tremp	411	32.8 (10)	456	22.8 (18)	109	18.3 (56)	337	22.6 (6)	431	12.1 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 7F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
Vernon Cty	342	33.9 (11)	381	21.3 (12)	100	30.0 (19)	291	21.6 (5)	370	12.7 (7)
Vilas	165	50.9 (60)	172	30.2 (49)	65	* *	135	31.1 (39)	158	20.3 (45)
Walworth Cty	1,111	34.4 (15)	1,311	20.4 (10)	311	38.3 (3)	920	22.7 (7)	1,270	13.5 (10)
Wash/Oz	1,085	48.8 (52)	1,241	26.3 (27)	433	36.7 (4)	895	28.5 (26)	1,173	15.3 (15)
Washburn Cty	234	42.3 (35)	273	29.7 (45)	93	* *	207	30.0 (28)	258	18.6 (36)
Waukesha Cty	1,476	40.2 (25)	1,710	24.4 (21)	443	34.5 (7)	1,170	25.1 (15)	1,642	15.2 (14)
Waupaca Cty	470	49.4 (55)	571	33.1 (63)	188	21.3 (52)	357	38.7 (60)	512	25.4 (65)
Wee Care	4,887	21.2 (3)	5,693	12.7 (2)	849	35.2 (6)	4,158	25.7 (18)	5,558	17.4 (25)
West Allis	772	45.2 (43)	892	27.6 (33)	276	34.1 (8)	642	34.3 (48)	852	20.3 (45)
Winnebago	1,517	43.1 (37)	1,732	30.7 (52)	529	27.0 (35)	1,258	30.0 (28)	1,664	18.9 (37)
Wood Cty	863	40.9 (30)	960	28.1 (35)	286	28.7 (24)	710	33.0 (45)	926	17.5 (27)
Unknown Local Agencies	3	* *	3	* *	1	* *	3	* *	3	* *
<b>Wisconsin</b>	<b>72,330</b>	<b>36.0</b>	<b>81,912</b>	<b>22.8</b>	<b>21,313</b>	<b>28.2</b>	<b>60,678</b>	<b>28.5</b>	<b>78,738</b>	<b>17.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	<b>Birthweight</b>			<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>		
	<b>Low</b> (4)	<b>High</b> (5)		<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	<b>Number</b>	<b>%</b> (Rank)	
	<b>Number</b>	<b>%</b> (Rank)	<b>%</b> (Rank)							
16th Street	4,259	6.3 (18)	8.9 (42)	4,055	10.5 (41)	3,577	2.5 (27)	4,326	83.6 (1)	
Aurora	1,847	10.9 (68)	5.1 (3)	1,799	13.3 (67)	1,492	4.8 (68)	1,947	50.1 (67)	
Barron Cty	731	7.3 (42)	10.0 (56)	721	11.4 (54)	606	2.8 (35)	761	66.0 (37)	
Bay Area	559	4.7 (5)	13.6 (69)	536	9.0 (11)	483	2.3 (19)	567	69.8 (22)	
Buffalo	243	4.5 (4)	9.5 (47)	229	9.2 (15)	203	1.5 (4)	228	76.3 (4)	
Burnett Cty	247	8.1 (54)	6.9 (10)	240	10.8 (47)	210	3.3 (56)	250	74.4 (9)	
Calumet Cty	382	8.1 (54)	7.9 (24)	364	11.5 (55)	317	1.9 (9)	386	60.1 (54)	
Chippewa	939	6.5 (20)	7.5 (19)	909	10.1 (34)	810	2.2 (15)	951	70.1 (20)	
City Mad	3,409	7.3 (42)	8.0 (26)	3,277	9.3 (22)	2,876	3.0 (47)	3,482	76.0 (5)	
Clark Cty	492	7.3 (42)	8.7 (39)	443	8.1 (4)	403	3.0 (47)	495	69.5 (26)	
Columbia Cty	553	7.1 (36)	11.6 (65)	538	8.9 (10)	482	3.3 (56)	558	69.9 (21)	
Cudahy	1,619	7.1 (36)	9.3 (46)	1,599	10.6 (43)	1,379	2.1 (12)	1,664	64.9 (40)	
Dane Cty	1,404	6.7 (26)	9.7 (51)	1,331	10.1 (34)	1,184	2.4 (24)	1,422	75.0 (7)	
Dodge Cty	903	8.2 (58)	7.1 (11)	888	12.2 (62)	762	2.2 (15)	923	75.1 (6)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Door Cty	275	5.1 (8)	7.6	(20)	267	9.7 (25)	239	2.1 (12)	277	70.4 (19)
Dunn Cty	605	5.3 (11)	10.4	(59)	571	8.8 (9)	513	2.3 (19)	613	67.2 (34)
Eau Claire	1,509	6.2 (17)	7.6	(20)	1,400	9.8 (26)	1,252	2.7 (31)	1,521	68.5 (30)
Family Health	686	6.7 (26)	8.3	(34)	685	10.7 (44)	594	1.9 (9)	705	57.9 (56)
Family Plan	2,343	7.1 (36)	7.4	(16)	2,242	9.2 (15)	1,991	2.9 (40)	2,382	56.5 (62)
Florence Cty	57	* * *	*	*	57	* *	48	* *	60	* *
Fond du Lac	1,303	8.1 (54)	6.4	(8)	1,276	11.8 (59)	1,099	2.9 (40)	1,328	67.7 (32)
Forest Cty	152	8.6 (61)	7.2	(12)	147	13.6 (69)	125	2.4 (24)	158	53.2 (65)
GLITC	722	5.5 (12)	13.3	(68)	679	10.9 (49)	591	2.2 (15)	719	60.5 (53)
Grant Cty	715	6.0 (14)	9.5	(47)	688	8.4 (6)	613	2.3 (19)	729	61.7 (47)
Green Cty	415	6.5 (20)	8.2	(29)	398	10.3 (38)	354	2.5 (27)	418	61.2 (48)
Health	2,412	7.8 (50)	8.2	(29)	2,276	10.2 (37)	2,007	3.3 (56)	2,410	60.9 (50)
Jackson	369	7.6 (47)	10.6	(60)	354	11.6 (56)	309	2.6 (29)	375	73.1 (11)
Jefferson Cty	1,008	4.4 (3)	8.2	(29)	965	9.2 (15)	854	1.6 (5)	1,022	71.4 (15)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Juneau	691	6.7 (26)	8.0 (26)	656	8.2 (5)	596	2.9 (40)	694	65.9 (38)
Kenosha Cty	2,675	8.3 (60)	6.5 (9)	2,564	9.9 (31)	2,266	3.8 (62)	2,718	57.8 (57)
Kewaunee Cty	190	6.3 (18)	8.4 (36)	187	7.0 (2)	169	1.8 (8)	192	70.8 (18)
La Crosse Cty	1,598	6.8 (29)	10.6 (60)	1,531	9.8 (26)	1,353	3.1 (52)	1,613	71.4 (15)
Marinette	658	6.8 (29)	9.7 (51)	632	8.5 (7)	564	3.0 (47)	669	48.7 (70)
Menominee	324	9.0 (64)	11.4 (64)	304	13.2 (66)	258	3.9 (64)	326	68.1 (31)
Mil. City	5,724	10.2 (67)	5.3 (4)	5,468	12.8 (65)	4,634	3.9 (64)	5,953	49.9 (68)
MLK	2,115	12.3 (70)	4.0 (1)	2,048	14.0 (70)	1,689	5.2 (69)	2,208	54.1 (64)
Monitowoc	1,040	7.1 (36)	7.7 (23)	1,012	10.4 (40)	887	2.9 (40)	1,065	57.2 (59)
Monroe Cty	739	6.5 (20)	9.6 (49)	701	9.3 (22)	620	2.4 (24)	756	69.4 (27)
NEW Comm	3,711	7.3 (42)	8.1 (28)	3,573	11.6 (56)	3,089	2.8 (35)	3,780	62.7 (44)
NHA	2,803	7.9 (51)	8.9 (42)	2,745	10.7 (44)	2,395	2.8 (35)	2,855	54.3 (63)
NWCSA	739	4.7 (5)	10.0 (56)	710	11.0 (50)	625	1.4 (3)	743	68.9 (28)
Oconto Cty	396	9.3 (65)	7.3 (14)	377	10.1 (34)	332	6.0 (70)	400	52.3 (66)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Oneida	357	4.8 (7)	19.6 (70)	346	9.2 (15)	307	0.0 (1)	364	61.0 (49)	
Oneida Cty	396	8.1 (54)	5.6 (6)	390	10.8 (47)	345	3.2 (54)	401	60.6 (52)	
Outagamie	1,689	7.7 (48)	7.6 (20)	1,664	9.9 (31)	1,452	2.8 (35)	1,728	63.0 (42)	
Pierce Cty	433	5.1 (8)	8.3 (34)	418	7.4 (3)	378	1.9 (9)	444	62.6 (45)	
Polk Cty	641	7.0 (34)	7.3 (14)	614	9.9 (31)	543	2.9 (40)	655	73.7 (10)	
Portage Cty	768	6.5 (20)	11.1 (63)	742	9.8 (26)	656	3.2 (54)	781	67.2 (34)	
Price Cty	243	3.3 (1)	9.9 (55)	238	6.7 (1)	218	0.9 (2)	247	69.6 (24)	
Racine	944	9.6 (66)	6.0 (7)	898	12.0 (61)	774	3.4 (59)	960	49.5 (69)	
Rusk Cty	271	7.7 (48)	12.2 (66)	257	11.7 (58)	225	2.7 (31)	272	59.6 (55)	
Sauk Cty	742	6.1 (15)	10.2 (58)	741	9.2 (15)	660	2.3 (19)	758	78.2 (2)	
Sawyer Cty	260	6.9 (31)	9.6 (49)	250	9.6 (24)	220	2.7 (31)	263	68.8 (29)	
Seeds	2,190	6.5 (20)	9.0 (44)	2,106	8.5 (7)	1,907	2.9 (40)	2,219	73.1 (11)	
Shawano	520	7.9 (51)	9.8 (53)	493	11.0 (50)	427	3.5 (60)	533	64.2 (41)	
Sheboygan	1,466	5.7 (13)	7.4 (16)	1,406	9.0 (11)	1,268	2.3 (19)	1,480	65.2 (39)	

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
St. Croix Cty	718	6.1 (15)	11.0 (62)	668	10.3 (38)	592	1.7 (6)	730	71.8 (13)
SWCAP	860	7.0 (34)	9.8 (53)	822	9.2 (15)	728	3.7 (61)	881	69.6 (24)
Taylor Cty	273	5.1 (8)	8.8 (41)	272	10.7 (44)	239	1.7 (6)	276	67.0 (36)
Tremp	449	7.1 (36)	8.5 (38)	426	12.4 (63)	361	3.0 (47)	457	71.6 (14)
Vernon Cty	379	6.9 (31)	12.4 (67)	358	9.8 (26)	320	2.2 (15)	384	74.5 (8)
Vilas	167	4.2 (2)	5.4 (5)	164	9.1 (14)	146	2.1 (12)	166	57.2 (59)
Walworth Cty	1,342	7.4 (46)	8.4 (36)	1,263	9.8 (26)	1,116	2.8 (35)	1,379	77.2 (3)
Wash/Oz	1,228	6.9 (31)	8.2 (29)	1,207	12.6 (64)	1,036	2.9 (40)	1,253	70.9 (17)
Washburn Cty	277	8.7 (62)	7.9 (24)	266	10.5 (41)	236	3.8 (62)	278	62.9 (43)
Waukesha Cty	1,727	6.5 (20)	8.2 (29)	1,654	9.2 (15)	1,476	2.6 (29)	1,751	69.8 (22)
Waupaca Cty	529	8.9 (63)	7.2 (12)	539	11.3 (53)	450	4.4 (66)	552	60.7 (51)
Wee Care	5,668	11.3 (69)	4.8 (2)	5,443	13.4 (68)	4,570	4.6 (67)	5,786	57.5 (58)
West Allis	901	8.2 (58)	8.7 (39)	848	11.8 (59)	738	3.1 (52)	913	67.5 (33)
Winnebago	1,755	8.0 (53)	7.4 (16)	1,686	11.2 (52)	1,464	2.7 (31)	1,780	62.0 (46)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.

**Table 8F**  
**2005 - 2007 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Wood Cty	956	7.2 (41)	9.0	(44)	919	9.0 (11)	801	3.0 (47)	972	57.1 (61)
Unknown Local Agencies	3	* * *	*	*	3	* *	2	* *	3	* *
<b>Wisconsin</b>	<b>81,713</b>	<b>7.8</b>	<b>7.9</b>		<b>78,543</b>	<b>10.7</b>	<b>68,505</b>	<b>3.1</b>	<b>83,315</b>	<b>64.2</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

