

Table 3B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Adams Cty	149	87.9	0.0	6.0	0.7	1.3	4.0	0.0
Ashland	150	75.3	0.0	0.7	21.3	0.0	2.7	0.0
Barron	323	84.2	6.8	6.8	0.6	1.2	0.3	0.0
Bayfield C	98	*	*	*	*	*	*	*
Brown	1,770	52.9	6.2	23.6	8.9	6.2	2.2	0.0
Buffalo	74	*	*	*	*	*	*	*
Burnett	124	75.0	0.8	1.6	17.7	0.8	4.0	0.0
Calumet	171	71.9	2.9	14.6	0.0	9.4	1.2	0.0
Chippewa	376	94.9	0.3	2.1	0.3	1.1	1.3	0.0
Clark	201	76.6	0.0	21.4	1.0	0.0	1.0	0.0
Columbia	257	84.8	1.2	10.9	1.6	0.4	1.2	0.0
Crawford C	100	98.0	0.0	1.0	0.0	0.0	1.0	0.0
Dane	2,176	41.5	21.6	27.2	0.5	7.0	2.2	0.0
Dodge	365	80.5	0.3	17.5	0.0	0.5	1.1	0.0
Door	126	85.7	1.6	8.7	0.8	1.6	1.6	0.0
Douglas Ct	353	89.8	0.3	2.8	1.4	1.1	4.5	0.0
Dunn	243	87.2	0.8	4.1	0.0	6.6	1.2	0.0
Eau Claire	636	81.4	0.9	5.0	0.5	9.7	2.4	0.0
Florence	25	*	*	*	*	*	*	*
Fond du La	531	73.4	3.0	16.8	0.2	5.1	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Forest	78	*	*	*	*	*	*	*
Grant Cty	322	92.5	0.6	5.6	0.0	0.3	0.9	0.0
Green Cty	155	86.5	0.0	12.9	0.0	0.6	0.0	0.0
Green Lake	80	*	*	*	*	*	*	*
Iowa Cty	111	93.7	0.9	4.5	0.0	0.0	0.9	0.0
Iron Cty	38	*	*	*	*	*	*	*
Jackson	161	79.5	0.0	3.1	16.1	0.6	0.6	0.0
Jefferson	444	61.7	1.6	34.9	0.2	0.7	0.9	0.0
Juneau	179	93.9	0.6	2.8	0.0	0.0	2.8	0.0
Kenosha	1,248	47.4	17.2	32.4	0.2	1.0	1.8	0.0
Kewaunee	83	*	*	*	*	*	*	*
La Crosse	677	76.8	3.7	2.1	1.0	13.7	2.7	0.0
Lafayette	60	*	*	*	*	*	*	*
Langlade C	126	91.3	0.8	7.1	0.0	0.0	0.8	0.0
Lincoln Ct	127	89.8	0.0	7.1	0.8	0.8	1.6	0.0
Manitowoc	460	72.0	1.3	13.0	1.1	10.4	2.2	0.0
Marathon C	677	65.4	2.1	6.2	0.3	24.5	1.5	0.0
Marinette	254	93.7	0.0	4.7	0.0	0.8	0.8	0.0
Marquette	67	*	*	*	*	*	*	*
Menominee	151	4.6	0.0	6.6	29.1	0.0	59.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Milwaukee	10,502	17.9	49.5	26.4	0.5	4.4	1.3	0.0
Monroe	308	81.5	0.6	15.6	0.3	1.9	0.0	0.0
Oconto	146	94.5	0.7	4.1	0.7	0.0	0.0	0.0
Oneida	185	93.0	0.0	4.9	0.0	0.5	1.6	0.0
Outagamie	754	66.4	2.3	17.0	0.9	11.8	1.6	0.0
Ozaukee	141	86.5	2.1	7.8	0.0	2.1	1.4	0.0
Pepin	34	*	*	*	*	*	*	*
Pierce	181	87.8	0.6	6.1	0.6	1.7	3.3	0.0
Polk	289	92.7	0.0	4.5	0.7	0.7	1.4	0.0
Portage	320	75.6	1.3	10.3	0.0	12.2	0.6	0.0
Price	96	*	*	*	*	*	*	*
Racine Cou	1,582	41.2	27.5	28.5	0.3	0.8	1.8	0.0
Richland C	114	93.0	0.0	7.0	0.0	0.0	0.0	0.0
Rock	1,185	68.8	9.7	18.2	0.2	0.9	2.2	0.0
Rusk	98	*	*	*	*	*	*	*
Sauk	387	82.2	0.3	12.9	3.4	0.0	1.3	0.0
Sawyer	164	56.1	0.0	4.3	37.2	1.2	1.2	0.0
Shawano	227	76.2	0.0	4.8	12.3	0.4	6.2	0.0
Sheboygan	614	59.4	2.4	20.4	0.0	17.4	0.3	0.0
St. Croix	341	88.9	0.3	6.2	0.6	3.5	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Taylor	126	95.2	0.0	2.4	0.0	0.8	1.6	0.0
Trempealea	217	72.4	0.5	25.3	0.0	0.0	1.8	0.0
Vernon	196	89.8	0.5	6.6	0.0	1.5	1.5	0.0
Vilas	135	57.0	0.7	3.0	36.3	0.0	3.0	0.0
Walworth	579	56.5	0.7	39.9	1.2	0.9	0.9	0.0
Washburn	110	94.5	0.0	4.5	0.0	0.9	0.0	0.0
Washington	444	86.3	1.8	9.5	0.2	1.4	0.9	0.0
Waukesha	701	67.5	4.0	24.5	0.6	2.0	1.4	0.0
Waupaca	241	90.5	0.0	8.3	0.0	0.4	0.8	0.0
Waushara C	124	66.9	0.0	32.3	0.0	0.0	0.8	0.0
Winnebago	775	76.4	4.1	10.3	0.3	7.0	1.9	0.0
Wood	487	83.8	0.8	8.6	2.3	3.7	0.8	0.0
Wisconsin	35,549	53.7	19.1	19.2	1.7	4.5	1.8	0.0
Nation (Prior Year)	1,252,728	41.7	23.3	27.9	1.1	3.2	1.0	1.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams Cty	149	0.0	11.4	17.4	57.7	13.4	0.0	0.0
Ashland	150	0.0	4.7	15.3	54.0	22.7	3.3	0.0
Barron	323	0.0	2.8	16.1	62.2	17.3	1.5	0.0
Bayfield C	98	*	*	*	*	*	*	*
Brown	1,770	0.5	6.7	12.8	61.1	17.5	1.4	0.1
Buffalo	74	*	*	*	*	*	*	*
Burnett	124	0.0	4.8	14.5	70.2	10.5	0.0	0.0
Calumet	171	0.0	3.5	13.5	64.9	17.0	1.2	0.0
Chippewa	376	0.0	3.2	13.6	62.2	19.1	1.9	0.0
Clark	201	0.5	3.5	10.9	60.2	23.9	1.0	0.0
Columbia	257	0.0	3.9	13.6	65.8	14.4	2.3	0.0
Crawford C	100	0.0	7.0	15.0	66.0	11.0	1.0	0.0
Dane	2,176	0.3	4.9	10.7	60.6	22.2	1.2	0.0
Dodge	365	0.3	3.0	13.4	65.5	15.9	1.9	0.0
Door	126	0.0	4.0	8.7	65.1	19.0	3.2	0.0
Douglas Ct	353	0.3	5.1	12.5	65.7	15.3	1.1	0.0
Dunn	243	0.0	6.2	14.0	62.1	17.3	0.4	0.0
Eau Claire	636	0.0	3.3	13.5	67.8	14.6	0.8	0.0
Florence	25	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Fond du La	531	0.2	6.4	14.3	62.7	15.4	0.9	0.0
Forest	78	*	*	*	*	*	*	*
Grant Cty	322	0.0	4.7	11.2	64.9	18.0	1.2	0.0
Green Cty	155	0.6	5.2	16.8	64.5	12.3	0.6	0.0
Green Lake	80	*	*	*	*	*	*	*
Iowa Cty	111	0.0	4.5	9.0	66.7	18.9	0.9	0.0
Iron Cty	38	*	*	*	*	*	*	*
Jackson	161	0.0	6.2	16.1	61.5	15.5	0.6	0.0
Jefferson	444	0.5	3.2	14.2	64.6	16.0	1.6	0.0
Juneau	179	0.0	3.9	16.2	64.2	14.5	1.1	0.0
Kenosha	1,248	0.4	7.1	15.5	60.7	15.4	0.9	0.0
Kewaunee	83	*	*	*	*	*	*	*
La Crosse	677	0.0	3.7	12.3	67.2	15.4	1.5	0.0
Lafayette	60	*	*	*	*	*	*	*
Langlade C	126	0.0	3.2	13.5	65.1	18.3	0.0	0.0
Lincoln Ct	127	0.0	6.3	11.8	68.5	13.4	0.0	0.0
Manitowoc	460	0.0	5.7	13.5	63.3	16.7	0.9	0.0
Marathon C	677	0.3	5.5	14.8	61.7	16.2	1.3	0.1
Marinette	254	0.8	4.7	12.6	65.4	15.7	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marquette	67	*	*	*	*	*	*	*
Menominee	151	1.3	13.2	17.2	51.0	17.2	0.0	0.0
Milwaukee	10,502	0.5	8.5	13.8	58.8	17.0	1.2	0.0
Monroe	308	0.0	3.9	16.2	63.6	15.3	1.0	0.0
Oconto	146	0.0	5.5	15.8	60.3	18.5	0.0	0.0
Oneida	185	0.0	3.8	16.8	57.8	20.0	1.6	0.0
Outagamie	754	0.0	5.3	11.3	64.6	17.4	1.5	0.0
Ozaukee	141	0.0	1.4	12.1	63.1	22.0	1.4	0.0
Pepin	34	*	*	*	*	*	*	*
Pierce	181	0.0	3.9	13.8	65.2	16.6	0.6	0.0
Polk	289	0.3	5.2	12.8	66.1	14.5	1.0	0.0
Portage	320	0.0	7.2	12.5	66.6	13.1	0.6	0.0
Price	96	*	*	*	*	*	*	*
Racine Cou	1,582	0.6	7.5	13.8	61.0	16.1	0.9	0.0
Richland C	114	0.0	7.0	15.8	57.0	20.2	0.0	0.0
Rock	1,185	0.7	6.8	14.3	60.0	17.6	0.7	0.0
Rusk	98	*	*	*	*	*	*	*
Sauk	387	0.0	5.2	14.2	65.4	13.4	1.6	0.3
Sawyer	164	0.6	4.3	12.8	65.2	14.6	2.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Shawano	227	0.0	5.7	14.5	63.0	16.3	0.4	0.0
Sheboygan	614	0.0	6.4	12.7	60.7	19.2	0.8	0.2
St. Croix	341	0.3	3.8	10.6	67.4	17.9	0.0	0.0
Taylor	126	0.0	4.0	8.7	67.5	18.3	1.6	0.0
Trempealea	217	0.5	3.2	11.5	64.1	20.3	0.5	0.0
Vernon	196	0.0	2.6	7.1	63.8	26.0	0.5	0.0
Vilas	135	0.0	4.4	11.1	59.3	23.0	2.2	0.0
Walworth	579	0.0	4.0	13.0	62.0	19.7	1.4	0.0
Washburn	110	0.0	3.6	9.1	67.3	19.1	0.9	0.0
Washington	444	0.0	5.9	16.7	55.9	20.9	0.7	0.0
Waukesha	701	0.0	2.6	11.0	66.2	19.3	1.0	0.0
Waupaca	241	0.4	3.3	18.3	61.8	15.4	0.8	0.0
Waushara C	124	0.0	2.4	17.7	63.7	14.5	1.6	0.0
Winnebago	775	0.0	5.3	14.5	65.5	14.2	0.4	0.1
Wood	487	0.4	3.5	16.2	65.3	12.7	1.8	0.0
Wisconsin	35,549	0.3	6.2	13.4	61.6	17.2	1.1	0.0
Nation (Prior Year)	1,252,728	0.4	6.5	12.7	59.9	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	147	6.1 (3)	51.7 (55)	69	* *	* *	17	* *	124	17.7(26)	137	2.9 (13)	137	8.0(53)
Ashland	149	6.7 (5)	46.3 (29)	124	23.4 (27)	48.4 (32)	27	* *	124	16.9(21)	140	8.6 (57)	140	2.9(11)
Barron	317	11.7 (44)	50.8 (53)	281	21.7 (20)	53.7 (45)	48	* *	259	21.6(34)	293	4.4 (40)	293	3.8(22)
Bayfield C	97	* *	* *	80	* *	* *	14	* *	63	* *	84	* *	84	* *
Brown	1,751	10.5 (36)	43.5 (13)	1,072	24.4 (33)	45.4 (20)	274	36.1 (11)	1,548	29.1(51)	1,621	5.4 (49)	1,621	6.5(37)
Buffalo	74	* *	* *	66	* *	* *	13	* *	63	* *	69	* *	69	* *
Burnett	124	10.5 (36)	46.8 (34)	76	* *	* *	15	* *	104	20.2(31)	111	3.6 (26)	111	0.9(1)
Calumet	168	11.9 (46)	43.5 (13)	151	19.2 (8)	43.7 (11)	34	* *	143	22.4(38)	162	8.6 (57)	162	4.9(28)
Chippewa	375	7.5 (12)	49.1 (46)	342	21.6 (19)	50.3 (39)	59	* *	324	22.2(36)	346	2.9 (13)	346	7.8(51)
Clark	201	6.5 (4)	45.8 (25)	179	20.7 (15)	45.8 (22)	40	* *	171	43.3(54)	184	2.2 (11)	184	7.1(45)
Columbia	257	8.9 (19)	46.7 (32)	173	17.9 (3)	48.6 (35)	32	* *	208	11.5 (6)	230	3.0 (18)	230	9.1(54)
Crawford C	99	* *	* *	60	* *	* *	21	* *	76	* *	89	* *	89	* *
Dane	2,144	9.4 (23)	44.7 (17)	1,251	26.0 (40)	42.9 (8)	403	31.5 (8)	1,620	17.2(23)	1,940	5.4 (49)	1,940	4.9(28)
Dodge	362	13.0 (53)	45.3 (22)	327	24.2 (31)	43.7 (11)	70	* *	325	26.5(48)	337	5.0 (46)	337	2.7(9)
Door	126	9.5 (25)	50.0 (50)	113	21.2 (16)	46.9 (30)	24	* *	111	13.5(14)	114	7.0 (54)	114	14.0(58)
Douglas Ct	351	9.1 (20)	47.3 (36)	261	19.9 (10)	52.5 (43)	54	* *	296	28.4(50)	307	7.8 (55)	307	7.2(46)
Dunn	238	6.7 (5)	49.2 (47)	133	18.8 (6)	57.9 (48)	35	* *	160	20.0(30)	209	4.3 (37)	209	6.7(40)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)															
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Eau Claire	634	10.1 (32)	45.4 (23)	54	*	*	80	*	*	551	25.6(46)	596	3.9 (29)	596	3.4(17)
Florence	25	*	*	22	*	*	4	*	*	20	*	23	*	23	*
Fond du La	520	10.2 (34)	46.3 (29)	459	25.5 (37)	44.7 (14)	96	*	*	427	23.7(41)	471	3.4 (23)	471	5.7(33)
Forest	78	*	*	65	*	*	13	*	*	50	*	67	*	67	*
Grant Cty	320	9.7 (27)	49.7 (48)	159	15.7 (1)	50.3 (39)	57	*	*	271	16.2(19)	307	5.5 (52)	307	10.1(57)
Green Cty	155	12.9 (52)	45.2 (20)	134	23.9 (29)	44.8 (15)	25	*	*	126	37.3(53)	134	0.7 (1)	134	3.7(21)
Green Lake	75	*	*	65	*	*	16	*	*	55	*	70	*	70	*
Iowa Cty	111	9.9 (31)	52.3 (56)	89	*	*	21	*	*	93	*	107	4.7 (44)	107	6.5(37)
Iron Cty	36	*	*	27	*	*	3	*	*	21	*	33	*	33	*
Jackson	160	11.9 (46)	47.5 (40)	137	27.7 (46)	46.0 (24)	27	*	*	128	22.7(39)	146	4.1 (31)	146	4.1(24)
Jefferson	438	8.2 (15)	44.7 (17)	226	22.6 (24)	45.1 (18)	85	*	*	375	12.5 (9)	400	3.5 (24)	400	6.8(43)
Juneau	177	11.3 (42)	45.8 (25)	88	*	*	20	*	*	146	13.7(16)	160	1.9 (9)	160	6.9(44)
Kenosha	1,220	9.8 (30)	47.5 (40)	1,121	23.0 (26)	46.1 (26)	222	32.4 (9)	1,047	27.0(49)	1,123	3.2 (20)	1,123	6.1(35)	
Kewaunee	83	*	*	75	*	*	14	*	*	71	*	79	*	79	*
La Crosse	675	10.8 (40)	42.5 (8)	250	26.4 (42)	42.8 (7)	110	22.7 (3)	550	21.8(35)	611	4.1 (31)	611	3.3(15)	
Lafayette	58	*	*	37	*	*	12	*	*	47	*	54	*	54	*
Langlade C	126	11.9 (46)	44.4 (16)	111	30.6 (48)	42.3 (5)	23	*	*	107	24.3(42)	114	3.5 (24)	114	2.6(7)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
							Number	%	%	Number	%	%	Number	%
	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Lincoln Ct	126	7.1 (9)	48.4 (43)	109	22.9 (25)	46.8 (29)	19	* *	110	8.2 (2)	122	4.1 (31)	122	3.3(15)
Manitowoc	459	10.7 (39)	42.9 (12)	147	27.9 (47)	43.5 (10)	51	* *	385	16.9(21)	416	4.6 (43)	416	7.5(49)
Marathon C	674	10.1 (32)	43.5 (13)	554	26.9 (45)	41.0 (3)	106	13.2 (1)	593	10.6 (4)	632	3.3 (22)	632	1.9(5)
Marinette	248	14.9 (58)	33.1 (1)	168	17.9 (3)	57.1 (47)	39	* *	152	12.5 (9)	221	5.0 (46)	221	5.4(30)
Marquette	65	* *	* *	57	* *	* *	15	* *	47	* *	58	* *	58	* *
Menominee	151	6.0 (2)	62.9 (59)	135	20.0 (11)	50.4 (41)	18	* *	140	12.9(12)	141	4.3 (37)	141	6.4(36)
Milwaukee	10,390	8.4 (17)	50.3 (52)	6,950	25.5 (37)	45.3 (19)	1,789	44.7 (13)	8,866	34.9(52)	9,546	4.1 (31)	9,546	6.6(39)
Monroe	297	10.4 (35)	47.1 (35)	0	* *	* *	48	* *	262	17.2(23)	280	2.9 (13)	280	1.1(2)
Oconto	146	12.3 (50)	48.6 (44)	139	20.1 (12)	53.2 (44)	33	* *	77	* *	141	2.1 (10)	141	3.5(19)
Oneida	185	14.6 (57)	40.5 (4)	81	* *	* *	27	* *	159	11.3 (5)	173	1.2 (3)	173	5.8(34)
Outagamie	738	11.1 (41)	42.3 (7)	612	26.3 (41)	43.0 (9)	128	28.1 (7)	638	26.2(47)	680	5.4 (49)	680	7.9(52)
Ozaukee	138	13.0 (53)	47.8 (42)	122	18.0 (5)	48.4 (32)	24	* *	124	10.5 (3)	125	3.2 (20)	125	7.2(46)
Pepin	34	* *	* *	31	* *	* *	2	* *	29	* *	32	* *	32	* *
Pierce	180	9.4 (23)	42.8 (11)	128	25.8 (39)	48.4 (32)	16	* *	141	18.4(28)	157	4.5 (41)	157	9.6(56)
Polk	286	11.5 (43)	45.5 (24)	183	19.7 (9)	48.6 (35)	37	* *	238	24.4(43)	264	4.5 (41)	264	9.5(55)
Portage	316	13.9 (56)	38.0 (2)	125	22.4 (23)	46.4 (27)	52	* *	270	12.2 (7)	292	2.7 (12)	292	5.5(32)
Price	95	* *	* *	83	* *	* *	12	* *	79	* *	83	* *	83	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Racine Cou	1,567	9.3 (22)	49.8 (49)	1,369	25.3 (35)	45.6 (21)	286	35.3 (10)	1,348	24.6(44)	1,430	3.8 (28)	1,430	6.7(40)
Richland C	114	13.2 (55)	47.4 (38)	86	* *	* *	16	* *	84	* *	105	2.9 (13)	105	7.6(50)
Rock	1,154	9.6 (26)	47.3 (36)	258	23.6 (28)	36.8 (1)	226	37.2 (12)	990	25.2(45)	1,056	1.6 (5)	1,056	2.6(7)
Rusk	98	* *	* *	87	* *	* *	12	* *	77	* *	89	* *	89	* *
Sauk	379	12.1 (49)	45.9 (27)	339	24.2 (31)	44.5 (13)	63	* *	325	13.5(14)	348	2.9 (13)	348	4.0(23)
Sawyer	164	15.9 (59)	40.2 (3)	128	20.3 (13)	41.4 (4)	20	* *	115	18.3(27)	141	0.7 (1)	141	2.1(6)
Shawano	226	12.4 (51)	42.5 (8)	176	25.0 (34)	44.9 (17)	33	* *	101	20.8(33)	208	4.8 (45)	208	2.9(11)
Sheboygan	613	9.1 (20)	45.2 (20)	530	26.4 (42)	38.5 (2)	129	20.2 (2)	552	16.5(20)	571	4.2 (35)	571	3.0(13)
St. Croix	329	8.8 (18)	46.5 (31)	288	19.1 (7)	51.7 (42)	51	* *	222	6.8 (1)	284	5.6 (53)	284	7.4(48)
Taylor	126	7.9 (14)	56.3 (58)	117	20.5 (14)	48.7 (37)	19	* *	110	17.3(25)	118	4.2 (35)	118	3.4(17)
Trempealea	216	7.4 (11)	50.9 (54)	67	* *	* *	31	* *	178	20.2(31)	197	3.0 (18)	197	3.0(13)
Vernon	196	8.2 (15)	50.0 (50)	105	16.2 (2)	56.2 (46)	29	* *	164	12.2 (7)	185	4.3 (37)	185	1.6(4)
Vilas	135	6.7 (5)	48.9 (45)	82	* *	* *	22	* *	101	15.8(18)	122	1.6 (5)	122	4.1(24)
Walworth	572	7.7 (13)	47.4 (38)	506	21.5 (18)	49.6 (38)	90	* *	487	13.3(13)	526	1.5 (4)	526	1.5(3)
Washburn	110	5.5 (1)	54.5 (57)	86	* *	* *	10	* *	93	* *	96	* *	96	* *
Washington	440	11.8 (45)	40.9 (5)	391	22.3 (22)	45.8 (22)	60	* *	400	12.8(11)	416	1.7 (7)	416	6.7(40)
Waukesha	696	10.5 (36)	40.9 (5)	625	21.4 (17)	47.8 (31)	138	23.9 (4)	546	15.0(17)	647	3.6 (26)	647	5.4(30)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Waupaca	239	7.1 (9)	42.7 (10)	213	25.4 (36)	46.0 (24)	38	* *	187	23.5(40)	217	5.1 (48)	217	4.1(24)
Waushara C	118	6.8 (8)	44.9 (19)	105	21.9 (21)	44.8 (15)	23	* *	97	* *	107	8.4 (56)	107	2.8(10)
Winnebago	765	9.7 (27)	46.7 (32)	600	24.0 (30)	46.5 (28)	139	24.5 (5)	684	18.6(29)	708	4.0 (30)	708	4.1(24)
Wood	485	9.7 (27)	46.2 (28)	430	26.7 (44)	42.3 (5)	124	27.4 (6)	382	22.3(37)	448	1.8 (8)	448	3.6(20)
Wisconsin	35,141	9.5	47.2	24,089	24.2	45.8	5,983	32.5	29,357	24.8	32,250	4.0	32,250	5.6
Nation (Prior Year)	1,175,574	10.9	44.5	922,658	25.0	42.8	180,176	33.5	414,735	29.8	230,547	3.9	280,574	4.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	139	87.1 (38)	2.2 (34)	143	42.0 (11)	144	25.7 (12)	128	87.5 (14)
Ashland	139	89.2 (24)	1.4 (24)	145	34.5 (28)	147	27.2 (8)	124	86.3 (26)
Barron	302	91.7 (11)	1.3 (22)	319	45.1 (8)	314	23.6 (15)	271	83.4 (35)
Bayfield C	87	* *	* *	95	* *	95	* *	83	* *
Brown	1,608	81.3 (53)	2.9 (46)	1,689	35.3 (27)	1,687	18.0 (34)	1,448	79.0 (47)
Buffalo	70	* *	* *	73	* *	74	* *	55	* *
Burnett	111	89.2 (24)	1.8 (29)	117	51.3 (3)	122	13.9 (49)	111	82.0 (39)
Calumet	164	87.2 (37)	2.4 (38)	166	28.9 (44)	169	28.4 (4)	143	83.9 (34)
Chippewa	348	92.0 (10)	0.6 (9)	370	38.6 (18)	369	23.8 (14)	308	80.8 (40)
Clark	186	88.7 (29)	1.1 (19)	190	24.2 (56)	201	30.3 (2)	168	87.5 (14)
Columbia	229	83.0 (49)	1.3 (22)	250	37.2 (24)	252	39.3 (1)	202	89.6 (9)
Crawford C	89	* *	* *	96	* *	96	* *	67	* *
Dane	1,946	85.4 (45)	1.6 (25)	2,099	26.7 (50)	2,040	15.9 (43)	1,685	82.6 (37)
Dodge	343	91.5 (12)	0.6 (9)	354	28.5 (45)	360	20.6 (22)	276	85.1 (32)
Door	112	89.3 (23)	0.0 (1)	124	32.3 (33)	125	28.0 (5)	105	92.4 (3)
Douglas Ct	314	87.9 (33)	1.0 (18)	349	45.6 (7)	343	21.6 (20)	302	88.1 (11)
Dunn	213	87.3 (36)	0.0 (1)	240	39.6 (14)	237	19.8 (23)	191	87.4 (17)
Eau Claire	598	86.3 (41)	0.8 (14)	622	38.6 (18)	632	16.0 (42)	522	85.8 (28)
Florence	23	* *	* *	25	* *	24	* *	19	* *
Fond du La	479	88.5 (30)	1.9 (30)	522	40.2 (12)	517	15.7 (46)	438	79.9 (44)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregpregnancy (4)		During Pregpregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Forest	67	* *	* *	* *	77	* *	74	* *	65	* *
Grant Cty	306	89.9 (20)	2.0 (31)		311	29.6 (43)	316	19.6 (24)	245	87.3 (19)
Green Cty	136	89.7 (22)	2.2 (34)		153	39.2 (15)	155	16.1 (41)	138	90.6 (6)
Green Lake	73	* *	* *	* *	78	* *	75	* *	53	* *
Iowa Cty	107	96.3 (2)	0.9 (15)		110	30.0 (41)	109	26.6 (10)	75	* *
Iron Cty	34	* *	* *	* *	35	* *	36	* *	30	* *
Jackson	147	93.9 (5)	2.7 (43)		150	38.7 (16)	160	19.4 (27)	132	72.0 (51)
Jefferson	391	85.4 (45)	3.8 (52)		432	30.8 (39)	391	15.9 (43)	367	86.4 (24)
Juneau	162	94.4 (3)	2.5 (39)		175	44.6 (9)	174	23.0 (18)	147	85.7 (30)
Kenosha	1,151	77.4 (57)	3.6 (50)		1,217	31.5 (35)	1,190	16.8 (37)	913	79.7 (45)
Kewaunee	78	* *	* *	* *	80	* *	83	* *	61	* *
La Crosse	611	93.3 (6)	0.7 (12)		654	38.4 (20)	662	18.7 (30)	535	50.5 (53)
Lafayette	53	* *	* *	* *	60	* *	56	* *	39	* *
Langlade C	113	92.9 (8)	2.7 (43)		123	24.4 (54)	125	17.6 (36)	109	91.7 (4)
Lincoln Ct	119	90.8 (15)	4.2 (55)		124	37.1 (25)	126	13.5 (50)	102	93.1 (2)
Manitowoc	407	82.3 (51)	3.9 (53)		450	48.9 (4)	455	18.9 (28)	385	82.6 (37)
Marathon C	634	82.2 (52)	6.3 (58)		663	34.1 (29)	640	13.0 (51)	551	85.3 (31)
Marinette	228	93.0 (7)	2.6 (40)		243	38.7 (16)	243	23.5 (16)	204	87.3 (19)
Marquette	61	* *	* *	* *	65	* *	63	* *	45	* *
Menominee	144	76.4 (58)	0.7 (12)		144	51.4 (2)	151	7.3 (58)	139	89.9 (8)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Milwaukee	9,695	78.8 (56)	2.6 (40)	10,130	32.9 (31)	10,284	11.9 (55)	8,419	79.5 (46)	
Monroe	285	88.1 (32)	1.1 (19)	299	27.1 (49)	288	17.7 (35)	230	77.4 (50)	
Oconto	141	92.2 (9)	2.1 (32)	142	27.5 (48)	145	26.2 (11)	117	88.0 (12)	
Oneida	176	90.3 (17)	2.8 (45)	177	44.1 (10)	175	18.3 (33)	154	89.6 (9)	
Outagamie	618	79.1 (55)	2.6 (40)	738	27.8 (47)	733	19.6 (24)	569	80.8 (40)	
Ozaukee	129	89.9 (20)	0.0 (1)	136	24.3 (55)	136	23.5 (16)	92	* *	
Pepin	31	* *	* *	32	* *	32	* *	24	* *	
Pierce	162	90.1 (18)	0.0 (1)	174	37.9 (23)	171	27.5 (6)	142	80.3 (43)	
Polk	259	85.7 (44)	5.4 (57)	285	48.1 (5)	282	18.8 (29)	241	85.9 (27)	
Portage	294	89.1 (26)	1.7 (26)	313	31.0 (37)	192	15.1 (47)	250	83.2 (36)	
Price	84	* *	* *	95	* *	95	* *	83	* *	
Racine Cou	1,426	81.0 (54)	2.1 (32)	1,542	35.9 (26)	1,554	12.9 (52)	1,302	78.1 (48)	
Richland C	105	91.4 (13)	0.0 (1)	110	28.2 (46)	109	22.0 (19)	90	* *	
Rock	1,082	88.5 (30)	4.9 (56)	1,159	29.7 (42)	1,157	10.6 (56)	954	86.9 (22)	
Rusk	89	* *	* *	93	* *	98	* *	81	* *	
Sauk	353	85.8 (43)	2.3 (36)	364	32.1 (34)	379	12.7 (53)	305	87.5 (14)	
Sawyer	143	90.9 (14)	3.5 (48)	160	47.5 (6)	156	14.1 (48)	141	85.1 (32)	
Shawano	204	87.7 (35)	3.9 (53)	214	40.2 (12)	220	24.5 (13)	191	93.2 (1)	
Sheboygan	574	82.8 (50)	1.7 (26)	605	18.0 (59)	608	16.3 (40)	475	86.5 (23)	
St. Croix	302	90.7 (16)	1.7 (26)	334	32.6 (32)	327	27.2 (8)	242	90.5 (7)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	118	94.1 (4)	0.0	(1)	124	30.6 (40)	126	30.2 (3)	99	* *
Trempealea	198	88.9 (28)	0.5	(8)	203	31.5 (35)	210	12.4 (54)	176	67.0 (52)
Vernon	166	96.4 (1)	0.6	(9)	186	38.2 (22)	154	5.8 (59)	152	77.6 (49)
Vilas	115	83.5 (48)	3.5	(48)	129	31.0 (37)	126	16.7 (38)	107	90.7 (5)
Walworth	531	86.1 (42)	0.9	(15)	560	26.1 (51)	565	20.9 (21)	428	86.4 (24)
Washburn	98	* *	* *		107	55.1 (1)	110	27.3 (7)	87	* *
Washington	416	90.1 (18)	3.1	(47)	433	38.3 (21)	442	16.5 (39)	334	87.4 (17)
Waukesha	650	86.6 (40)	0.3	(7)	687	23.9 (57)	690	18.6 (31)	498	87.8 (13)
Waupaca	215	87.9 (33)	2.3	(36)	231	33.3 (30)	232	9.5 (57)	183	85.8 (28)
Waushara C	112	84.8 (47)	0.9	(15)	121	24.8 (53)	118	19.5 (26)	94	* *
Winnebago	699	86.7 (39)	3.6	(50)	754	24.9 (52)	759	15.9 (43)	598	80.4 (42)
Wood	445	89.0 (27)	1.1	(19)	464	20.3 (58)	481	18.5 (32)	380	87.1 (21)
Wisconsin	32,437	84.0	2.3		34,433	33.0	34,386	16.3	28,219	82.0
Nation (Prior Year)	612,462	79.4	4.7		1,034,046	31.3	98,170	15.7	238,509	75.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	146	55.5 (52)	137	32.8 (47)	64	* *	132	38.6 (53)	137	21.9 (58)
Ashland	147	52.4 (47)	139	32.4 (45)	68	* *	127	26.0 (37)	140	5.7 (30)
Barron	312	45.2 (28)	293	28.0 (32)	120	20.0 (26)	275	21.5 (25)	293	5.1 (20)
Bayfield C	94	* *	83	* *	36	* *	85	* *	83	* *
Brown	1,692	39.4 (19)	1,612	21.0 (17)	517	32.9 (9)	1,462	20.7 (21)	1,613	9.8 (51)
Buffalo	70	* *	69	* *	21	* *	56	* *	69	* *
Burnett	120	58.3 (57)	111	46.8 (58)	55	* *	112	35.7 (50)	111	3.6 (6)
Calumet	165	38.2 (15)	162	21.0 (17)	51	* *	142	16.9 (8)	162	4.3 (12)
Chippewa	368	55.7 (53)	346	32.4 (45)	164	35.4 (4)	310	23.5 (34)	346	4.3 (12)
Clark	200	38.0 (14)	184	19.6 (12)	55	* *	170	18.8 (14)	184	5.4 (26)
Columbia	251	57.0 (54)	227	28.2 (34)	100	21.0 (24)	208	23.6 (35)	228	7.5 (40)
Crawford C	94	* *	89	* *	27	* *	69	* *	89	* *
Dane	2,097	28.2 (2)	1,933	15.6 (4)	402	33.8 (7)	1,720	14.1 (1)	1,934	5.3 (22)
Dodge	357	47.3 (35)	337	30.6 (40)	125	28.8 (14)	281	19.9 (18)	337	6.8 (37)
Door	124	39.5 (20)	114	15.8 (5)	40	* *	101	15.8 (4)	114	4.4 (15)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Douglas Ct	335	57.0 (54)	307	33.6 (48)	159	23.9 (21)	306	27.8 (43)	307	3.9 (8)
Dunn	235	46.4 (32)	209	24.4 (24)	76	* *	194	17.5 (11)	209	4.3 (12)
Eau Claire	615	47.6 (37)	596	27.2 (30)	239	30.1 (13)	531	18.8 (14)	596	2.9 (2)
Florence	22	* *	23	* *	9	* *	19	* *	23	* *
Fond du La	515	44.1 (26)	470	21.9 (19)	173	26.6 (18)	449	21.2 (24)	471	5.1 (20)
Forest	76	* *	67	* *	28	* *	66	* *	67	* *
Grant Cty	311	34.4 (7)	307	19.9 (14)	88	* *	250	18.4 (12)	307	5.5 (28)
Green Cty	153	53.6 (51)	134	34.3 (51)	64	* *	139	21.6 (27)	134	9.0 (47)
Green Lake	71	* *	70	* *	15	* *	53	* *	70	* *
Iowa Cty	107	53.3 (50)	107	37.4 (55)	46	* *	77	* *	107	4.7 (17)
Iron Cty	35	* *	33	* *	18	* *	29	* *	33	* *
Jackson	157	57.3 (56)	144	35.4 (54)	72	* *	137	27.7 (42)	146	3.4 (5)
Jefferson	433	34.4 (7)	399	19.0 (10)	109	31.2 (11)	369	15.4 (3)	399	4.8 (18)
Juneau	175	52.0 (45)	159	37.7 (56)	71	* *	150	36.7 (52)	160	15.0 (56)
Kenosha	1,113	33.8 (3)	1,123	19.6 (12)	285	34.7 (5)	920	22.6 (31)	1,122	7.3 (39)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee	83	* *	79	* *	29	* *	66	* *	79	* *
La Crosse	656	38.3 (16)	603	19.1 (11)	201	27.9 (16)	562	17.3 (10)	603	2.0 (1)
Lafayette	55	* *	53	* *	18	* *	39	* *	53	* *
Langlade C	123	50.4 (40)	113	33.6 (48)	50	* *	109	32.1 (47)	114	10.5 (54)
Lincoln Ct	126	50.8 (43)	121	30.6 (40)	48	* *	103	34.0 (48)	122	6.6 (35)
Manitowoc	450	45.3 (30)	416	29.1 (37)	159	20.8 (25)	389	22.4 (29)	416	5.5 (28)
Marathon C	661	36.9 (12)	632	25.2 (26)	192	21.9 (23)	556	21.0 (23)	632	6.0 (31)
Marinette	238	52.1 (46)	221	34.4 (52)	102	24.5 (20)	209	29.2 (45)	221	8.1 (45)
Marquette	64	* *	58	* *	21	* *	46	* *	58	* *
Menominee	149	65.8 (59)	141	28.4 (36)	87	* *	140	26.4 (38)	141	5.0 (19)
Milwaukee	9,931	24.4 (1)	9,543	12.7 (1)	1,812	35.9 (3)	8,531	22.3 (28)	9,542	10.8 (55)
Monroe	283	40.6 (21)	280	30.7 (42)	98	* *	238	18.5 (13)	280	6.4 (34)
Oconto	142	50.7 (42)	141	34.8 (53)	63	* *	117	26.5 (39)	141	7.8 (43)
Oneida	182	63.7 (58)	171	39.2 (57)	92	* *	154	34.4 (49)	172	4.1 (10)
Outagamie	707	42.9 (25)	680	22.2 (20)	221	23.5 (22)	572	14.2 (2)	680	3.8 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ozaukee	136	39.0 (18)	125	14.4 (2)	33	* *	93	* *	125	3.2 (4)
Pepin	33	* *	31	* *	8	* *	24	* *	31	* *
Pierce	166	41.0 (24)	156	22.4 (21)	53	* *	143	28.0 (44)	156	9.0 (47)
Polk	280	52.9 (49)	264	31.1 (43)	125	24.8 (19)	244	26.6 (40)	264	5.3 (22)
Portage	308	34.7 (9)	290	16.9 (6)	84	* *	254	16.5 (6)	290	4.1 (10)
Price	93	* *	83	* *	34	* *	88	* *	83	* *
Racine Cou	1,516	34.3 (6)	1,428	17.4 (7)	411	31.9 (10)	1,311	23.4 (33)	1,428	7.6 (42)
Richland C	112	47.3 (35)	104	26.9 (29)	42	* *	92	* *	104	2.9 (2)
Rock	1,133	40.6 (21)	1,056	25.4 (28)	358	30.4 (12)	960	24.2 (36)	1,056	10.4 (53)
Rusk	96	* *	89	* *	37	* *	82	* *	89	* *
Sauk	376	46.5 (33)	347	34.0 (50)	129	19.4 (27)	310	22.6 (31)	348	6.9 (38)
Sawyer	158	51.9 (44)	141	29.1 (37)	70	* *	142	29.6 (46)	141	9.9 (52)
Shawano	222	52.7 (48)	208	31.3 (44)	98	* *	193	20.2 (19)	208	6.3 (33)
Sheboygan	599	35.4 (10)	571	20.3 (15)	163	27.6 (17)	484	16.5 (6)	571	7.5 (40)
St. Croix	321	45.2 (28)	284	20.4 (16)	99	* *	253	15.8 (4)	284	3.9 (8)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	126	48.4 (38)	117	28.2 (34)	44	* *	99	* *	118	8.5 (46)
Trempealea	205	34.1 (4)	197	14.7 (3)	60	* *	181	18.8 (14)	197	4.6 (16)
Vernon	191	40.8 (23)	184	29.3 (39)	65	* *	161	22.4 (29)	185	5.4 (26)
Vilas	123	47.2 (34)	119	25.2 (26)	43	* *	104	26.9 (41)	119	6.7 (36)
Walworth	551	36.5 (11)	526	18.1 (8)	147	42.9 (1)	431	16.9 (8)	526	5.3 (22)
Washburn	106	38.7 (17)	96	* *	31	* *	91	* *	96	* *
Washington	435	49.2 (39)	416	24.8 (25)	159	34.6 (6)	339	21.5 (25)	416	9.1 (49)
Waukesha	679	44.6 (27)	647	18.1 (8)	207	37.7 (2)	505	20.4 (20)	647	5.3 (22)
Waupaca	235	46.0 (31)	217	27.6 (31)	79	* *	187	35.8 (51)	217	16.6 (57)
Waushara C	111	34.2 (5)	107	22.4 (21)	27	* *	96	* *	107	9.3 (50)
Winnebago	746	50.4 (40)	703	28.0 (32)	269	33.1 (8)	600	19.3 (17)	705	6.1 (32)
Wood	472	36.9 (12)	448	22.5 (23)	128	28.1 (15)	382	20.7 (21)	448	7.8 (43)
Wisconsin	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
Nation (Prior Year)	895,227	26.6	716,629	15.8	161,864	35.1	572,120	20.9	486,636	22.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	Number	% (Rank)						
Adams Cty	131	11.5 (58)	15.3 (58)		128	12.5 (52)	109	3.7 (44)	132	63.6 (44)
Ashland	133	2.3 (1)	13.5 (51)		135	7.4 (9)	120	0.0 (1)	138	73.2 (16)
Barron	286	7.7 (38)	9.8 (40)		294	10.2 (34)	254	2.4 (22)	293	73.0 (17)
Bayfield C	77	*	*	*	79	*	*	*	79	*
Brown	1,543	6.9 (27)	9.2 (32)		1,539	11.4 (47)	1,307	2.4 (22)	1,564	61.3 (48)
Buffalo	68	*	*	*	68	*	*	*	68	*
Burnett	106	4.7 (8)	9.4 (33)		105	10.5 (35)	90	*	108	70.4 (25)
Calumet	156	8.3 (44)	7.7 (16)		155	6.5 (4)	142	2.8 (32)	159	66.7 (38)
Chippewa	334	7.5 (36)	8.1 (19)		341	10.0 (31)	296	2.4 (22)	337	64.7 (41)
Clark	177	7.3 (34)	9.6 (37)		170	11.2 (43)	149	4.0 (47)	178	79.2 (4)
Columbia	215	7.0 (30)	10.7 (43)		221	7.7 (13)	195	5.1 (50)	222	71.6 (20)
Crawford C	88	*	*	*	86	*	*	*	89	*
Dane	1,846	6.4 (25)	9.4 (33)		1,877	9.7 (28)	1,609	2.8 (32)	1,882	78.7 (6)
Dodge	322	6.2 (21)	6.5 (4)		327	8.3 (15)	291	2.1 (18)	328	78.0 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Door	111	5.4 (11)	11.7 (46)		111	15.3 (58)	93	* *	110	70.9 (24)
Douglas Ct	303	3.3 (3)	13.2 (50)		308	9.7 (28)	274	0.4 (4)	305	68.9 (32)
Dunn	206	5.3 (10)	9.7 (39)		212	7.5 (11)	187	2.1 (18)	211	76.8 (10)
Eau Claire	582	5.8 (16)	9.6 (37)		581	6.7 (5)	530	2.6 (27)	588	71.1 (22)
Florence	23	* *	* *		23	* *	21	* *	23	* *
Fond du La	445	7.6 (37)	8.8 (26)		465	10.5 (35)	396	3.5 (43)	455	68.4 (35)
Forest	61	* *	* *		66	* *	51	* *	62	* *
Grant Cty	286	5.9 (18)	12.2 (47)		294	10.9 (39)	245	2.0 (16)	297	63.3 (45)
Green Cty	130	9.2 (53)	14.6 (56)		131	12.2 (49)	113	2.7 (30)	133	60.2 (53)
Green Lake	69	* *	* *		71	* *	61	* *	71	* *
Iowa Cty	105	9.5 (56)	6.7 (6)		106	5.7 (2)	98	* *	105	69.5 (30)
Iron Cty	31	* *	* *		30	* *	25	* *	32	* *
Jackson	138	5.8 (16)	15.2 (57)		132	8.3 (15)	116	0.9 (6)	140	70.0 (29)
Jefferson	380	5.5 (13)	8.2 (22)		390	11.3 (44)	330	1.2 (7)	388	70.4 (25)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Juneau	158	5.7 (15)	7.6 (15)	159	11.3 (44)	137	1.5 (9)	159	72.3 (18)
Kenosha	1,093	9.3 (54)	6.5 (4)	1,113	11.0 (42)	956	3.7 (44)	1,102	65.3 (40)
Kewaunee	77	* * *	* *	75	* *	72	* *	77	* *
La Crosse	572	7.7 (38)	8.4 (24)	584	9.2 (26)	502	3.2 (39)	580	71.4 (21)
Lafayette	54	* * *	* *	55	* *	50	* *	54	* *
Langlade C	107	8.4 (45)	8.4 (24)	111	9.0 (24)	95	* *	109	61.5 (47)
Lincoln Ct	116	3.4 (4)	9.5 (36)	120	5.0 (1)	108	1.9 (14)	119	56.3 (55)
Manitowoc	406	9.4 (55)	6.7 (6)	411	12.7 (54)	349	5.7 (51)	412	61.9 (46)
Marathon C	606	6.9 (27)	5.1 (1)	622	10.1 (33)	537	2.6 (27)	618	60.7 (52)
Marinette	222	7.2 (33)	8.1 (19)	215	8.8 (22)	192	2.6 (27)	222	52.7 (57)
Marquette	58	* * *	* *	58	* *	51	* *	59	* *
Menominee	140	2.9 (2)	14.3 (55)	134	7.5 (11)	123	0.0 (1)	140	68.6 (34)
Milwaukee	9,159	9.0 (51)	6.1 (3)	9,235	10.7 (38)	7,898	3.7 (44)	9,272	64.6 (42)
Monroe	270	7.4 (35)	10.0 (41)	282	8.9 (23)	240	4.2 (49)	280	70.4 (25)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Oconto	136	8.1 (42)	8.8 (26)	137	10.9 (39)	120	1.7 (10)	138	60.9 (51)
Oneida	166	8.4 (45)	9.0 (31)	163	8.0 (14)	146	3.4 (42)	167	61.1 (49)
Outagamie	658	8.7 (48)	6.7 (6)	674	11.3 (44)	573	3.0 (36)	674	68.7 (33)
Ozaukee	125	8.8 (49)	13.6 (52)	125	13.6 (57)	104	0.0 (1)	126	79.4 (3)
Pepin	30	* * *	* *	29	* *	24	* *	30	* *
Pierce	155	6.5 (26)	10.3 (42)	155	13.5 (56)	129	0.8 (5)	159	75.5 (11)
Polk	256	5.5 (13)	5.9 (2)	262	7.3 (6)	235	2.1 (18)	260	78.8 (5)
Portage	288	6.9 (27)	6.9 (11)	290	9.0 (24)	257	3.1 (38)	292	72.3 (18)
Price	79	* * *	* *	81	* *	67	* *	82	* *
Racine Cou	1,388	9.9 (57)	6.8 (9)	1,393	12.3 (50)	1,186	4.1 (48)	1,401	61.0 (50)
Richland C	101	4.0 (6)	13.9 (53)	100	6.0 (3)	91	* *	103	81.6 (1)
Rock	1,040	6.2 (21)	8.1 (19)	1,054	8.5 (17)	927	2.7 (30)	1,056	50.6 (58)
Rusk	84	* * *	* *	82	* *	72	* *	85	* *
Sauk	339	4.1 (7)	8.8 (26)	327	8.6 (19)	293	1.7 (10)	343	78.1 (7)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Sawyer	136	8.1 (42)	8.8 (26)	137	11.7 (48)	117	1.7 (10)	138	73.9 (14)
Shawano	206	6.3 (24)	7.8 (17)	198	12.6 (53)	171	1.2 (7)	206	67.5 (36)
Sheboygan	546	7.7 (38)	7.0 (13)	558	10.0 (31)	487	3.3 (40)	552	66.1 (39)
St. Croix	281	7.1 (31)	8.2 (22)	288	7.3 (6)	253	2.8 (32)	291	77.3 (9)
Taylor	112	7.1 (31)	12.5 (48)	115	8.7 (21)	100	3.0 (36)	114	70.2 (28)
Trempealea	184	6.0 (20)	11.4 (45)	186	8.6 (19)	159	2.5 (26)	190	71.1 (22)
Vernon	177	6.2 (21)	11.3 (44)	176	7.4 (9)	156	1.9 (14)	180	73.9 (14)
Vilas	110	9.1 (52)	12.7 (49)	113	13.3 (55)	91	* *	111	58.6 (54)
Walworth	514	4.9 (9)	8.9 (30)	518	7.3 (6)	461	2.4 (22)	522	79.7 (2)
Washburn	92	* *	* *	92	* *	77	* *	93	* *
Washington	393	5.9 (18)	6.9 (11)	403	9.9 (30)	348	2.0 (16)	406	74.1 (13)
Waukesha	633	5.4 (11)	7.4 (14)	638	8.5 (17)	567	2.1 (18)	642	74.6 (12)
Waupaca	203	8.9 (50)	9.4 (33)	209	12.4 (51)	170	1.8 (13)	208	64.4 (43)
Waushara C	106	3.8 (5)	14.2 (54)	110	10.9 (39)	92	* *	110	69.1 (31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Winnebago	690	8.4 (45)	6.8 (9)	691	10.6 (37)	600	3.3 (40)	697	67.0 (37)
Wood	415	8.0 (41)	8.0 (18)	421	9.5 (27)	357	2.8 (32)	423	56.0 (56)
Wisconsin	31,033	7.7	7.8	31,344	10.2	27,048	3.1	31,499	67.0
Nation (Prior Year)	1,039,036	8.2	6.7	947,705	11.9	856,467	3.7	936,194	64.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Adams Cty	309	89.3	0.0	5.2	0.6	0.6	4.2	0.0
Ashland	463	72.6	0.0	0.9	22.2	0.6	2.8	0.9
Barron	942	86.4	5.3	5.9	0.4	0.5	0.4	1.0
Bayfield C	268	63.8	0.0	4.1	27.6	1.1	3.0	0.4
Brown	5,017	52.2	4.6	25.0	9.5	6.6	1.6	0.5
Buffalo	204	92.6	0.0	5.4	0.0	0.0	1.5	0.5
Burnett	372	74.7	0.5	1.9	17.2	0.5	4.6	0.5
Calumet	476	74.6	2.1	15.1	0.0	7.1	0.6	0.4
Chippewa	1,077	95.4	0.2	1.7	0.3	1.2	0.9	0.4
Clark	560	76.6	0.0	20.0	1.3	0.0	1.6	0.5
Columbia	701	82.5	1.1	13.1	0.7	0.4	1.7	0.4
Crawford C	264	97.3	0.0	0.8	0.4	0.4	0.8	0.4
Dane	6,166	40.6	20.8	27.2	0.5	7.8	2.4	0.7
Dodge	1,044	78.3	0.9	18.7	0.1	0.6	1.0	0.6
Door	340	86.5	0.9	9.4	0.3	1.2	1.2	0.6
Douglas Ct	945	88.1	0.7	2.8	1.5	0.8	4.9	1.2
Dunn	704	84.7	0.7	3.8	0.4	7.7	1.7	1.0
Eau Claire	1,825	79.2	1.3	4.3	0.9	10.6	2.8	0.9
Florence	76	*	*	*	*	*	*	*
Fond du La	1,560	73.5	2.4	17.8	0.1	4.2	1.7	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Forest	238	79.4	0.0	2.9	13.0	0.8	3.8	0.0
Grant Cty	867	91.6	0.3	5.7	0.0	0.8	1.3	0.3
Green Cty	489	87.5	0.2	11.0	0.0	0.2	0.2	0.8
Green Lake	263	73.8	0.4	21.7	0.8	1.5	1.1	0.8
Iowa Cty	319	92.8	0.6	4.1	0.0	0.0	0.9	1.6
Iron Cty	72	*	*	*	*	*	*	*
Jackson	503	78.1	0.0	3.8	16.7	0.2	0.4	0.8
Jefferson	1,303	63.0	1.0	34.3	0.2	0.5	0.8	0.2
Juneau	568	90.8	0.5	4.4	0.2	0.5	3.3	0.2
Kenosha	3,452	48.6	17.1	31.4	0.2	1.0	1.7	0.1
Kewaunee	240	82.1	0.4	13.8	1.7	0.0	1.7	0.4
La Crosse	1,951	76.3	3.7	3.3	0.9	13.2	2.0	0.6
Lafayette	173	85.5	0.0	14.5	0.0	0.0	0.0	0.0
Langlade C	366	93.2	0.3	4.1	0.3	0.0	1.4	0.8
Lincoln Ct	379	91.0	0.5	5.5	0.8	0.3	1.3	0.5
Manitowoc	1,257	73.2	1.5	12.7	0.6	9.4	1.8	0.7
Marathon C	2,066	66.8	1.4	5.3	0.3	25.1	0.9	0.2
Marinette	774	93.4	0.0	3.7	0.1	0.4	1.0	1.3
Marquette	184	83.7	2.2	11.4	0.0	0.0	1.1	1.6
Menominee	423	4.7	0.0	5.7	48.7	0.0	40.2	0.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Milwaukee	30,323	17.9	49.6	26.2	0.5	4.0	1.5	0.4
Monroe	917	82.4	1.1	14.2	0.4	1.1	0.5	0.2
Oconto	458	95.4	0.4	2.8	0.7	0.4	0.0	0.2
Oneida	513	92.8	0.2	3.9	0.4	0.8	1.4	0.6
Outagamie	2,144	66.6	2.7	15.7	1.2	12.0	1.2	0.7
Ozaukee	411	84.7	1.7	8.5	0.2	2.2	1.5	1.2
Pepin	96	*	*	*	*	*	*	*
Pierce	525	89.3	1.0	5.1	1.1	1.0	1.9	0.6
Polk	804	92.5	0.0	3.4	0.9	0.4	1.4	1.5
Portage	952	76.2	0.8	9.8	0.3	12.1	0.5	0.3
Price	281	94.7	0.4	1.8	0.4	2.8	0.0	0.0
Racine Cou	4,442	39.6	27.7	29.5	0.3	0.7	1.7	0.6
Richland C	308	91.6	0.0	7.1	0.0	0.0	0.3	1.0
Rock	3,412	67.7	10.3	18.5	0.1	1.0	1.8	0.7
Rusk	310	93.2	0.0	1.3	0.0	0.6	3.2	1.6
Sauk	992	78.9	0.5	15.2	4.3	0.0	0.6	0.4
Sawyer	479	54.7	0.2	4.0	37.2	0.6	1.9	1.5
Shawano	689	75.2	0.6	6.8	12.6	0.4	4.1	0.3
Sheboygan	1,769	58.7	2.4	20.4	0.2	17.7	0.5	0.2
St. Croix	936	87.5	0.6	5.6	1.0	3.4	1.0	1.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Taylor	320	90.9	0.0	6.6	0.0	0.6	1.6	0.3
Trempealea	600	72.2	0.3	24.5	0.3	0.3	1.7	0.7
Vernon	491	91.4	0.4	5.7	0.2	0.6	1.0	0.6
Vilas	414	56.0	0.2	3.6	37.0	0.0	2.4	0.7
Walworth	1,639	55.2	1.3	40.1	0.9	0.7	1.0	0.9
Washburn	332	92.2	0.6	4.5	0.3	0.6	1.2	0.6
Washington	1,199	85.3	1.3	10.5	0.3	0.8	1.3	0.4
Waukesha	2,058	66.4	3.8	25.9	0.4	1.6	1.5	0.4
Waupaca	694	89.0	0.1	8.9	0.0	0.3	1.0	0.6
Waushara C	362	65.7	0.3	32.9	0.0	0.0	1.1	0.0
Winnebago	2,198	76.0	3.7	11.3	0.3	7.0	1.6	0.1
Wood	1,292	82.7	0.9	8.4	1.9	4.3	1.1	0.8
Wisconsin	101,560	53.1	19.1	19.2	1.9	4.4	1.7	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams Cty	309	0.3	10.7	17.2	59.5	11.7	0.6	0.0
Ashland	463	0.2	4.1	12.5	59.6	20.3	3.2	0.0
Barron	942	0.1	2.9	16.5	65.0	14.6	1.0	0.0
Bayfield C	268	0.0	4.9	10.1	58.2	25.7	1.1	0.0
Brown	5,017	0.4	6.6	12.9	61.9	16.8	1.3	0.1
Buffalo	204	0.0	3.4	10.3	62.3	22.5	1.5	0.0
Burnett	372	0.0	6.2	14.2	64.8	14.5	0.3	0.0
Calumet	476	0.0	4.0	13.9	63.7	17.4	1.1	0.0
Chippewa	1,077	0.0	4.3	11.6	67.0	16.0	1.1	0.0
Clark	560	0.4	5.0	13.4	61.8	18.9	0.4	0.2
Columbia	701	0.1	5.0	14.4	62.8	15.8	1.9	0.0
Crawford C	264	0.0	4.2	15.5	64.0	14.8	1.5	0.0
Dane	6,166	0.3	5.7	11.4	61.5	19.9	1.1	0.1
Dodge	1,044	0.2	5.0	13.9	63.2	16.4	1.2	0.1
Door	340	0.0	4.4	10.9	63.5	19.4	1.8	0.0
Douglas Ct	945	0.2	6.1	12.4	66.9	13.2	1.2	0.0
Dunn	704	0.0	6.1	15.3	61.5	15.5	1.6	0.0
Eau Claire	1,825	0.2	4.0	13.4	67.5	14.1	0.8	0.1
Florence	76	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Fond du La	1,560	0.1	6.3	14.4	63.9	14.1	1.1	0.1
Forest	238	0.0	3.4	14.3	66.4	15.1	0.8	0.0
Grant Cty	867	0.0	4.3	11.9	65.3	17.2	1.4	0.0
Green Cty	489	0.4	5.5	15.7	64.4	13.5	0.4	0.0
Green Lake	263	0.4	5.3	14.4	62.7	16.3	0.8	0.0
Iowa Cty	319	0.0	4.7	12.9	66.5	15.4	0.6	0.0
Iron Cty	72	*	*	*	*	*	*	*
Jackson	503	0.2	5.2	17.1	62.0	14.5	0.8	0.2
Jefferson	1,303	0.3	4.5	12.3	64.6	16.7	1.6	0.0
Juneau	568	0.2	7.6	14.3	63.7	13.2	1.1	0.0
Kenosha	3,452	0.3	7.3	14.9	61.4	15.2	0.9	0.0
Kewaunee	240	0.0	5.0	12.5	60.4	20.0	1.7	0.4
La Crosse	1,951	0.1	5.0	13.3	64.4	15.6	1.5	0.0
Lafayette	173	0.0	2.9	12.7	66.5	15.6	2.3	0.0
Langlade C	366	0.0	3.6	14.5	66.9	14.5	0.5	0.0
Lincoln Ct	379	0.3	4.5	16.4	62.5	15.3	1.1	0.0
Manitowoc	1,257	0.2	5.5	15.8	62.0	15.3	1.3	0.0
Marathon C	2,066	0.1	5.5	14.2	63.2	15.9	1.1	0.0
Marinette	774	0.3	4.1	14.1	66.5	14.2	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marquette	184	0.5	4.3	14.1	62.0	17.4	1.6	0.0
Menominee	423	1.2	13.5	15.4	54.4	14.7	0.9	0.0
Milwaukee	30,323	0.6	8.8	14.0	58.8	16.5	1.2	0.0
Monroe	917	0.2	4.6	15.9	64.8	13.4	1.1	0.0
Oconto	458	0.2	4.6	17.0	61.4	15.7	0.9	0.2
Oneida	513	0.0	2.3	12.9	64.1	18.7	1.9	0.0
Outagamie	2,144	0.2	5.3	13.2	64.6	15.7	1.2	0.0
Ozaukee	411	0.0	3.2	10.0	68.6	17.5	0.7	0.0
Pepin	96	*	*	*	*	*	*	*
Pierce	525	0.4	3.2	15.0	62.7	17.1	1.5	0.0
Polk	804	0.1	6.0	15.3	63.2	13.9	1.4	0.1
Portage	952	0.0	6.5	13.8	66.2	12.3	1.3	0.0
Price	281	0.0	3.9	8.9	66.2	19.6	1.4	0.0
Racine Cou	4,442	0.5	7.7	14.9	60.8	15.1	1.0	0.0
Richland C	308	0.0	4.9	12.3	64.9	16.2	1.6	0.0
Rock	3,412	0.5	6.9	15.6	60.8	15.5	0.7	0.1
Rusk	310	0.0	3.5	14.2	61.6	18.7	1.6	0.3
Sauk	992	0.0	5.6	14.2	63.2	15.6	1.2	0.1
Sawyer	479	0.2	6.7	15.0	61.8	14.8	1.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Shawano	689	0.3	4.4	13.9	63.0	17.3	1.0	0.1
Sheboygan	1,769	0.1	6.5	13.3	61.4	17.5	1.0	0.2
St. Croix	936	0.4	4.4	10.4	65.7	18.6	0.5	0.0
Taylor	320	0.3	4.7	10.3	66.6	17.5	0.6	0.0
Trempealea	600	0.2	3.0	13.8	65.2	17.5	0.3	0.0
Vernon	491	0.0	2.0	11.0	63.5	22.2	1.2	0.0
Vilas	414	0.5	5.6	12.8	58.9	20.3	1.9	0.0
Walworth	1,639	0.3	4.8	13.2	62.4	18.2	1.0	0.1
Washburn	332	0.0	4.5	9.0	68.1	17.5	0.9	0.0
Washington	1,199	0.0	4.5	15.2	62.2	17.3	0.8	0.1
Waukesha	2,058	0.1	3.5	12.8	63.3	19.0	1.2	0.0
Waupaca	694	0.1	4.6	16.9	62.7	14.3	1.3	0.1
Waushara C	362	0.3	4.7	15.7	64.4	13.3	1.4	0.3
Winnebago	2,198	0.0	5.1	14.2	65.7	13.9	1.0	0.1
Wood	1,292	0.2	4.1	14.5	66.1	13.9	1.1	0.1
Wisconsin	101,560	0.3	6.5	13.8	61.8	16.3	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	307	8.1 (9)	52.4 (64)	146	21.9 (24)	51.4 (62)	43	* *	250	20.8(41)	260	2.7 (9)	260	9.2(64)
Ashland	460	6.1 (2)	47.6 (47)	376	23.7 (38)	46.0 (31)	65	* *	359	17.0(24)	379	6.6 (64)	382	4.2(20)
Barron	927	10.1 (39)	49.3 (53)	799	21.7 (21)	53.3 (64)	118	29.7 (22)	758	21.0(44)	675	5.5 (58)	671	4.6(23)
Bayfield C	267	3.0 (1)	52.8 (66)	214	22.9 (32)	48.6 (49)	32	* *	183	12.6 (8)	213	8.0 (69)	212	5.7(32)
Brown	4,965	10.9 (50)	43.5 (19)	2,996	25.8 (57)	45.6 (25)	744	36.2 (30)	4,379	28.2(64)	4,227	5.3 (54)	4,239	6.6(47)
Buffalo	202	6.4 (3)	52.0 (62)	173	19.1 (4)	48.6 (49)	36	* *	164	16.5(22)	143	3.5 (22)	144	6.9(49)
Burnett	371	9.2 (20)	44.7 (28)	217	19.8 (8)	55.3 (68)	57	* *	297	19.2(36)	289	2.1 (3)	290	1.4(1)
Calumet	466	10.5 (44)	40.8 (3)	402	23.1 (33)	43.0 (8)	80	* *	406	17.2(25)	416	7.7 (68)	418	5.7(32)
Chippewa	1,071	9.9 (33)	46.0 (35)	982	25.4 (51)	45.4 (22)	153	22.9 (9)	919	17.8(27)	911	4.6 (41)	906	7.1(53)
Clark	560	8.9 (19)	46.4 (39)	471	23.8 (39)	43.5 (10)	104	28.8 (20)	486	36.0(69)	509	2.8 (11)	512	5.7(32)
Columbia	699	11.7 (61)	42.3 (6)	521	20.2 (13)	49.5 (53)	100	23.0 (10)	572	12.4 (6)	598	3.5 (22)	601	8.3(60)
Crawford C	261	8.4 (12)	47.1 (43)	193	25.4 (51)	44.0 (12)	42	* *	200	23.0(51)	227	4.8 (45)	234	12.8(69)
Dane	6,079	10.5 (44)	43.2 (16)	3,664	26.9 (64)	43.0 (8)	1,134	28.9 (21)	4,143	19.8(38)	4,977	5.0 (48)	4,987	5.4(28)
Dodge	1,027	13.2 (66)	42.6 (10)	733	23.1 (33)	42.6 (7)	189	30.2 (23)	910	24.9(60)	933	4.1 (32)	934	4.1(18)
Door	340	11.2 (57)	42.6 (10)	307	18.6 (3)	45.3 (21)	61	* *	296	15.2(14)	264	5.7 (61)	267	10.9(68)
Douglas Ct	938	8.8 (17)	47.0 (42)	680	19.1 (4)	54.0 (66)	138	32.6 (26)	785	30.2(66)	708	7.3 (65)	712	8.0(58)
Dunn	695	8.6 (16)	49.6 (55)	413	20.3 (14)	51.1 (61)	91	* *	503	14.9(13)	500	4.8 (45)	506	6.9(49)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eau Claire	1,817	11.0 (53)	43.4 (17)	128	26.6 (62)	48.4 (45)	253	27.3 (18)	1,557	24.7 (59)	1,609	4.0 (30)	1,604	3.4 (14)
Florence	75	* *	* *	63	* *	* *	13	* *	54	* *	60	* *	61	* *
Fond du La	1,536	11.3 (58)	43.8 (20)	1,352	27.1 (65)	41.4 (4)	228	35.5 (28)	1,269	24.3 (57)	1,220	5.3 (54)	1,218	6.3 (39)
Forest	238	10.9 (50)	52.9 (67)	201	24.4 (43)	53.7 (65)	28	* *	159	20.8 (41)	191	5.2 (52)	192	8.3 (60)
Grant Cty	859	10.0 (35)	48.8 (52)	623	19.4 (6)	48.6 (49)	150	23.3 (11)	740	17.2 (25)	721	4.6 (41)	727	7.7 (57)
Green Cty	489	11.0 (53)	44.6 (26)	435	21.8 (22)	47.4 (38)	74	* *	411	30.9 (67)	412	1.7 (1)	416	2.9 (11)
Green Lake	251	8.4 (12)	52.6 (65)	212	20.8 (16)	48.1 (43)	48	* *	191	18.3 (30)	199	3.0 (15)	198	2.5 (9)
Iowa Cty	315	8.3 (11)	47.9 (49)	258	24.0 (42)	50.0 (56)	57	* *	232	19.0 (34)	269	3.3 (17)	275	7.6 (56)
Iron Cty	67	* *	* *	52	* *	* *	5	* *	50	* *	58	* *	60	* *
Jackson	501	9.4 (24)	48.7 (51)	433	25.2 (49)	42.5 (6)	75	* *	403	19.9 (39)	339	2.9 (12)	338	2.4 (8)
Jefferson	1,285	10.0 (35)	44.4 (23)	683	22.1 (26)	44.8 (17)	225	15.6 (2)	1,087	12.5 (7)	1,078	4.4 (38)	1,078	6.3 (39)
Juneau	565	9.7 (30)	44.6 (26)	251	26.3 (59)	44.2 (13)	69	* *	452	13.7 (10)	372	3.8 (27)	370	6.2 (38)
Kenosha	3,379	9.6 (27)	46.7 (40)	2,921	24.9 (47)	45.2 (19)	597	36.7 (31)	2,906	29.5 (65)	2,444	3.4 (20)	2,449	6.3 (39)
Kewaunee	240	9.6 (27)	50.8 (60)	215	20.0 (10)	54.4 (67)	30	* *	205	23.9 (55)	179	3.9 (28)	178	1.7 (4)
La Crosse	1,946	11.8 (63)	41.8 (4)	1,089	23.4 (37)	46.1 (32)	250	25.2 (14)	1,574	21.5 (47)	1,732	3.6 (25)	1,722	1.7 (4)
Lafayette	170	7.6 (6)	50.0 (57)	121	21.5 (20)	49.6 (54)	30	* *	124	25.8 (63)	145	2.1 (3)	150	10.7 (67)
Langlade C	364	10.7 (48)	47.5 (46)	317	25.2 (49)	44.5 (15)	64	* *	316	19.0 (34)	315	2.5 (8)	318	5.7 (32)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lincoln Ct	377	7.7 (7)	45.4 (32)	327	22.0 (25)	47.4 (38)	63	* *	329	6.7 (1)	338	3.6 (25)	339	4.4(21)
Manitowoc	1,246	11.7 (61)	42.0 (5)	418	25.6 (54)	48.3 (44)	153	25.5 (15)	1,002	21.0(44)	1,069	4.1 (32)	1,075	6.5(46)
Marathon C	2,048	10.1 (39)	43.1 (14)	1,763	27.5 (66)	39.6 (1)	330	17.9 (4)	1,793	9.8 (2)	1,775	3.5 (22)	1,767	2.7(10)
Marinette	759	14.9 (69)	36.5 (1)	594	18.4 (2)	50.7 (57)	113	17.7 (3)	485	16.7(23)	632	4.7 (43)	641	6.4(42)
Marquette	180	9.4 (24)	47.2 (44)	152	28.3 (67)	45.4 (22)	31	* *	138	19.6(37)	140	4.3 (34)	140	6.4(42)
Menominee	421	7.1 (4)	53.2 (69)	316	25.0 (48)	47.5 (40)	39	* *	370	18.1(29)	362	7.5 (67)	365	9.6(66)
Milwaukee	30,018	8.5 (15)	50.1 (58)	20,688	26.6 (62)	43.8 (11)	4,925	42.7 (33)	25,251	34.7(68)	24,969	4.3 (34)	25,078	6.9(49)
Monroe	893	10.2 (41)	46.1 (37)	5	* *	* *	155	38.1 (32)	796	22.2(48)	729	4.4 (38)	727	2.3(7)
Oconto	454	14.3 (68)	46.0 (35)	431	21.8 (22)	50.8 (58)	78	* *	314	15.3(15)	394	4.8 (45)	390	5.6(31)
Oneida	511	11.0 (53)	43.4 (17)	181	22.7 (29)	49.7 (55)	78	* *	422	12.1 (5)	445	4.5 (40)	448	6.0(36)
Outagamie	2,108	11.9 (64)	42.7 (12)	1,681	24.4 (43)	44.7 (16)	350	28.3 (19)	1,803	24.3(57)	1,834	5.6 (59)	1,823	6.4(42)
Ozaukee	402	13.9 (67)	43.8 (20)	353	16.7 (1)	47.3 (37)	72	* *	339	11.8 (4)	341	5.9 (62)	337	7.1(53)
Pepin	95	* *	* *	85	* *	* *	11	* *	82	* *	68	* *	66	* *
Pierce	523	9.9 (33)	42.4 (8)	349	20.3 (14)	47.9 (41)	55	* *	420	21.0(44)	451	3.3 (17)	453	7.3(55)
Polk	793	11.9 (64)	45.1 (31)	524	19.8 (8)	46.9 (36)	101	35.6 (29)	669	22.4(50)	626	5.4 (57)	621	9.0(62)
Portage	945	10.7 (48)	42.3 (6)	648	22.7 (29)	45.4 (22)	133	26.3 (16)	784	14.2(11)	734	2.7 (9)	739	4.6(23)
Price	279	7.2 (5)	52.0 (62)	255	20.0 (10)	52.5 (63)	36	* *	245	15.9(18)	238	2.9 (12)	237	4.6(23)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Racine Cou	4,393	9.7 (30)	50.1 (58)	3,826	25.7 (56)	45.9 (29)	726	31.5 (25)	3,767	23.4 (53)	3,042	5.2 (52)	3,068	8.0 (58)
Richland C	306	9.2 (20)	52.9 (67)	244	25.4 (51)	48.0 (42)	48	* *	222	23.4 (53)	262	5.0 (48)	267	9.0 (62)
Rock	3,336	10.5 (44)	47.7 (48)	926	20.1 (12)	41.9 (5)	599	33.2 (27)	2,799	25.3 (62)	2,194	2.2 (5)	2,199	3.3 (13)
Rusk	308	8.8 (17)	47.4 (45)	246	26.4 (60)	45.9 (29)	37	* *	251	13.5 (9)	244	3.3 (17)	246	2.0 (6)
Sauk	972	11.3 (58)	48.6 (50)	824	22.8 (31)	46.6 (35)	171	22.8 (8)	825	15.9 (18)	836	5.1 (50)	840	5.5 (30)
Sawyer	478	10.9 (50)	42.7 (12)	367	21.3 (18)	48.5 (46)	62	* *	358	24.9 (60)	296	1.7 (1)	296	1.4 (1)
Shawano	683	9.2 (20)	44.9 (29)	515	22.3 (27)	49.1 (52)	98	* *	368	15.8 (16)	580	4.3 (34)	582	5.3 (27)
Sheboygan	1,765	9.5 (26)	44.1 (22)	1,376	26.2 (58)	40.9 (2)	351	21.4 (6)	1,570	17.9 (28)	1,377	5.3 (54)	1,374	3.5 (15)
St. Croix	911	10.0 (35)	46.2 (38)	817	19.6 (7)	50.8 (58)	128	14.8 (1)	643	10.6 (3)	658	6.5 (63)	654	9.2 (64)
Taylor	319	10.0 (35)	49.8 (56)	291	22.3 (27)	48.5 (46)	45	* *	286	19.9 (39)	266	3.4 (20)	267	4.1 (18)
Trempealea	592	10.3 (42)	42.4 (8)	181	31.5 (68)	40.9 (2)	93	* *	502	23.3 (52)	447	4.7 (43)	447	5.1 (26)
Vernon	490	9.6 (27)	51.6 (61)	341	21.4 (19)	51.0 (60)	68	* *	399	15.8 (16)	431	3.9 (28)	432	3.2 (12)
Vilas	414	8.0 (8)	45.9 (34)	250	26.4 (60)	45.6 (25)	67	* *	314	22.3 (49)	344	2.9 (12)	350	6.0 (36)
Walworth	1,601	8.4 (12)	45.0 (30)	1,430	24.4 (43)	45.8 (27)	250	23.6 (12)	1,345	16.0 (20)	1,194	2.2 (5)	1,188	1.4 (1)
Washburn	331	8.2 (10)	49.5 (54)	264	21.2 (17)	48.5 (46)	38	* *	279	16.1 (21)	226	4.0 (30)	225	4.0 (17)
Washington	1,189	10.3 (42)	43.1 (14)	1,032	23.1 (33)	46.5 (34)	187	19.8 (5)	1,030	14.8 (12)	1,004	3.2 (16)	1,007	6.6 (47)
Waukesha	2,042	11.1 (56)	40.4 (2)	1,826	23.9 (41)	46.2 (33)	401	22.4 (7)	1,584	18.3 (30)	1,688	4.3 (34)	1,706	6.4 (42)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	
Waupaca	688	11.6 (60)	44.5 (25)	620	25.6 (54)	44.4 (14)	103	27.2 (17)	517	18.4 (32)	467	5.6 (59)	465	6.9 (49)
Waushara C	346	9.2 (20)	45.7 (33)	301	24.6 (46)	45.2 (19)	64	* *	274	18.6 (33)	257	7.4 (66)	260	5.4 (28)
Winnebago	2,169	10.6 (47)	44.4 (23)	1,717	23.1 (33)	45.8 (27)	441	23.8 (13)	1,927	20.8 (41)	1,753	5.1 (50)	1,745	4.4 (21)
Wood	1,283	9.8 (32)	46.8 (41)	1,091	23.8 (39)	44.9 (18)	298	30.9 (24)	1,002	24.0 (56)	1,050	2.3 (7)	1,056	3.6 (16)
Wisconsin	100,400	9.8	46.7	69,896	24.7	45.2	16,381	31.9	82,844	25.2	81,333	4.4	81,549	5.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	282	87.2 (41)	3.9 (67)	294	38.8 (18)	256	24.6 (19)	225	86.7 (36)
Ashland	342	86.8 (46)	0.9 (23)	421	40.1 (13)	291	25.4 (16)	264	86.0 (42)
Barron	683	91.4 (6)	0.9 (23)	880	42.3 (11)	569	24.1 (23)	498	86.3 (38)
Bayfield C	203	91.6 (4)	1.0 (28)	254	43.3 (8)	163	22.7 (28)	148	93.9 (2)
Brown	3,746	80.5 (66)	2.3 (52)	4,680	36.5 (24)	3,222	17.0 (54)	2,840	78.5 (62)
Buffalo	153	84.3 (54)	0.0 (1)	190	30.5 (42)	137	32.1 (3)	108	85.2 (44)
Burnett	268	82.8 (58)	0.7 (17)	344	43.3 (8)	218	16.1 (59)	193	83.9 (50)
Calumet	372	86.3 (47)	2.4 (53)	445	25.8 (58)	315	25.4 (16)	249	84.3 (47)
Chippewa	794	91.2 (9)	0.6 (13)	1,032	30.0 (50)	678	26.3 (13)	594	84.5 (46)
Clark	425	88.7 (30)	0.5 (11)	512	22.7 (66)	363	31.1 (4)	300	90.7 (14)
Columbia	558	81.5 (63)	1.4 (36)	680	32.6 (36)	468	39.5 (1)	381	90.6 (15)
Crawford C	186	90.9 (11)	0.0 (1)	236	30.5 (42)	162	29.0 (7)	123	82.1 (57)
Dane	4,573	84.5 (53)	1.4 (36)	5,783	25.2 (61)	3,898	17.1 (53)	3,228	83.5 (52)
Dodge	765	90.3 (16)	0.8 (19)	992	26.0 (57)	679	18.6 (41)	527	86.1 (41)
Door	245	89.4 (23)	0.0 (1)	325	28.0 (52)	218	30.7 (6)	189	93.7 (3)
Douglas Ct	659	87.9 (34)	0.9 (23)	897	40.0 (14)	609	23.2 (25)	533	88.6 (25)
Dunn	515	87.2 (41)	0.4 (8)	667	39.4 (16)	446	22.9 (26)	382	90.6 (15)
Eau Claire	1,290	87.8 (35)	0.4 (8)	1,735	36.7 (23)	1,163	14.8 (62)	1,029	85.0 (45)
Florence	57	* * *	* *	73	* *	45	* *	42	* *
Fond du La	1,138	87.2 (41)	2.6 (56)	1,472	37.0 (22)	999	17.5 (51)	849	81.6 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Forest	179	87.7 (37)	1.7 (45)	225	46.7 (4)	150	27.3 (10)	135	92.6 (6)
Grant Cty	658	90.7 (13)	1.2 (32)	808	30.6 (41)	562	22.6 (29)	454	88.8 (24)
Green Cty	336	89.6 (20)	2.1 (50)	466	36.1 (26)	305	18.0 (47)	277	89.9 (19)
Green Lake	202	85.6 (51)	0.0 (1)	248	25.8 (58)	163	20.9 (35)	124	91.9 (9)
Iowa Cty	230	90.9 (11)	1.7 (45)	304	30.3 (46)	203	20.7 (36)	147	89.1 (22)
Iron Cty	55	* *	* *	66	* *	51	* *	40	* *
Jackson	367	89.1 (26)	1.6 (44)	459	35.7 (28)	311	21.2 (34)	263	73.0 (65)
Jefferson	920	83.7 (56)	4.5 (68)	1,223	32.0 (38)	700	18.0 (47)	721	85.9 (43)
Juneau	462	90.5 (14)	3.2 (60)	538	41.6 (12)	346	27.2 (11)	306	86.3 (38)
Kenosha	2,452	79.5 (67)	3.2 (60)	3,258	30.1 (49)	2,149	16.4 (56)	1,709	82.3 (55)
Kewaunee	178	86.0 (48)	0.6 (13)	228	35.1 (30)	155	28.4 (8)	120	88.3 (26)
La Crosse	1,574	91.1 (10)	0.6 (13)	1,824	39.4 (16)	1,201	19.7 (38)	1,017	56.5 (67)
Lafayette	122	90.2 (18)	0.0 (1)	166	23.5 (65)	102	17.6 (49)	83	* *
Langlade C	271	87.5 (38)	3.7 (64)	343	27.7 (54)	233	17.6 (49)	200	92.0 (8)
Lincoln Ct	286	87.4 (39)	3.1 (58)	366	31.4 (39)	236	17.4 (52)	198	92.9 (4)
Manitowoc	894	83.4 (57)	3.2 (60)	1,187	46.1 (6)	824	18.3 (42)	741	83.5 (52)
Marathon C	1,669	82.0 (59)	4.7 (69)	1,943	30.4 (44)	1,268	16.2 (58)	1,129	86.3 (38)
Marinette	593	91.4 (6)	2.5 (54)	726	39.5 (15)	452	23.9 (24)	406	87.4 (31)
Marquette	148	87.8 (35)	0.7 (17)	175	25.7 (60)	122	31.1 (4)	93	* *
Menominee	327	74.9 (69)	1.8 (49)	387	52.7 (2)	284	7.7 (69)	262	92.4 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Milwaukee	22,395	78.7 (68)	2.2 (51)	28,196	32.6 (36)	19,496	12.6 (65)	16,389	80.1 (60)
Monroe	654	87.0 (45)	0.5 (11)	860	26.4 (55)	559	19.0 (39)	452	78.8 (61)
Oconto	351	89.5 (21)	1.1 (31)	427	33.7 (32)	294	24.5 (20)	240	90.8 (13)
Oneida	392	88.5 (32)	2.6 (56)	485	38.8 (18)	314	25.5 (15)	281	90.4 (17)
Outagamie	1,438	81.8 (61)	1.7 (45)	2,047	30.4 (44)	1,362	22.0 (33)	1,113	80.9 (59)
Ozaukee	295	91.5 (5)	0.3 (7)	387	24.5 (63)	254	24.8 (18)	189	91.0 (11)
Pepin	73	* *	* *	90	* *	59	* *	51	* *
Pierce	382	89.3 (24)	1.0 (28)	492	35.2 (29)	319	22.3 (30)	274	83.9 (50)
Polk	564	87.1 (44)	3.7 (64)	772	42.6 (10)	494	22.1 (32)	422	87.9 (28)
Portage	760	87.4 (39)	1.3 (33)	900	35.9 (27)	343	15.5 (60)	490	84.3 (47)
Price	208	89.9 (19)	1.4 (36)	269	55.0 (1)	188	18.1 (45)	170	82.9 (54)
Racine Cou	3,102	80.9 (64)	1.5 (42)	4,188	34.7 (31)	2,827	12.1 (67)	2,434	78.1 (63)
Richland C	232	92.2 (3)	0.0 (1)	287	26.1 (56)	193	22.8 (27)	160	91.9 (9)
Rock	3,004	89.1 (26)	3.8 (66)	3,248	29.9 (51)	2,222	12.6 (65)	1,839	87.2 (32)
Rusk	218	90.4 (15)	1.4 (36)	286	46.5 (5)	210	32.4 (2)	177	92.7 (5)
Sauk	864	88.4 (33)	1.5 (42)	943	30.3 (46)	692	13.6 (64)	495	88.3 (26)
Sawyer	398	80.7 (65)	3.5 (63)	444	45.0 (7)	295	18.3 (42)	268	86.9 (34)
Shawano	485	89.5 (21)	3.1 (58)	637	37.2 (21)	428	27.1 (12)	379	94.2 (1)
Sheboygan	1,319	81.8 (61)	1.4 (36)	1,678	20.4 (68)	1,181	16.4 (56)	929	86.8 (35)
St. Croix	671	89.3 (24)	0.9 (23)	888	33.0 (34)	585	26.0 (14)	457	91.0 (11)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	277	91.3 (8)	0.4	(8)	309	32.7 (35)	245	28.2 (9)	205	90.2 (18)
Trempealea	486	88.9 (28)	0.8	(19)	553	33.3 (33)	400	14.3 (63)	334	69.8 (66)
Vernon	362	95.9 (1)	0.8	(19)	464	38.8 (18)	282	22.3 (30)	278	77.0 (64)
Vilas	320	81.9 (60)	1.3	(33)	387	30.2 (48)	253	19.0 (39)	233	87.6 (29)
Walworth	1,133	83.9 (55)	0.9	(23)	1,543	27.9 (53)	1,011	20.6 (37)	817	84.2 (49)
Washburn	240	94.2 (2)	1.3	(33)	314	48.1 (3)	215	24.2 (22)	188	89.9 (19)
Washington	888	90.3 (16)	1.7	(45)	1,138	36.4 (25)	805	16.6 (55)	654	87.5 (30)
Waukesha	1,542	86.0 (48)	0.6	(13)	1,965	22.7 (66)	1,305	18.2 (44)	975	89.1 (22)
Waupaca	514	88.9 (28)	1.4	(36)	659	31.0 (40)	435	9.2 (68)	353	87.0 (33)
Waushara C	292	84.6 (52)	1.0	(28)	351	24.8 (62)	241	24.5 (20)	186	89.2 (21)
Winnebago	1,602	86.0 (48)	2.5	(54)	2,063	23.9 (64)	1,449	15.4 (61)	1,174	82.2 (56)
Wood	987	88.7 (30)	0.8	(19)	1,185	19.5 (69)	875	18.1 (45)	699	86.6 (37)
Wisconsin	75,625	83.7	2.0		95,307	32.4	64,547	16.9	54,432	82.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	293	49.1 (55)	282	32.6 (58)	117	24.8 (38)	245	38.8 (68)	254	21.3 (69)
Ashland	409	52.1 (61)	422	35.3 (65)	183	26.8 (28)	348	25.0 (40)	395	6.1 (22)
Barron	807	45.6 (42)	857	30.1 (44)	318	22.0 (45)	670	21.3 (27)	791	5.6 (15)
Bayfield C	230	52.2 (62)	237	32.5 (56)	102	30.4 (14)	194	21.1 (26)	218	5.5 (13)
Brown	4,438	38.0 (18)	4,567	20.8 (12)	1,367	32.5 (8)	3,616	20.4 (23)	4,192	8.2 (41)
Buffalo	165	46.1 (44)	185	31.9 (54)	64	* *	135	16.3 (4)	172	5.8 (17)
Burnett	323	56.7 (67)	337	39.2 (69)	148	16.9 (54)	269	35.7 (67)	301	9.3 (50)
Calumet	415	38.8 (19)	446	25.3 (28)	129	27.9 (26)	320	18.8 (15)	402	5.5 (13)
Chippewa	935	53.9 (65)	1,001	31.4 (50)	422	31.0 (12)	761	24.8 (38)	931	6.2 (24)
Clark	513	39.0 (20)	526	23.6 (20)	159	25.2 (36)	403	20.1 (22)	480	7.7 (38)
Columbia	624	50.5 (57)	629	28.6 (38)	232	24.1 (39)	483	27.5 (47)	563	8.3 (43)
Crawford C	220	40.5 (22)	241	27.8 (35)	67	* *	173	19.7 (20)	224	5.8 (17)
Dane	5,341	27.4 (2)	5,477	16.4 (2)	1,027	30.6 (13)	4,216	15.3 (2)	5,056	5.9 (20)
Dodge	908	41.7 (25)	957	27.1 (32)	291	23.0 (41)	695	20.9 (25)	885	6.2 (24)
Door	289	43.6 (32)	310	21.6 (13)	108	39.8 (3)	225	15.6 (3)	286	4.2 (6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Douglas Ct	812	56.3 (66)	824	32.5 (56)	381	26.2 (30)	699	29.3 (52)	770	6.4 (28)
Dunn	609	45.0 (39)	620	24.0 (22)	215	30.2 (16)	498	20.7 (24)	575	4.9 (8)
Eau Claire	1,560	44.0 (35)	1,681	27.2 (33)	573	27.7 (27)	1,270	19.7 (20)	1,529	3.9 (4)
Florence	63	* *	65	* *	24	* *	50	* *	59	* *
Fond du La	1,364	44.7 (37)	1,415	24.7 (24)	494	28.1 (24)	1,104	23.1 (32)	1,292	7.4 (35)
Forest	210	45.2 (40)	212	29.2 (41)	71	* *	174	32.2 (62)	197	11.7 (58)
Grant Cty	741	35.1 (12)	813	22.5 (18)	218	26.1 (32)	584	18.5 (12)	760	6.1 (22)
Green Cty	429	53.4 (63)	433	31.4 (50)	188	31.4 (11)	360	32.5 (64)	391	13.0 (65)
Green Lake	204	36.8 (16)	230	25.7 (30)	51	* *	156	26.9 (44)	216	11.1 (56)
Iowa Cty	274	48.9 (53)	286	37.4 (68)	104	11.5 (57)	204	25.5 (42)	264	7.2 (34)
Iron Cty	60	* *	63	* *	26	* *	48	* *	59	* *
Jackson	443	49.4 (56)	460	34.3 (61)	185	16.2 (55)	365	30.7 (60)	426	7.0 (31)
Jefferson	1,128	35.8 (14)	1,169	22.2 (15)	316	29.7 (20)	914	17.9 (11)	1,071	7.5 (36)
Juneau	505	43.4 (30)	519	31.6 (52)	175	13.1 (56)	416	34.1 (65)	478	16.9 (66)
Kenosha	2,887	33.1 (5)	3,116	20.7 (11)	743	29.9 (18)	2,278	27.9 (49)	2,855	10.8 (55)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee	224	34.8 (10)	224	20.1 (8)	63	* *	176	23.9 (36)	207	6.8 (30)
La Crosse	1,694	35.0 (11)	1,753	20.1 (8)	472	22.2 (44)	1,370	15.2 (1)	1,599	2.8 (1)
Lafayette	139	43.9 (34)	153	22.2 (15)	39	* *	103	19.4 (17)	145	6.2 (24)
Langlade C	313	48.9 (53)	336	34.5 (62)	127	17.3 (53)	249	29.7 (55)	313	11.5 (57)
Lincoln Ct	330	46.7 (48)	362	31.8 (53)	123	19.5 (51)	263	38.8 (68)	340	12.6 (64)
Manitowoc	1,115	46.4 (45)	1,138	31.2 (48)	428	21.5 (46)	908	26.4 (43)	1,043	9.7 (51)
Marathon C	1,849	36.0 (15)	1,924	24.7 (24)	536	20.3 (49)	1,457	18.5 (12)	1,750	6.3 (27)
Marinette	640	51.1 (60)	685	33.3 (60)	274	23.0 (41)	510	27.6 (48)	611	9.2 (48)
Marquette	161	46.6 (47)	161	36.0 (66)	53	* *	120	30.0 (58)	145	9.7 (51)
Menominee	373	66.2 (69)	389	35.0 (64)	219	39.7 (4)	321	29.3 (52)	358	5.3 (11)
Milwaukee	26,150	23.8 (1)	27,532	13.1 (1)	4,812	34.0 (6)	21,456	23.4 (33)	25,327	12.1 (62)
Monroe	775	45.5 (41)	845	32.3 (55)	296	19.6 (50)	627	23.6 (35)	770	8.2 (41)
Oconto	389	48.8 (52)	434	36.9 (67)	166	18.1 (52)	316	31.3 (61)	408	12.0 (61)
Oneida	454	60.4 (68)	461	34.7 (63)	211	29.9 (18)	354	29.4 (54)	420	6.0 (21)
Outagamie	1,818	42.1 (28)	1,934	22.2 (15)	592	30.2 (16)	1,438	17.3 (9)	1,766	5.2 (10)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ozaukee	336	43.5 (31)	355	17.5 (3)	104	45.2 (1)	247	18.6 (14)	329	4.6 (7)
Pepin	85	* *	89	* *	26	* *	65	* *	82	* *
Pierce	443	48.1 (49)	466	29.2 (41)	168	26.8 (28)	355	30.1 (59)	429	8.9 (47)
Polk	676	51.0 (59)	731	30.6 (46)	290	22.8 (43)	539	28.8 (51)	671	5.8 (17)
Portage	831	33.6 (8)	859	18.4 (5)	228	28.5 (23)	662	16.5 (5)	786	4.1 (5)
Price	256	41.0 (24)	256	24.2 (23)	89	* *	227	22.9 (30)	239	9.2 (48)
Racine Cou	3,790	33.0 (4)	4,001	18.8 (6)	998	29.6 (21)	3,086	23.0 (31)	3,652	7.5 (36)
Richland C	270	45.6 (42)	276	30.8 (47)	97	* *	205	24.9 (39)	250	3.6 (2)
Rock	2,915	40.8 (23)	3,056	25.6 (29)	959	26.1 (32)	2,328	26.9 (44)	2,773	10.5 (54)
Rusk	283	42.0 (27)	282	28.4 (36)	98	* *	236	25.4 (41)	267	8.6 (46)
Sauk	902	44.3 (36)	889	31.2 (48)	303	20.5 (48)	688	32.4 (63)	798	17.4 (68)
Sawyer	428	50.7 (58)	413	27.6 (34)	176	32.4 (9)	351	27.1 (46)	378	10.1 (53)
Shawano	610	48.2 (50)	636	29.2 (41)	248	31.5 (10)	489	23.5 (34)	578	7.1 (32)
Sheboygan	1,564	33.2 (7)	1,632	20.2 (10)	385	25.2 (36)	1,233	19.1 (16)	1,500	8.3 (43)
St. Croix	797	42.8 (29)	812	23.3 (19)	254	30.3 (15)	610	16.6 (6)	744	6.5 (29)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	307	43.6 (32)	303	28.7 (39)	107	26.2 (30)	238	28.6 (50)	280	11.8 (60)
Trempealea	525	32.2 (3)	545	20.0 (7)	140	20.7 (47)	426	16.9 (7)	498	3.6 (2)
Vernon	442	35.1 (12)	454	23.6 (20)	129	23.3 (40)	356	19.4 (17)	423	5.4 (12)
Vilas	368	53.8 (64)	363	32.8 (59)	161	25.5 (35)	304	29.9 (57)	331	12.1 (62)
Walworth	1,378	34.5 (9)	1,469	18.1 (4)	360	41.1 (2)	1,064	16.9 (7)	1,355	4.9 (8)
Washburn	280	39.6 (21)	296	28.4 (36)	96	* *	238	17.6 (10)	273	7.7 (38)
Washington	1,056	48.7 (51)	1,094	25.0 (26)	409	35.5 (5)	822	22.5 (29)	1,009	7.1 (32)
Waukesha	1,777	41.8 (26)	1,881	21.9 (14)	520	33.8 (7)	1,292	19.4 (17)	1,718	5.7 (16)
Waupaca	589	46.5 (46)	634	30.4 (45)	210	25.7 (34)	445	34.2 (66)	586	17.1 (67)
Waushara C	290	33.1 (5)	311	27.0 (31)	72	* *	229	29.7 (55)	290	11.7 (58)
Winnebago	1,938	44.9 (38)	1,990	28.7 (39)	656	29.1 (22)	1,532	22.4 (28)	1,838	8.4 (45)
Wood	1,141	37.6 (17)	1,167	25.1 (27)	329	28.0 (25)	905	24.3 (37)	1,091	7.9 (40)
Wisconsin	87,900	35.4	91,971	21.1	24,521	28.8	70,713	22.8	84,462	9.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	267	9.4 (68)	11.6 (62)	263	11.8 (64)	226	4.0 (63)	269	65.1 (44)
Ashland	408	3.7 (1)	12.0 (64)	384	8.1 (8)	344	2.3 (17)	416	67.3 (38)
Barron	804	7.3 (46)	9.7 (48)	794	10.6 (48)	674	2.5 (23)	835	69.6 (29)
Bayfield C	219	5.5 (9)	12.8 (67)	216	8.3 (10)	193	2.6 (26)	225	74.2 (12)
Brown	4,391	7.2 (43)	9.1 (40)	4,234	11.7 (62)	3,627	2.8 (38)	4,463	62.9 (51)
Buffalo	178	7.3 (46)	6.7 (5)	172	11.0 (52)	149	0.7 (1)	180	74.4 (9)
Burnett	317	5.7 (11)	7.6 (19)	305	9.8 (36)	266	1.9 (11)	319	70.5 (25)
Calumet	435	8.5 (60)	8.0 (25)	413	9.2 (26)	370	2.4 (20)	440	64.5 (48)
Chippewa	972	7.3 (46)	7.2 (11)	954	10.0 (38)	843	2.5 (23)	984	67.4 (37)
Clark	505	8.5 (60)	7.9 (21)	468	10.5 (47)	415	3.9 (62)	508	74.8 (8)
Columbia	590	7.5 (51)	10.5 (56)	592	7.9 (7)	529	4.2 (67)	602	70.3 (27)
Crawford C	242	6.6 (28)	8.3 (27)	219	9.1 (21)	198	1.5 (6)	244	66.0 (41)
Dane	5,246	6.8 (31)	8.9 (36)	5,112	9.3 (28)	4,458	2.7 (33)	5,355	77.7 (3)
Dodge	917	7.5 (51)	7.5 (16)	908	11.1 (54)	781	1.8 (10)	940	76.1 (7)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Door	297	4.7 (3)	9.1 (40)	289	11.4 (57)	253	2.4 (20)	298	72.5 (16)	
Douglas Ct	807	4.3 (2)	11.5 (61)	790	9.9 (37)	702	1.0 (2)	812	70.9 (24)	
Dunn	599	5.3 (7)	10.7 (59)	586	7.7 (4)	524	2.7 (33)	611	71.2 (19)	
Eau Claire	1,646	6.0 (16)	8.4 (29)	1,592	8.5 (11)	1,437	2.7 (33)	1,661	71.0 (23)	
Florence	63	* * *	* *	62	* *	55	* *	64	* *	
Fond du La	1,341	7.8 (54)	7.1 (9)	1,331	10.7 (50)	1,145	3.3 (53)	1,370	68.2 (34)	
Forest	193	8.8 (65)	7.3 (13)	191	11.5 (60)	161	3.7 (58)	198	52.0 (68)	
Grant Cty	772	6.2 (21)	9.5 (45)	748	9.1 (21)	651	2.3 (17)	794	63.4 (49)	
Green Cty	419	7.4 (49)	10.3 (54)	403	11.9 (66)	351	2.6 (26)	425	60.5 (57)	
Green Lake	225	6.2 (21)	10.2 (52)	220	8.6 (14)	195	3.1 (46)	231	55.8 (65)	
Iowa Cty	279	8.2 (58)	7.9 (21)	270	7.8 (6)	244	5.3 (69)	284	67.6 (35)	
Iron Cty	62	* * *	* *	61	* *	52	* *	65	* *	
Jackson	440	6.1 (19)	12.7 (66)	416	10.1 (41)	362	1.7 (9)	452	73.9 (13)	
Jefferson	1,114	4.9 (5)	8.4 (29)	1,086	10.0 (38)	941	1.4 (4)	1,133	71.2 (19)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Juneau	504	5.6 (10)	8.3 (27)	490	8.6 (14)	442	2.0 (14)	508	69.1 (30)
Kenosha	3,034	8.5 (60)	6.0 (3)	2,953	10.4 (43)	2,575	3.8 (59)	3,076	61.3 (55)
Kewaunee	213	6.6 (28)	7.5 (16)	209	5.7 (1)	191	3.1 (46)	216	72.7 (14)
La Crosse	1,673	7.1 (40)	10.5 (56)	1,614	9.5 (30)	1,416	3.2 (51)	1,696	72.2 (18)
Lafayette	149	4.7 (3)	5.4 (1)	148	8.1 (8)	132	3.8 (59)	149	67.1 (39)
Langlade C	316	7.9 (55)	9.5 (45)	312	7.7 (4)	273	4.4 (68)	328	57.6 (63)
Lincoln Ct	344	5.8 (12)	9.0 (38)	344	6.4 (2)	312	2.9 (41)	349	60.2 (58)
Manitowoc	1,111	7.9 (55)	7.5 (16)	1,085	11.0 (52)	942	4.0 (63)	1,139	59.4 (59)
Marathon C	1,846	6.6 (28)	5.9 (2)	1,800	8.7 (16)	1,596	2.6 (26)	1,878	58.4 (62)
Marinette	665	7.1 (40)	9.0 (38)	641	9.7 (33)	565	2.8 (38)	672	50.9 (69)
Marquette	157	7.0 (34)	7.0 (7)	154	10.4 (43)	136	2.9 (41)	161	62.1 (52)
Menominee	373	7.0 (34)	12.1 (65)	345	10.4 (43)	304	2.3 (17)	376	67.6 (35)
Milwaukee	26,020	9.2 (67)	6.1 (4)	25,157	11.4 (57)	21,446	3.8 (59)	26,671	63.1 (50)
Monroe	807	6.8 (31)	8.6 (32)	795	9.3 (28)	690	2.9 (41)	829	68.9 (31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Oconto	419	8.1 (57)	7.9 (21)	401	9.0 (19)	357	3.6 (57)	425	53.9 (66)
Oneida	444	7.7 (53)	7.9 (21)	437	8.9 (17)	391	3.1 (46)	450	61.6 (53)
Outagamie	1,834	8.6 (64)	7.1 (9)	1,825	10.4 (43)	1,567	3.2 (51)	1,884	66.0 (41)
Ozaukee	354	5.9 (14)	10.2 (52)	343	11.1 (54)	299	2.0 (14)	362	77.3 (4)
Pepin	84	*	*	81	*	73	*	88	*
Pierce	461	5.4 (8)	8.9 (36)	443	10.2 (42)	390	1.5 (6)	470	68.3 (33)
Polk	712	6.3 (24)	7.2 (11)	701	7.6 (3)	633	2.7 (33)	723	76.9 (5)
Portage	832	6.9 (33)	9.6 (47)	810	10.0 (38)	710	3.1 (46)	847	68.7 (32)
Price	243	5.8 (12)	9.1 (40)	241	9.1 (21)	213	1.4 (4)	250	70.0 (28)
Racine Cou	3,877	9.0 (66)	7.0 (7)	3,749	11.7 (62)	3,234	3.3 (53)	3,927	59.4 (59)
Richland C	265	6.0 (16)	15.1 (69)	253	9.5 (30)	223	2.2 (16)	272	76.8 (6)
Rock	2,977	7.2 (43)	8.7 (33)	2,933	9.6 (32)	2,572	2.9 (41)	3,033	52.1 (67)
Rusk	270	7.0 (34)	11.9 (63)	255	10.6 (48)	224	4.0 (63)	272	61.4 (54)
Sauk	858	5.2 (6)	10.6 (58)	846	9.7 (33)	748	1.9 (11)	876	78.2 (2)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Sawyer	404	6.2 (21)	9.7 (48)	384	8.9 (17)	342	2.6 (26)	408	66.7 (40)	
Shawano	607	7.4 (49)	8.7 (33)	579	12.1 (67)	496	2.8 (38)	617	65.5 (43)	
Sheboygan	1,569	6.4 (26)	6.9 (6)	1,525	9.0 (19)	1,360	2.9 (41)	1,590	64.7 (46)	
St. Croix	803	7.0 (34)	9.7 (48)	779	9.2 (26)	687	2.5 (23)	825	74.4 (9)	
Taylor	286	5.9 (14)	10.8 (60)	289	9.7 (33)	252	1.6 (8)	291	70.4 (26)	
Trempealea	511	6.3 (24)	10.0 (51)	497	11.3 (56)	419	2.6 (26)	525	72.6 (15)	
Vernon	434	7.1 (40)	12.9 (68)	424	8.5 (11)	378	2.6 (26)	442	74.4 (9)	
Vilas	348	6.0 (16)	10.3 (54)	338	12.1 (67)	288	1.0 (2)	351	57.5 (64)	
Walworth	1,436	6.5 (27)	9.1 (40)	1,398	8.5 (11)	1,243	2.6 (26)	1,472	79.0 (1)	
Washburn	289	8.3 (59)	8.7 (33)	278	12.9 (69)	238	3.4 (56)	291	71.1 (22)	
Washington	1,023	7.2 (43)	7.4 (15)	1,023	10.9 (51)	879	3.1 (46)	1,054	71.2 (19)	
Waukesha	1,818	6.1 (19)	7.8 (20)	1,780	9.1 (21)	1,577	2.4 (20)	1,851	72.4 (17)	
Waupaca	577	9.9 (69)	8.1 (26)	591	11.8 (64)	483	4.1 (66)	600	61.3 (55)	
Waushara C	302	7.0 (34)	9.3 (44)	312	11.5 (60)	260	1.9 (11)	313	64.9 (45)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Winnebago	1,936	8.5 (60)	7.3 (13)	1,869	11.4 (57)	1,610	3.3 (53)	1,971	64.7 (46)
Wood	1,087	7.0 (34)	8.4 (29)	1,053	9.1 (21)	902	2.7 (33)	1,112	58.6 (61)
Wisconsin	88,020	7.7	7.8	85,563	10.4	74,214	3.1	89,818	65.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

