

Table 3G
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	2,176	41.5	21.6	27.2	0.5	7.0	2.2	0.0
Mil. Cty	10,502	17.9	49.5	26.4	0.5	4.4	1.3	0.0
Racine	1,582	41.2	27.5	28.5	0.3	0.8	1.8	0.0
Rest of WI	21,289	73.5	3.2	14.1	2.6	4.5	2.1	0.0
Wisconsin	35,549	53.7	19.1	19.2	1.7	4.5	1.8	0.0
Nation (Prior Year)	1,252,728	41.7	23.3	27.9	1.1	3.2	1.0	1.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4G
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	2,176	0.3	4.9	10.7	60.6	22.2	1.2	0.0
Mil. Cty	10,502	0.5	8.5	13.8	58.8	17.0	1.2	0.0
Racine	1,582	0.6	7.5	13.8	61.0	16.1	0.9	0.0
Rest of WI	21,289	0.2	5.1	13.5	63.2	16.9	1.1	0.0
Wisconsin	35,549	0.3	6.2	13.4	61.6	17.2	1.1	0.0
Nation (Prior Year)	1,252,728	0.4	6.5	12.7	59.9	19.0	1.4	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5G
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area

Comparison of Maternal Health Indicators ^(2, 3)														
Metro Area	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	2,144	9.4 (3)	44.7 (1)	1,251	26.0 (4)	42.9 (1)	403	31.5 (2)	1,620	17.2 (1)	1,940	5.4 (4)	1,940	4.9(1)
Mil. Cty	10,390	8.4 (1)	50.3 (4)	6,950	25.5 (3)	45.3 (2)	1,789	44.7 (4)	8,866	34.9 (4)	9,546	4.1 (3)	9,546	6.6(3)
Racine	1,567	9.3 (2)	49.8 (3)	1,369	25.3 (2)	45.6 (3)	286	35.3 (3)	1,348	24.6 (3)	1,430	3.8 (1)	1,430	6.7(4)
Rest of WI	21,040	10.0 (4)	45.8 (2)	14,519	23.3 (1)	46.3 (4)	3,505	26.2 (1)	17,523	20.4 (2)	19,334	3.8 (1)	19,334	5.1(2)
Wisconsin	35,141	9.5	47.2	24,089	24.2	45.8	5,983	32.5	29,357	24.8	32,250	4.0	32,250	5.6
Nation (Prior Year)	1,175,574	10.9	44.5	922,658	25.0	42.8	180,176	33.5	414,735	29.8	230,547	3.9	280,574	4.3

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
- (4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
- (5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6G
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area

Comparison of Maternal Behavioral Indicators ^(2, 3)													
Metro Area	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Pregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
Dane Cty	1,946	85.4 (2)	1.6	(1)	2,099	26.7	(4)	2,040	15.9	(2)	1,685	82.6	(2)
Mil. Cty	9,695	78.8 (4)	2.6	(4)	10,130	32.9	(3)	10,284	11.9	(4)	8,419	79.5	(3)
Racine	1,426	81.0 (3)	2.1	(2)	1,542	35.9	(1)	1,554	12.9	(3)	1,302	78.1	(4)
Rest of WI	19,370	86.8 (1)	2.3	(3)	20,662	33.6	(2)	20,508	18.7	(1)	16,813	83.5	(1)
Wisconsin	32,437	84.0	2.3		34,433	33.0		34,386	16.3		28,219	82.0	
Nation (Prior Year)	612,462	79.4	4.7		1,034,046	31.3		98,170	15.7		238,509	75.9	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7G
2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	2,097	28.2 (2)	1,933	15.6 (2)	402	33.8 (2)	1,720	14.1 (1)	1,934	5.3 (1)
Mil. Cty	9,931	24.4 (1)	9,543	12.7 (1)	1,812	35.9 (1)	8,531	22.3 (3)	9,542	10.8 (4)
Racine	1,516	34.3 (3)	1,428	17.4 (3)	411	31.9 (3)	1,311	23.4 (4)	1,428	7.6 (3)
Rest of WI	20,425	43.6 (4)	19,286	25.0 (4)	6,868	29.4 (4)	17,057	21.9 (2)	19,300	6.7 (2)
Wisconsin	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
Nation (Prior Year)	895,227	26.6	716,629	15.8	161,864	35.1	572,120	20.9	486,636	22.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8G
2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area

Comparison of Infant Health Indicators ^(2, 3)										
Metro Area	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	1,846	6.4 (1)	9.4 (4)		1,877	9.7 (1)	1,609	2.8 (2)	1,882	78.7 (1)
Mil. Cty	9,159	9.0 (3)	6.1 (1)		9,235	10.7 (3)	7,898	3.7 (3)	9,272	64.6 (3)
Racine	1,388	9.9 (4)	6.8 (2)		1,393	12.3 (4)	1,186	4.1 (4)	1,401	61.0 (4)
Rest of WI	18,640	7.0 (2)	8.6 (3)		18,839	9.8 (2)	16,355	2.7 (1)	18,944	67.4 (2)
Wisconsin	31,033	7.7	7.8		31,344	10.2	27,048	3.1	31,499	67.0
Nation (Prior Year)	1,039,036	8.2	6.7		947,705	11.9	856,467	3.7	936,194	64.6

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3G
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Metro Area
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dane Cty	6,166	40.6	20.8	27.2	0.5	7.8	2.4	0.7
Mil. Cty	30,323	17.9	49.6	26.2	0.5	4.0	1.5	0.4
Racine	4,442	39.6	27.7	29.5	0.3	0.7	1.7	0.6
Rest of WI	60,629	73.0	3.1	14.2	2.9	4.5	1.8	0.5
Wisconsin	101,560	53.1	19.1	19.2	1.9	4.4	1.7	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4G
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Metro Area
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
Metro Area	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Dane Cty	6,166	0.3	5.7	11.4	61.5	19.9	1.1	0.1
Mil. Cty	30,323	0.6	8.8	14.0	58.8	16.5	1.2	0.0
Racine	4,442	0.5	7.7	14.9	60.8	15.1	1.0	0.0
Rest of WI	60,629	0.2	5.3	13.8	63.5	16.0	1.1	0.1
Wisconsin	101,560	0.3	6.5	13.8	61.8	16.3	1.1	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5G
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
Metro Area	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	6,079	10.5 (4)	43.2 (1)	3,664	26.9 (4)	43.0 (1)	1,134	28.9 (2)	4,143	19.8 (1)	4,977	5.0 (3)	4,987	5.4 (2)
Mil. Cty	30,018	8.5 (1)	50.1 (3)	20,688	26.6 (3)	43.8 (2)	4,925	42.7 (4)	25,251	34.7 (4)	24,969	4.3 (1)	25,078	6.9 (3)
Racine	4,393	9.7 (2)	50.1 (3)	3,826	25.7 (2)	45.9 (3)	726	31.5 (3)	3,767	23.4 (3)	3,042	5.2 (4)	3,068	8.0 (4)
Rest of WI	59,910	10.4 (3)	45.0 (2)	41,718	23.6 (1)	46.1 (4)	9,596	26.7 (1)	49,683	20.9 (2)	48,345	4.3 (1)	48,416	5.3 (1)
Wisconsin	100,400	9.8	46.7	69,896	24.7	45.2	16,381	31.9	82,844	25.2	81,333	4.4	81,549	5.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6G
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by Metro Area
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)												
Metro Area	Medical Care				WIC Enrollment			Multivitamin Consumption				
	1st Trimester		No Care		1st Trimester			Pregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	% (Rank)
Dane Cty	4,573	84.5 (2)	1.4	(1)	5,783	25.2	(4)	3,898	17.1	(2)	3,228	83.5 (2)
Mil. Cty	22,395	78.7 (4)	2.2	(4)	28,196	32.6	(3)	19,496	12.6	(3)	16,389	80.1 (3)
Racine	3,102	80.9 (3)	1.5	(2)	4,188	34.7	(1)	2,827	12.1	(4)	2,434	78.1 (4)
Rest of WI	45,555	86.3 (1)	1.9	(3)	57,140	32.8	(2)	38,326	19.4	(1)	32,381	84.4 (1)
Wisconsin	75,625	83.7	2.0		95,307	32.4		64,547	16.9		54,432	82.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7G
2006 - 2008 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Metro Area
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
Metro Area	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	5,341	27.4 (2)	5,477	16.4 (2)	1,027	30.6 (2)	4,216	15.3 (1)	5,056	5.9 (1)
Mil. Cty	26,150	23.8 (1)	27,532	13.1 (1)	4,812	34.0 (1)	21,456	23.4 (4)	25,327	12.1 (4)
Racine	3,790	33.0 (3)	4,001	18.8 (3)	998	29.6 (3)	3,086	23.0 (2)	3,652	7.5 (2)
Rest of WI	52,619	42.2 (4)	54,961	25.7 (4)	17,684	27.2 (4)	41,955	23.2 (3)	50,427	7.8 (3)
Wisconsin	87,900	35.4	91,971	21.1	24,521	28.8	70,713	22.8	84,462	9.0

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8G
2006 - 2008 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Metro Area
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
Metro Area	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dane Cty	5,246	6.8 (1)	8.9 (4)	5,112	9.3 (1)	4,458	2.7 (1)	5,355	77.7 (1)
Mil. Cty	26,020	9.2 (4)	6.1 (1)	25,157	11.4 (3)	21,446	3.8 (4)	26,671	63.1 (3)
Racine	3,877	9.0 (3)	7.0 (2)	3,749	11.7 (4)	3,234	3.3 (3)	3,927	59.4 (4)
Rest of WI	52,877	7.0 (2)	8.5 (3)	51,545	9.9 (2)	45,076	2.8 (2)	53,865	66.2 (2)
Wisconsin	88,020	7.7	7.8	85,563	10.4	74,214	3.1	89,818	65.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one metro area's rate to other metro areas. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

