

**Table 3E**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Region**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Region</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
NERO	6,599	66.5	3.1	16.2	4.1	6.9	3.1	0.0
NRO	2,807	77.1	0.9	6.0	6.1	8.3	1.5	0.0
SERO	15,641	30.1	37.7	27.1	0.5	3.3	1.4	0.0
SRO	5,756	66.3	10.3	18.0	0.5	3.0	1.8	0.0
WRO	4,746	84.3	1.3	6.5	1.6	4.4	1.8	0.0
<b>Wisconsin</b>	<b>35,549</b>	<b>53.7</b>	<b>19.1</b>	<b>19.2</b>	<b>1.7</b>	<b>4.5</b>	<b>1.8</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,252,728</b>	<b>41.7</b>	<b>23.3</b>	<b>27.9</b>	<b>1.1</b>	<b>3.2</b>	<b>1.0</b>	<b>1.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4E**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Region**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Region</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
NERO	6,599	0.2	5.8	13.5	62.4	16.9	1.0	0.0
NRO	2,807	0.2	4.9	13.5	63.2	16.7	1.4	0.0
SERO	15,641	0.5	7.6	13.9	59.8	17.1	1.2	0.0
SRO	5,756	0.3	5.3	12.6	62.1	18.6	1.1	0.0
WRO	4,746	0.1	4.0	13.0	65.2	16.7	1.0	0.0
<b>Wisconsin</b>	<b>35,549</b>	<b>0.3</b>	<b>6.2</b>	<b>13.4</b>	<b>61.6</b>	<b>17.2</b>	<b>1.1</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,252,728</b>	<b>0.4</b>	<b>6.5</b>	<b>12.7</b>	<b>59.9</b>	<b>19.0</b>	<b>1.4</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5E**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Region**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Region</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	6,516	10.2 (4)	44.8 (1)	4,839	24.3 (3)	45.4 (3)	1,112	26.9 (2)	5,431	22.2 (4)	6,010	4.9 (5)	6,010	5.7(4)
NRO	2,792	10.7 (5)	45.2 (2)	2,116	25.2 (5)	44.0 (1)	477	21.6 (1)	2,288	15.1 (1)	2,567	3.2 (1)	2,567	3.5(1)
SERO	15,461	8.8 (1)	49.0 (5)	11,310	24.6 (4)	45.8 (4)	2,694	39.0 (5)	13,193	30.1 (5)	14,213	3.8 (2)	14,213	6.3(5)
SRO	5,673	10.0 (3)	46.3 (3)	3,175	23.9 (2)	44.3 (2)	1,012	30.2 (4)	4,599	19.7 (2)	5,189	3.9 (3)	5,189	5.0(2)
WRO	4,699	9.4 (2)	46.8 (4)	2,649	22.0 (1)	49.9 (5)	688	27.0 (3)	3,846	22.0 (3)	4,271	4.1 (4)	4,271	5.1(3)
<b>Wisconsin</b>	<b>35,141</b>	<b>9.5</b>	<b>47.2</b>	<b>24,089</b>	<b>24.2</b>	<b>45.8</b>	<b>5,983</b>	<b>32.5</b>	<b>29,357</b>	<b>24.8</b>	<b>32,250</b>	<b>4.0</b>	<b>32,250</b>	<b>5.6</b>
<b>Nation (Prior Year)</b>	<b>1,175,574</b>	<b>10.9</b>	<b>44.5</b>	<b>922,658</b>	<b>25.0</b>	<b>42.8</b>	<b>180,176</b>	<b>33.5</b>	<b>414,735</b>	<b>29.8</b>	<b>230,547</b>	<b>3.9</b>	<b>280,574</b>	<b>4.3</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6E**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Region**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>													
<b>Region</b>	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
NERO	5,942	84.0 (4)	2.5	(3)	6,389	32.6	(3)	6,408	18.3	(2)	5,270	82.9	(3)
NRO	2,566	88.3 (2)	3.1	(5)	2,726	33.9	(2)	2,593	18.1	(3)	2,275	86.9	(1)
SERO	14,389	80.1 (5)	2.5	(3)	15,137	32.4	(4)	15,252	13.3	(5)	12,353	80.5	(4)
SRO	5,216	87.9 (3)	2.2	(2)	5,570	30.1	(5)	5,501	17.0	(4)	4,503	85.3	(2)
WRO	4,324	89.8 (1)	1.2	(1)	4,611	38.7	(1)	4,632	21.0	(1)	3,818	78.9	(5)
<b>Wisconsin</b>	<b>32,437</b>	<b>84.0</b>	<b>2.3</b>		<b>34,433</b>	<b>33.0</b>		<b>34,386</b>	<b>16.3</b>		<b>28,219</b>	<b>82.0</b>	
<b>Nation (Prior Year)</b>	<b>612,462</b>	<b>79.4</b>	<b>4.7</b>		<b>1,034,046</b>	<b>31.3</b>		<b>98,170</b>	<b>15.7</b>		<b>238,509</b>	<b>75.9</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7E**  
**2008 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Region**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Region</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>			<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	6,338	43.7 (4)	5,995	24.4 (3)	2,127	30.7 (2)	5,328	21.0 (2)	5,999	7.4 (3)
NRO	2,721	43.6 (3)	2,555	26.1 (4)	931	26.7 (5)	2,295	24.5 (5)	2,560	6.7 (2)
SERO	14,794	28.7 (1)	14,207	14.7 (1)	3,163	35.6 (1)	12,499	22.0 (4)	14,205	9.5 (5)
SRO	5,558	38.7 (2)	5,174	23.7 (2)	1,599	28.6 (3)	4,588	20.4 (1)	5,179	7.4 (3)
WRO	4,558	46.0 (5)	4,259	26.6 (5)	1,673	26.9 (4)	3,909	21.2 (3)	4,261	4.4 (1)
<b>Wisconsin</b>	<b>33,969</b>	<b>36.6</b>	<b>32,190</b>	<b>20.4</b>	<b>9,493</b>	<b>30.9</b>	<b>28,619</b>	<b>21.6</b>	<b>32,204</b>	<b>7.9</b>
<b>Nation (Prior Year)</b>	<b>895,227</b>	<b>26.6</b>	<b>716,629</b>	<b>15.8</b>	<b>161,864</b>	<b>35.1</b>	<b>572,120</b>	<b>20.9</b>	<b>486,636</b>	<b>22.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8E**  
**2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Region**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Region</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
NERO	5,796	7.6 (4)	8.3 (3)		5,834	10.8 (5)	5,020	3.0 (4)	5,879	64.7 (4)
NRO	2,436	7.3 (3)	8.0 (2)		2,483	9.5 (3)	2,144	2.7 (2)	2,482	63.5 (5)
SERO	13,685	8.6 (5)	6.5 (1)		13,815	10.6 (4)	11,850	3.5 (5)	13,859	65.9 (3)
SRO	4,992	6.4 (1)	9.4 (4)		5,041	9.2 (2)	4,390	2.7 (2)	5,083	69.8 (2)
WRO	4,124	6.7 (2)	9.6 (5)		4,171	9.1 (1)	3,644	2.5 (1)	4,196	72.3 (1)
<b>Wisconsin</b>	<b>31,033</b>	<b>7.7</b>	<b>7.8</b>		<b>31,344</b>	<b>10.2</b>	<b>27,048</b>	<b>3.1</b>	<b>31,499</b>	<b>67.0</b>
<b>Nation (Prior Year)</b>	<b>1,039,036</b>	<b>8.2</b>	<b>6.7</b>		<b>947,705</b>	<b>11.9</b>	<b>856,467</b>	<b>3.7</b>	<b>936,194</b>	<b>64.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Region**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Region</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
NERO	18,924	66.4	2.6	16.6	4.8	6.8	2.3	0.5
NRO	8,103	76.8	0.7	5.6	6.2	8.8	1.4	0.5
SERO	44,827	29.7	37.9	27.1	0.4	3.0	1.5	0.4
SRO	16,103	64.8	10.4	18.5	0.5	3.3	1.8	0.6
WRO	13,603	83.8	1.4	6.1	1.8	4.3	1.9	0.8
<b>Wisconsin</b>	<b>101,560</b>	<b>53.1</b>	<b>19.1</b>	<b>19.2</b>	<b>1.9</b>	<b>4.4</b>	<b>1.7</b>	<b>0.5</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Region**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Region</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
NERO	18,924	0.2	5.8	13.9	62.9	15.9	1.1	0.1
NRO	8,103	0.1	4.9	13.5	64.0	16.1	1.3	0.0
SERO	44,827	0.5	7.9	14.0	59.9	16.5	1.2	0.0
SRO	16,103	0.3	5.7	13.2	62.3	17.3	1.1	0.0
WRO	13,603	0.2	4.6	13.6	65.0	15.6	1.1	0.0
<b>Wisconsin</b>	<b>101,560</b>	<b>0.3</b>	<b>6.5</b>	<b>13.8</b>	<b>61.8</b>	<b>16.3</b>	<b>1.1</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Region</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	18,693	10.9 (5)	43.8 (1)	13,660	24.5 (4)	45.3 (4)	3,026	27.4 (3)	15,546	22.6 (4)	15,433	5.3 (5)	15,442	5.9(4)
NRO	8,049	9.7 (2)	45.3 (2)	6,341	24.3 (3)	45.2 (3)	1,305	22.8 (1)	6,622	15.8 (1)	6,639	3.4 (1)	6,660	4.1(1)
SERO	44,309	9.0 (1)	48.8 (5)	32,759	25.8 (5)	44.5 (1)	7,383	37.8 (5)	37,309	30.3 (5)	35,760	4.2 (3)	35,911	6.7(5)
SRO	15,875	10.5 (4)	45.7 (4)	9,280	23.9 (2)	45.0 (2)	2,774	28.7 (4)	12,279	20.9 (2)	12,637	4.1 (2)	12,692	5.5(3)
WRO	13,474	10.2 (3)	45.5 (3)	7,856	22.4 (1)	48.3 (5)	1,893	26.9 (2)	11,088	21.6 (3)	10,864	4.4 (4)	10,844	4.9(2)
<b>Wisconsin</b>	<b>100,400</b>	<b>9.8</b>	<b>46.7</b>	<b>69,896</b>	<b>24.7</b>	<b>45.2</b>	<b>16,381</b>	<b>31.9</b>	<b>82,844</b>	<b>25.2</b>	<b>81,333</b>	<b>4.4</b>	<b>81,549</b>	<b>5.9</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight (BMI <19.8), overweight (BMI >26.0; includes overweight and obese women).  
(5) Based on 1990 IOM report, "Nutrition During Pregnancy": underweight prepregnancy (ideal weight gain = 28 to 40 lbs), normal weight prepregnancy (ideal weight gain = 25 to 35 lbs), overweight prepregnancy (ideal weight gain = 15 to 25 lbs), obese prepregnancy (ideal weight gain = 15 to 25 lbs).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>													
<b>Region</b>	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Pregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)			
NERO	13,916	83.9 (4)	2.1	(3)	17,804	32.8	(3)	12,193	18.6	(3)	10,294	83.4	(3)
NRO	6,332	86.0 (3)	2.5	(5)	7,601	33.1	(2)	4,901	19.6	(2)	4,455	87.6	(1)
SERO	32,727	80.0 (5)	2.1	(3)	41,898	32.0	(4)	28,547	13.7	(5)	23,888	81.0	(5)
SRO	12,634	87.6 (2)	2.0	(2)	15,209	29.0	(5)	10,370	18.4	(4)	8,523	85.8	(2)
WRO	10,016	89.0 (1)	0.9	(1)	12,795	36.5	(1)	8,536	21.7	(1)	7,272	81.2	(4)
<b>Wisconsin</b>	<b>75,625</b>	<b>83.7</b>	<b>2.0</b>		<b>95,307</b>	<b>32.4</b>		<b>64,547</b>	<b>16.9</b>		<b>54,432</b>	<b>82.8</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Region</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	16,486	42.2 (3)	17,210	25.3 (3)	5,555	28.7 (2)	13,195	22.5 (3)	15,787	8.2 (3)
NRO	7,184	42.5 (4)	7,369	26.8 (4)	2,453	25.7 (4)	5,767	23.7 (5)	6,787	7.8 (2)
SERO	38,502	27.8 (1)	40,617	15.5 (1)	8,262	33.5 (1)	31,159	23.1 (4)	37,316	10.7 (5)
SRO	14,003	36.9 (2)	14,465	23.3 (2)	3,946	25.8 (3)	11,056	22.3 (2)	13,264	8.5 (4)
WRO	11,725	44.7 (5)	12,310	27.5 (5)	4,305	24.3 (5)	9,536	22.1 (1)	11,308	5.7 (1)
<b>Wisconsin</b>	<b>87,900</b>	<b>35.4</b>	<b>91,971</b>	<b>21.1</b>	<b>24,521</b>	<b>28.8</b>	<b>70,713</b>	<b>22.8</b>	<b>84,462</b>	<b>9.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8E**  
**2006 - 2008 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>											
<b>Region</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)								
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
NERO	16,521	7.6 (4)	8.1 (2)		16,088	10.8 (4)		13,919	3.1 (4)	16,837	63.4 (4)
NRO	7,026	6.5 (1)	8.5 (3)		6,857	9.1 (1)		6,026	2.6 (1)	7,161	62.3 (5)
SERO	38,676	8.7 (5)	6.5 (1)		37,489	11.1 (5)		32,194	3.5 (5)	39,546	64.2 (3)
SRO	13,919	6.8 (3)	9.2 (4)		13,609	9.5 (2)		11,933	2.7 (3)	14,193	69.3 (2)
WRO	11,878	6.5 (1)	9.3 (5)		11,520	9.5 (2)		10,142	2.6 (1)	12,081	71.3 (1)
<b>Wisconsin</b>	<b>88,020</b>	<b>7.7</b>	<b>7.8</b>		<b>85,563</b>	<b>10.4</b>		<b>74,214</b>	<b>3.1</b>	<b>89,818</b>	<b>65.7</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

