

Table 3B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Adams Cty	144	91.7	0.0	3.5	0.7	0.0	4.2	0.0
Ashland	152	73.7	0.0	2.0	19.1	0.0	5.3	0.0
Barron	296	84.8	5.1	8.4	0.7	0.7	0.3	0.0
Bayfield C	92	*	*	*	*	*	*	*
Brown	1,851	50.2	5.7	25.9	9.1	6.5	2.6	0.0
Buffalo	71	*	*	*	*	*	*	*
Burnett	143	73.4	0.0	2.1	21.0	0.0	3.5	0.0
Calumet	167	70.1	2.4	20.4	0.0	6.6	0.6	0.0
Chippewa	340	95.6	0.0	0.9	0.0	1.5	2.1	0.0
Clark	206	76.7	0.5	20.4	0.5	0.0	1.9	0.0
Columbia	234	88.0	2.1	8.5	0.9	0.0	0.4	0.0
Crawford C	95	*	*	*	*	*	*	*
Dane	2,142	42.3	21.1	25.5	0.7	7.9	2.5	0.0
Dodge	376	78.5	0.3	17.8	0.5	1.1	1.9	0.0
Door	148	83.8	0.0	12.8	1.4	0.7	1.4	0.0
Douglas Ct	326	88.7	1.2	2.5	1.5	1.2	4.9	0.0
Dunn	223	84.8	0.9	3.6	0.0	9.9	0.9	0.0
Eau Claire	619	79.6	1.3	5.7	1.5	9.9	2.1	0.0
Florence	40	*	*	*	*	*	*	*
Fond du La	570	75.6	3.0	15.8	0.5	3.9	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Forest	77	*	*	*	*	*	*	*
Grant Cty	277	92.1	1.4	5.8	0.0	0.0	0.7	0.0
Green Cty	186	89.8	0.0	8.6	0.0	0.0	1.6	0.0
Green Lake	91	*	*	*	*	*	*	*
Iowa Cty	99	*	*	*	*	*	*	*
Iron Cty	28	*	*	*	*	*	*	*
Jackson	175	77.7	0.6	5.1	14.9	0.0	1.7	0.0
Jefferson	476	64.9	0.6	32.6	0.2	0.4	1.3	0.0
Juneau	193	91.2	0.5	3.6	1.0	0.5	3.1	0.0
Kenosha	1,297	49.6	16.4	30.5	0.5	1.2	1.9	0.0
Kewaunee	95	*	*	*	*	*	*	*
La Crosse	677	73.4	4.1	4.0	0.3	16.1	2.1	0.0
Lafayette	52	*	*	*	*	*	*	*
Langlade C	107	91.6	0.0	7.5	0.9	0.0	0.0	0.0
Lincoln Ct	136	95.6	0.0	3.7	0.0	0.0	0.7	0.0
Manitowoc	429	76.5	1.9	11.7	0.7	7.7	1.6	0.0
Marathon C	801	65.4	1.2	7.7	0.5	24.2	0.9	0.0
Marinette	256	96.5	0.0	3.1	0.4	0.0	0.0	0.0
Marquette	72	*	*	*	*	*	*	*
Menominee	121	5.0	0.0	9.1	26.4	0.0	59.5	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Milwaukee	10,703	18.1	49.7	25.6	0.7	4.3	1.6	0.0
Monroe	304	84.2	1.3	11.5	0.3	1.0	1.6	0.0
Oconto	194	96.4	0.0	1.5	1.0	0.5	0.5	0.0
Oneida	174	93.1	0.0	4.6	0.0	0.0	2.3	0.0
Outagamie	766	71.4	2.6	13.6	0.9	10.4	1.0	0.0
Ozaukee	147	81.0	5.4	8.2	0.0	2.7	2.7	0.0
Pepin	51	*	*	*	*	*	*	*
Pierce	171	94.7	0.0	2.3	0.6	0.6	1.8	0.0
Polk	276	95.7	0.0	2.9	0.4	0.4	0.7	0.0
Portage	360	77.2	0.3	8.6	0.8	12.2	0.8	0.0
Price	105	92.4	0.0	4.8	1.0	1.9	0.0	0.0
Racine Cou	1,493	41.2	27.3	28.5	0.1	0.3	2.6	0.0
Richland C	115	84.3	0.0	13.9	0.0	1.7	0.0	0.0
Rock	1,134	66.8	10.8	20.3	0.2	0.9	1.0	0.0
Rusk	110	96.4	0.0	2.7	0.0	0.0	0.9	0.0
Sauk	423	80.6	0.5	13.5	5.2	0.2	0.0	0.0
Sawyer	156	54.5	0.6	4.5	37.2	0.6	2.6	0.0
Shawano	218	77.1	0.5	4.1	15.1	0.0	3.2	0.0
Sheboygan	615	59.2	2.1	18.2	0.2	20.0	0.3	0.0
St. Croix	330	87.0	0.9	7.0	1.2	3.0	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Taylor	122	88.5	0.8	7.4	1.6	0.0	1.6	0.0
Trempealea	212	67.0	0.0	31.1	0.0	0.5	1.4	0.0
Vernon	177	93.2	0.6	5.1	0.0	1.1	0.0	0.0
Vilas	152	55.3	0.0	0.7	40.1	0.0	3.9	0.0
Walworth	547	57.2	1.3	38.2	1.1	0.7	1.5	0.0
Washburn	113	96.5	0.9	1.8	0.0	0.0	0.9	0.0
Washington	436	84.9	0.9	8.9	0.7	1.6	3.0	0.0
Waukesha	715	68.5	3.6	24.9	0.0	1.4	1.5	0.0
Waupaca	270	89.3	0.4	9.3	0.0	0.4	0.7	0.0
Waushara C	147	70.1	0.0	28.6	0.0	0.0	1.4	0.0
Winnebago	844	76.2	3.6	11.3	0.7	6.5	1.8	0.0
Wood	436	82.8	0.2	10.1	1.6	3.7	1.6	0.0
Wisconsin	35,896	53.7	19.1	18.9	1.9	4.5	1.9	0.0
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams Cty	144	0.0	5.6	19.4	59.7	14.6	0.7	0.0
Ashland	152	0.0	3.9	11.8	63.8	16.4	3.9	0.0
Barron	296	0.0	2.4	13.5	64.2	18.2	1.7	0.0
Bayfield C	92	*	*	*	*	*	*	*
Brown	1,851	0.5	6.1	12.7	62.0	17.1	1.3	0.2
Buffalo	71	*	*	*	*	*	*	*
Burnett	143	0.0	4.2	18.2	67.1	9.8	0.7	0.0
Calumet	167	0.0	3.6	13.2	64.1	17.4	1.8	0.0
Chippewa	340	0.0	2.6	12.4	67.9	15.6	1.5	0.0
Clark	206	0.0	5.3	14.1	64.6	14.6	1.5	0.0
Columbia	234	0.0	2.1	14.1	65.0	17.5	1.3	0.0
Crawford C	95	*	*	*	*	*	*	*
Dane	2,142	0.4	4.5	11.4	60.2	21.7	1.7	0.0
Dodge	376	0.0	2.1	12.2	65.2	18.6	1.9	0.0
Door	148	0.0	4.1	10.8	60.1	23.6	1.4	0.0
Douglas Ct	326	0.0	3.7	13.8	68.1	13.5	0.9	0.0
Dunn	223	0.0	4.9	10.8	64.6	18.4	1.3	0.0
Eau Claire	619	0.2	4.0	12.0	66.2	16.2	1.5	0.0
Florence	40	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Fond du La	570	0.4	3.0	17.5	60.0	18.2	0.9	0.0
Forest	77	*	*	*	*	*	*	*
Grant Cty	277	0.0	2.2	14.8	63.5	19.1	0.4	0.0
Green Cty	186	0.5	5.4	14.0	64.5	15.1	0.5	0.0
Green Lake	91	*	*	*	*	*	*	*
Iowa Cty	99	*	*	*	*	*	*	*
Iron Cty	28	*	*	*	*	*	*	*
Jackson	175	0.6	4.6	11.4	67.4	16.0	0.0	0.0
Jefferson	476	0.0	2.5	11.1	61.1	23.7	1.5	0.0
Juneau	193	0.0	4.1	9.3	63.7	20.7	2.1	0.0
Kenosha	1,297	0.4	6.3	15.3	59.2	17.7	1.0	0.0
Kewaunee	95	*	*	*	*	*	*	*
La Crosse	677	0.1	4.1	13.3	63.1	18.0	1.3	0.0
Lafayette	52	*	*	*	*	*	*	*
Langlade C	107	0.0	10.3	8.4	59.8	21.5	0.0	0.0
Lincoln Ct	136	0.7	4.4	19.9	59.6	14.0	1.5	0.0
Manitowoc	429	0.2	4.9	10.7	64.8	17.9	1.4	0.0
Marathon C	801	0.2	4.5	12.2	64.8	17.0	1.2	0.0
Marinette	256	0.0	5.5	10.5	64.5	18.4	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marquette	72	*	*	*	*	*	*	*
Menominee	121	0.0	9.9	16.5	54.5	19.0	0.0	0.0
Milwaukee	10,703	0.4	7.6	13.2	58.9	18.6	1.3	0.0
Monroe	304	0.3	5.9	13.5	65.8	13.8	0.7	0.0
Oconto	194	0.0	4.1	17.0	61.3	17.0	0.5	0.0
Oneida	174	0.0	4.6	10.3	66.7	17.8	0.6	0.0
Outagamie	766	0.1	5.0	12.7	64.5	16.8	0.9	0.0
Ozaukee	147	0.0	2.0	8.2	63.3	23.1	3.4	0.0
Pepin	51	*	*	*	*	*	*	*
Pierce	171	0.0	3.5	9.9	66.1	20.5	0.0	0.0
Polk	276	0.0	4.0	13.4	63.0	17.8	1.8	0.0
Portage	360	0.0	3.3	11.1	66.4	18.3	0.8	0.0
Price	105	0.0	1.0	9.5	66.7	22.9	0.0	0.0
Racine Cou	1,493	0.6	6.4	12.0	63.2	16.6	1.1	0.1
Richland C	115	0.0	1.7	10.4	67.8	19.1	0.9	0.0
Rock	1,134	0.2	6.3	17.3	59.1	15.9	1.1	0.1
Rusk	110	0.0	5.5	16.4	60.9	15.5	1.8	0.0
Sauk	423	0.0	2.8	14.9	67.4	13.9	0.9	0.0
Sawyer	156	0.6	3.8	13.5	62.2	19.2	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Shawano	218	0.0	4.1	10.6	63.3	21.1	0.9	0.0
Sheboygan	615	0.0	6.2	13.2	62.0	17.4	1.3	0.0
St. Croix	330	0.0	2.1	8.5	66.4	21.5	1.5	0.0
Taylor	122	0.0	4.1	12.3	59.8	23.0	0.8	0.0
Trempealea	212	0.5	4.7	9.0	65.1	18.9	1.9	0.0
Vernon	177	0.0	6.8	11.3	59.3	22.0	0.6	0.0
Vilas	152	0.0	2.6	19.7	61.8	14.5	1.3	0.0
Walworth	547	0.2	4.0	10.1	63.4	21.4	0.9	0.0
Washburn	113	0.0	3.5	12.4	61.9	21.2	0.9	0.0
Washington	436	0.0	3.4	12.4	64.0	19.3	0.9	0.0
Waukesha	715	0.3	4.5	11.5	61.7	20.3	1.8	0.0
Waupaca	270	0.4	4.8	14.8	62.2	17.0	0.7	0.0
Waushara C	147	0.0	4.8	13.6	63.3	18.4	0.0	0.0
Winnebago	844	0.1	3.8	13.7	64.3	16.9	1.1	0.0
Wood	436	0.0	2.3	14.0	66.1	16.5	1.1	0.0
Wisconsin	35,896	0.3	5.5	13.0	61.8	18.3	1.2	0.0
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	140	5.0 (49)	52.9 (20)	115	26.1 (46)	50.4 (22)	15	* *	111	15.3 (9)	133	5.3 (41)	133	11.3(59)
Ashland	151	1.3 (2)	55.0 (33)	135	25.2 (42)	47.4 (6)	17	* *	116	19.8(29)	136	8.8 (58)	136	5.9(38)
Barron	291	3.1 (18)	64.3 (60)	246	19.5 (20)	52.0 (27)	36	* *	220	21.4(36)	263	8.4 (56)	263	3.0(10)
Bayfield C	92	* *	* *	80	* *	* *	7	* *	64	* *	85	* *	85	* *
Brown	1,817	5.1 (52)	52.6 (19)	1,302	21.7 (29)	49.5 (16)	287	34.8 (10)	1,500	23.6(42)	1,689	5.1 (36)	1,689	6.6(45)
Buffalo	71	* *	* *	67	* *	* *	12	* *	60	* *	66	* *	66	* *
Burnett	142	3.5 (25)	62.7 (59)	91	* *	* *	13	* *	112	20.5(31)	127	3.1 (11)	127	0.0(1)
Calumet	167	6.0 (55)	47.3 (1)	152	19.1 (18)	48.7 (10)	27	* *	138	23.9(43)	158	5.1 (36)	158	5.7(36)
Chippewa	339	3.5 (25)	56.6 (41)	316	19.0 (17)	51.9 (26)	53	* *	297	22.2(40)	319	5.0 (35)	319	3.4(15)
Clark	205	4.9 (47)	57.1 (45)	194	25.8 (45)	48.5 (8)	48	* *	176	34.1(53)	195	3.1 (11)	195	4.1(22)
Columbia	233	4.7 (45)	54.1 (25)	193	15.5 (2)	59.6 (50)	30	* *	190	20.5(31)	208	6.7 (52)	208	4.8(25)
Crawford C	95	* *	* *	71	* *	* *	19	* *	70	* *	85	* *	85	* *
Dane	2,110	3.6 (28)	51.9 (17)	1,463	20.8 (24)	51.4 (24)	432	23.6 (3)	1,547	21.7(39)	1,911	6.0 (47)	1,911	6.0(41)
Dodge	373	4.0 (33)	50.1 (7)	306	19.6 (21)	49.0 (12)	66	* *	315	19.0(26)	343	5.2 (39)	343	5.2(28)
Door	148	4.7 (45)	54.1 (25)	138	18.8 (13)	54.3 (35)	26	* *	121	14.0 (3)	138	5.1 (36)	138	5.8(37)
Douglas Ct	325	3.4 (24)	52.9 (20)	236	17.8 (9)	54.7 (40)	49	* *	239	25.5(46)	285	5.3 (41)	285	5.6(35)
Dunn	218	5.0 (49)	57.3 (47)	113	30.1 (50)	43.4 (3)	26	* *	148	16.2(14)	192	3.1 (11)	192	4.7(24)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)															
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Eau Claire	617	5.2 (53)	47.6 (3)	41	*	*	83	*	*	516	19.4(27)	575	5.2 (39)	575	4.0(20)
Florence	39	*	*	28	*	*	4	*	*	28	*	32	*	32	*
Fond du La	557	3.2 (19)	53.0 (22)	501	22.6 (33)	46.1 (5)	88	*	*	468	19.7(28)	519	4.6 (29)	519	7.1(48)
Forest	75	*	*	65	*	*	7	*	*	47	*	71	*	71	*
Grant Cty	272	2.6 (9)	61.8 (58)	227	21.6 (27)	57.3 (47)	45	*	*	214	15.9(11)	257	4.3 (23)	257	8.2(57)
Green Cty	183	1.6 (6)	54.6 (30)	166	12.7 (1)	57.8 (48)	37	*	*	161	28.0(49)	166	2.4 (7)	166	3.0(10)
Green Lake	89	*	*	82	*	*	15	*	*	66	*	79	*	79	*
Iowa Cty	99	*	*	78	*	*	16	*	*	79	*	92	*	92	*
Iron Cty	27	*	*	25	*	*	2	*	*	17	*	26	*	26	*
Jackson	175	7.4 (60)	55.4 (34)	155	27.1 (49)	41.3 (1)	19	*	*	135	25.2(45)	156	3.8 (19)	156	5.1(27)
Jefferson	471	4.2 (34)	55.6 (36)	380	20.8 (24)	52.4 (30)	90	*	*	366	12.8 (1)	435	4.8 (32)	435	7.6(52)
Juneau	192	3.6 (28)	59.4 (50)	160	21.3 (26)	58.1 (49)	25	*	*	161	14.3 (5)	183	4.4 (26)	183	10.9(58)
Kenosha	1,271	2.7 (10)	56.9 (43)	1,188	20.0 (22)	50.8 (23)	237	32.1 (8)	1,055	28.4(50)	1,170	5.5 (44)	1,170	7.7(54)	
Kewaunee	95	*	*	83	*	*	12	*	*	78	*	89	*	89	*
La Crosse	675	4.3 (38)	51.0 (12)	170	26.5 (48)	49.4 (14)	101	20.8 (2)	508	29.5(51)	600	4.2 (21)	600	1.7(4)	
Lafayette	51	*	*	32	*	*	8	*	*	35	*	42	*	42	*
Langlade C	106	2.8 (11)	60.4 (55)	97	*	*	11	*	*	90	*	102	2.0 (3)	102	2.0(6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lincoln Ct	135	1.5 (5)	60.0 (52)	98	* *	* *	14	* *	117	22.2(40)	131	8.4 (56)	131	3.8(18)
Manitowoc	425	5.6 (54)	50.8 (11)	164	31.1 (51)	49.4 (14)	61	* *	356	16.0(13)	384	4.2 (21)	384	5.2(28)
Marathon C	797	4.3 (38)	51.8 (16)	613	23.2 (36)	50.2 (21)	132	28.8 (6)	678	21.4(36)	736	3.3 (14)	736	1.4(3)
Marinette	253	6.7 (59)	47.4 (2)	181	22.1 (32)	55.2 (41)	33	* *	150	20.7(33)	226	7.1 (54)	226	5.3(30)
Marquette	69	* *	* *	60	* *	* *	13	* *	48	* *	65	* *	65	* *
Menominee	121	3.3 (22)	60.3 (54)	107	26.2 (47)	49.5 (16)	9	* *	101	16.8(17)	110	3.6 (17)	110	5.5(33)
Milwaukee	10,591	3.2 (19)	57.1 (45)	7,824	23.6 (37)	49.2 (13)	1,685	42.7 (12)	8,844	33.9(52)	9,756	4.3 (23)	9,756	6.9(47)
Monroe	301	4.3 (38)	49.8 (4)	7	* *	* *	51	* *	248	21.0(34)	279	4.7 (30)	279	1.8(5)
Oconto	194	4.6 (42)	51.5 (13)	183	16.4 (5)	59.6 (50)	33	* *	56	* *	183	4.4 (26)	183	7.1(48)
Oneida	174	4.6 (42)	55.7 (37)	114	23.7 (38)	55.3 (42)	21	* *	140	14.3 (5)	158	4.4 (26)	158	3.8(18)
Outagamie	753	4.6 (42)	50.2 (8)	614	16.3 (4)	55.5 (43)	122	30.3 (7)	643	25.7(47)	692	5.6 (46)	692	6.5(44)
Ozaukee	146	6.2 (56)	50.0 (6)	126	15.9 (3)	55.6 (44)	34	* *	112	17.9(21)	127	9.4 (59)	127	6.3(42)
Pepin	51	* *	* *	48	* *	* *	7	* *	42	* *	48	* *	48	* *
Pierce	171	2.9 (15)	56.7 (42)	135	17.0 (7)	61.5 (52)	15	* *	138	15.9(11)	152	6.6 (51)	152	7.9(56)
Polk	276	3.6 (28)	60.1 (53)	206	22.8 (34)	50.0 (20)	31	* *	228	16.2(14)	252	7.1 (54)	252	7.1(48)
Portage	356	4.2 (34)	54.8 (31)	127	22.8 (34)	52.0 (27)	59	* *	282	12.8 (1)	325	2.2 (6)	325	5.5(33)
Price	104	2.9 (15)	59.6 (51)	98	* *	* *	10	* *	92	* *	99	* *	99	* *

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)															
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy		
	Under- weight	Over- weight		< Ideal	> Ideal		3rd Trimester		Postpartum						
		Number	%		%	Number	%	%	Number	%	Number	%	Number	%	Number
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)
Racine Cou	1,477	2.8 (11)	58.6 (48)	1,343	21.7 (29)	52.0 (27)	242	33.9 (9)	1,191	24.1 (44)	1,353	4.7 (30)	1,353	5.9 (38)	
Richland C	115	1.7 (7)	59.1 (49)	89	* *	* *	15	* *	91	* *	104	0.0 (1)	104	2.9 (7)	
Rock	1,104	3.8 (32)	56.3 (39)	815	17.9 (11)	48.6 (9)	198	34.8 (10)	915	26.6 (48)	1,021	2.0 (3)	1,021	3.0 (10)	
Rusk	110	3.6 (28)	54.5 (29)	101	25.7 (44)	51.5 (25)	9	* *	82	* *	104	1.0 (2)	104	4.8 (25)	
Sauk	409	3.2 (19)	54.8 (31)	365	17.8 (9)	54.5 (36)	55	* *	338	14.8 (8)	380	2.9 (8)	380	2.9 (7)	
Sawyer	156	1.3 (2)	56.4 (40)	116	24.1 (39)	56.0 (45)	25	* *	105	17.1 (19)	140	2.1 (5)	140	2.9 (7)	
Shawano	216	4.2 (34)	61.6 (57)	178	24.2 (40)	48.9 (11)	25	* *	92	* *	204	6.4 (49)	204	5.9 (38)	
Sheboygan	613	2.8 (11)	50.7 (9)	550	24.9 (41)	43.1 (2)	101	27.7 (5)	529	18.3 (24)	576	4.9 (33)	576	3.1 (13)	
St. Croix	322	0.6 (1)	54.3 (27)	288	21.9 (31)	48.3 (7)	45	* *	193	14.0 (3)	281	5.3 (41)	281	7.8 (55)	
Taylor	122	6.6 (58)	53.3 (24)	112	17.9 (11)	54.5 (36)	16	* *	100	18.0 (22)	114	7.0 (53)	114	5.3 (30)	
Trempealea	212	2.8 (11)	56.1 (38)	120	34.2 (52)	45.8 (4)	35	* *	161	21.1 (35)	187	3.7 (18)	187	4.3 (23)	
Vernon	176	4.5 (41)	51.7 (15)	117	18.8 (13)	53.0 (33)	25	* *	127	17.3 (20)	165	3.0 (10)	165	7.3 (51)	
Vilas	152	1.3 (2)	61.2 (56)	97	* *	* *	20	* *	112	14.3 (5)	141	4.3 (23)	141	3.5 (16)	
Walworth	542	3.3 (22)	55.5 (35)	283	21.6 (27)	49.8 (19)	88	* *	465	21.5 (38)	501	3.4 (16)	501	3.2 (14)	
Washburn	113	3.5 (25)	53.1 (23)	101	18.8 (13)	54.5 (36)	11	* *	95	* *	102	2.9 (8)	102	1.0 (2)	
Washington	432	2.3 (8)	50.7 (9)	391	17.1 (8)	52.7 (31)	61	* *	383	19.8 (29)	406	6.2 (48)	406	7.6 (52)	
Waukesha	709	4.2 (34)	51.5 (13)	625	19.2 (19)	54.6 (39)	113	27.4 (4)	485	18.8 (25)	651	5.5 (44)	651	6.6 (45)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Waupaca	267	6.4 (57)	49.8 (4)	237	20.3 (23)	53.2 (34)	34	* *	182	18.1 (23)	250	4.0 (20)	250	6.4 (43)
Waushara C	138	2.9 (15)	54.3 (27)	123	25.2 (42)	49.6 (18)	22	* *	111	16.2 (14)	124	6.5 (50)	124	4.0 (20)
Winnebago	839	5.0 (49)	52.4 (18)	719	16.4 (5)	56.6 (46)	166	20.5 (1)	670	16.9 (18)	793	4.9 (33)	793	5.4 (32)
Wood	430	4.9 (47)	57.0 (44)	383	18.8 (13)	52.7 (31)	98	* *	319	15.4 (10)	395	3.3 (14)	395	3.5 (16)
Wisconsin	35,482	3.7	55.1	26,554	21.6	50.9	5,697	31.9	28,469	25.0	32,642	4.6	32,642	5.8
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4	395,960	5.2

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)												
County	Medical Care				WIC Enrollment			Multivitamin Consumption				
	1st Trimester		No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)	Number	% (Rank)
Adams Cty	137	95.6 (2)	0.7	(11)	144	52.8	(2)	141	20.6	(29)	117	88.9 (22)
Ashland	141	85.1 (51)	2.8	(43)	152	46.7	(7)	149	29.5	(4)	137	94.9 (1)
Barron	274	93.8 (7)	0.7	(11)	296	49.7	(4)	290	23.4	(22)	238	86.6 (30)
Bayfield C	86	* *	* *		92	* *		91	* *		78	* *
Brown	1,685	79.9 (59)	4.3	(52)	1,841	37.2	(26)	1,739	16.7	(48)	1,482	79.6 (52)
Buffalo	68	* *	* *		71	* *		70	* *		49	* *
Burnett	130	92.3 (14)	2.3	(41)	140	45.7	(11)	140	21.4	(26)	120	83.3 (45)
Calumet	155	88.4 (38)	0.6	(8)	167	29.3	(50)	167	18.0	(43)	119	90.8 (12)
Chippewa	325	93.5 (9)	0.0	(1)	340	36.8	(29)	338	25.4	(12)	275	83.6 (43)
Clark	199	86.4 (46)	1.5	(28)	206	23.8	(58)	205	26.3	(8)	165	90.9 (11)
Columbia	210	83.8 (55)	1.0	(19)	234	32.1	(45)	232	38.4	(1)	185	86.5 (31)
Crawford C	88	* *	* *		95	* *		95	* *		75	* *
Dane	1,937	89.4 (32)	1.1	(22)	2,138	27.7	(55)	1,967	16.0	(49)	1,656	86.5 (31)
Dodge	357	89.6 (28)	2.0	(38)	375	32.5	(43)	371	24.5	(15)	289	86.9 (29)
Door	140	87.9 (41)	2.1	(39)	148	38.5	(22)	145	31.7	(3)	124	91.9 (4)
Douglas Ct	297	89.9 (26)	1.7	(33)	326	42.3	(14)	315	20.3	(31)	271	89.7 (20)
Dunn	205	93.7 (8)	1.5	(28)	223	46.2	(8)	213	25.4	(12)	185	87.6 (28)
Eau Claire	586	86.2 (48)	1.4	(26)	617	36.0	(34)	612	19.4	(36)	488	83.2 (46)
Florence	33	* *	* *		40	* *		39	* *		31	* *
Fond du La	535	86.0 (49)	4.1	(50)	570	37.2	(26)	555	17.5	(44)	456	81.6 (50)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Forest	72	* *	* *	* *	77	* *	76	* *	59	* *
Grant Cty	266	92.1 (17)	2.6 (42)		276	33.3 (41)	271	24.4 (16)	213	91.5 (7)
Green Cty	169	89.3 (35)	1.2 (23)		185	37.8 (24)	185	22.2 (24)	155	90.3 (15)
Green Lake	84	* *	* *	* *	91	* *	86	* *	65	* *
Iowa Cty	94	* *	* *	* *	99	* *	98	* *	77	* *
Iron Cty	25	* *	* *	* *	28	* *	25	* *	22	* *
Jackson	162	91.4 (22)	0.0 (1)		175	40.0 (19)	174	10.9 (59)	141	82.3 (48)
Jefferson	426	88.3 (39)	1.6 (31)		474	31.4 (46)	449	20.0 (33)	367	82.0 (49)
Juneau	189	92.1 (17)	0.5 (7)		193	36.3 (31)	190	21.6 (25)	153	90.2 (16)
Kenosha	1,218	76.0 (60)	8.6 (60)		1,294	28.7 (53)	1,246	20.3 (31)	932	85.9 (34)
Kewaunee	90	* *	* *	* *	95	* *	93	* *	60	* *
La Crosse	616	92.5 (12)	1.9 (36)		674	36.1 (32)	662	20.4 (30)	513	59.6 (56)
Lafayette	40	* *	* *	* *	52	* *	50	* *	37	* *
Langlade C	104	92.3 (14)	1.0 (19)		107	41.1 (18)	107	18.7 (40)	91	* *
Lincoln Ct	128	93.0 (10)	1.6 (31)		135	32.6 (42)	136	16.9 (47)	105	92.4 (3)
Manitowoc	397	84.4 (52)	6.0 (59)		427	43.3 (13)	420	23.3 (23)	362	84.3 (42)
Marathon C	748	86.4 (46)	4.5 (55)		801	31.3 (47)	776	13.4 (53)	639	84.4 (41)
Marinette	231	86.6 (45)	4.3 (52)		255	38.0 (23)	250	26.8 (7)	187	84.5 (40)
Marquette	64	* *	* *	* *	70	* *	69	* *	53	* *
Menominee	113	86.7 (44)	0.9 (17)		121	48.8 (6)	118	11.0 (58)	99	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Milwaukee	9,986	81.3 (58)	3.1 (48)	10,654	34.4 (38)	10,491	12.0 (56)	8,435	80.6 (51)
Monroe	281	88.6 (37)	1.4 (26)	303	29.4 (49)	288	17.4 (45)	224	77.2 (54)
Oconto	185	85.4 (50)	2.2 (40)	194	41.2 (17)	192	21.4 (26)	155	87.7 (27)
Oneida	163	93.9 (5)	0.6 (8)	173	39.9 (20)	166	29.5 (4)	132	90.2 (16)
Outagamie	633	83.3 (56)	0.8 (14)	763	32.2 (44)	746	19.2 (38)	584	82.9 (47)
Ozaukee	131	87.0 (43)	0.8 (14)	147	29.9 (48)	145	21.4 (26)	113	91.2 (9)
Pepin	49	* *	* *	51	* *	51	* *	43	* *
Pierce	157	93.0 (10)	1.3 (25)	170	45.9 (9)	166	24.1 (18)	137	89.1 (21)
Polk	258	89.5 (31)	4.3 (52)	275	39.3 (21)	269	25.7 (10)	205	90.7 (13)
Portage	337	89.6 (28)	0.6 (8)	360	37.8 (24)	224	19.2 (38)	292	88.7 (24)
Price	100	91.0 (23)	5.0 (57)	104	59.6 (1)	105	23.8 (21)	88	* *
Racine Cou	1,364	83.0 (57)	3.1 (48)	1,490	36.0 (34)	1,455	15.0 (52)	1,191	84.6 (39)
Richland C	106	94.3 (4)	0.0 (1)	115	35.7 (37)	115	24.3 (17)	91	* *
Rock	1,054	89.0 (36)	4.6 (56)	1,133	28.9 (52)	1,090	12.8 (54)	910	86.3 (33)
Rusk	105	92.4 (13)	2.9 (45)	110	51.8 (3)	109	25.7 (10)	102	90.2 (16)
Sauk	384	90.6 (24)	1.8 (34)	423	37.1 (28)	413	18.6 (41)	325	88.6 (25)
Sawyer	141	89.4 (32)	2.8 (43)	155	45.8 (10)	147	17.0 (46)	133	85.0 (37)
Shawano	206	91.7 (20)	1.9 (36)	217	33.6 (39)	215	28.4 (6)	167	91.0 (10)
Sheboygan	589	84.0 (54)	2.9 (45)	614	21.5 (60)	608	18.4 (42)	474	83.5 (44)
St. Croix	298	89.9 (26)	1.0 (19)	329	41.6 (16)	319	24.1 (18)	240	88.8 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care		1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	113	90.3 (25)	0.0	(1)	122	41.8 (15)	121	33.9 (2)	103	91.3 (8)
Trempealea	189	87.8 (42)	4.2	(51)	209	36.4 (30)	207	11.1 (57)	166	66.3 (55)
Vernon	152	94.7 (3)	0.0	(1)	176	44.9 (12)	175	2.9 (60)	142	78.9 (53)
Vilas	142	88.0 (40)	0.7	(11)	151	29.1 (51)	150	20.0 (33)	120	91.7 (5)
Walworth	509	84.1 (53)	5.5	(58)	542	33.6 (39)	540	25.4 (12)	409	87.8 (26)
Washburn	106	98.1 (1)	0.9	(17)	113	49.6 (5)	113	23.9 (20)	100	90.0 (19)
Washington	417	92.3 (14)	1.2	(23)	436	36.0 (34)	428	26.2 (9)	320	93.4 (2)
Waukesha	659	89.4 (32)	0.2	(6)	713	25.7 (56)	703	19.5 (35)	502	90.4 (14)
Waupaca	251	91.6 (21)	0.8	(14)	269	27.9 (54)	257	12.5 (55)	184	85.3 (35)
Waushara C	132	93.9 (5)	3.0	(47)	147	36.1 (32)	135	15.6 (50)	108	91.7 (5)
Winnebago	800	89.6 (28)	1.5	(28)	842	25.5 (57)	826	19.4 (36)	658	85.3 (35)
Wood	389	91.8 (19)	1.8	(34)	434	21.9 (59)	432	15.3 (51)	344	84.9 (38)
Wisconsin	33,210	85.6	2.8		35,778	34.1	34,756	17.3	28,067	83.8
Nation (Prior Year)	580,485	80.1	3.1		1,187,387	31.7	266,567	19.3	269,548	80.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	140	51.4 (53)	133	33.1 (54)	60	* *	122	35.2 (56)	133	14.3 (59)
Ashland	147	50.3 (50)	136	31.6 (50)	62	* *	139	23.0 (39)	136	5.1 (26)
Barron	286	47.9 (41)	263	28.5 (38)	106	15.1 (26)	244	23.8 (42)	263	3.8 (16)
Bayfield C	90	* *	85	* *	40	* *	78	* *	85	* *
Brown	1,757	38.8 (16)	1,681	19.6 (12)	533	32.3 (11)	1,494	19.6 (25)	1,682	9.3 (50)
Buffalo	67	* *	66	* *	21	* *	50	* *	66	* *
Burnett	138	51.4 (53)	127	34.6 (57)	59	* *	121	29.8 (52)	127	9.4 (51)
Calumet	164	37.8 (13)	157	19.7 (13)	42	* *	121	18.2 (20)	158	2.5 (4)
Chippewa	334	53.3 (56)	319	29.2 (40)	142	39.4 (3)	279	16.5 (12)	319	3.4 (10)
Clark	200	40.5 (18)	195	21.5 (18)	70	* *	168	22.0 (33)	195	6.7 (37)
Columbia	231	55.4 (58)	208	33.7 (55)	92	* *	187	21.4 (31)	208	8.7 (46)
Crawford C	91	* *	85	* *	27	* *	76	* *	85	* *
Dane	2,054	31.5 (4)	1,908	15.9 (4)	446	41.9 (2)	1,680	15.2 (8)	1,908	3.9 (20)
Dodge	367	45.5 (33)	343	25.9 (35)	129	29.5 (14)	291	23.7 (41)	343	6.7 (37)
Door	140	50.7 (52)	137	19.7 (13)	57	* *	121	15.7 (11)	137	3.6 (11)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Douglas Ct	303	54.8 (57)	283	31.1 (48)	132	26.5 (20)	276	26.8 (47)	285	9.5 (52)
Dunn	214	43.0 (24)	192	25.0 (30)	70	* *	186	13.4 (2)	192	3.6 (11)
Eau Claire	593	44.5 (29)	575	25.9 (35)	211	28.4 (17)	495	15.2 (8)	575	2.6 (5)
Florence	36	* *	32	* *	12	* *	31	* *	32	* *
Fond du La	547	44.8 (30)	519	25.8 (34)	192	25.5 (22)	461	20.8 (30)	518	5.6 (29)
Forest	76	* *	71	* *	32	* *	59	* *	71	* *
Grant Cty	257	40.5 (18)	257	24.1 (27)	87	* *	216	18.5 (21)	257	7.4 (41)
Green Cty	181	44.8 (30)	166	24.1 (27)	63	* *	155	20.6 (29)	166	7.8 (42)
Green Lake	85	* *	79	* *	29	* *	66	* *	79	* *
Iowa Cty	95	* *	92	* *	32	* *	77	* *	92	* *
Iron Cty	25	* *	24	* *	9	* *	21	* *	25	* *
Jackson	170	48.2 (44)	156	29.5 (41)	65	* *	147	20.4 (27)	156	3.8 (16)
Jefferson	457	36.5 (10)	434	22.1 (19)	130	28.5 (16)	370	17.3 (15)	435	6.0 (32)
Juneau	189	49.2 (47)	183	33.9 (56)	78	* *	157	24.8 (46)	183	9.8 (54)
Kenosha	1,140	33.4 (7)	1,168	18.5 (10)	276	30.4 (13)	934	20.2 (26)	1,170	5.3 (27)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee	92	* *	89	* *	33	* *	65	* *	89	* *
La Crosse	655	38.2 (14)	597	18.4 (9)	178	28.1 (18)	541	14.2 (4)	598	2.8 (7)
Lafayette	46	* *	42	* *	14	* *	37	* *	42	* *
Langlade C	106	49.1 (46)	102	35.3 (58)	40	* *	91	* *	102	13.7 (57)
Lincoln Ct	133	44.4 (28)	131	25.2 (31)	45	* *	108	23.1 (40)	131	3.1 (8)
Manitowoc	413	47.9 (41)	384	28.9 (39)	162	26.5 (20)	362	22.9 (37)	384	5.5 (28)
Marathon C	776	36.1 (9)	734	23.0 (24)	210	27.1 (19)	646	18.0 (19)	736	3.8 (16)
Marinette	238	47.5 (40)	226	31.0 (47)	90	* *	199	27.1 (48)	226	4.9 (25)
Marquette	71	* *	65	* *	26	* *	53	* *	65	* *
Menominee	115	65.2 (60)	110	32.7 (52)	62	* *	102	31.4 (53)	110	4.5 (23)
Milwaukee	10,023	23.9 (1)	9,753	12.3 (1)	1,798	37.9 (5)	8,500	21.4 (31)	9,753	9.1 (49)
Monroe	279	45.2 (32)	279	29.7 (42)	96	* *	227	22.9 (37)	279	6.8 (39)
Oconto	188	47.3 (38)	183	32.2 (51)	75	* *	155	28.4 (51)	183	9.8 (54)
Oneida	165	51.5 (55)	157	30.6 (45)	65	* *	135	27.4 (50)	157	6.4 (36)
Outagamie	718	46.0 (35)	692	21.4 (17)	249	20.1 (25)	592	16.6 (13)	692	2.7 (6)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ozaukee	142	43.0 (24)	127	22.8 (23)	48	* *	116	17.2 (14)	127	3.1 (8)
Pepin	51	* *	48	* *	13	* *	44	* *	48	* *
Pierce	163	46.6 (36)	149	23.5 (25)	54	* *	137	32.8 (54)	149	2.0 (2)
Polk	268	50.4 (51)	252	25.4 (32)	104	35.6 (7)	212	24.1 (43)	252	6.3 (35)
Portage	349	31.8 (5)	323	18.3 (8)	88	* *	297	13.8 (3)	323	2.2 (3)
Price	102	41.2 (20)	99	* *	33	* *	91	* *	99	* *
Racine Cou	1,429	32.5 (6)	1,353	15.6 (2)	341	32.0 (12)	1,211	22.0 (33)	1,353	9.5 (52)
Richland C	114	41.2 (20)	104	19.2 (11)	33	* *	95	* *	104	5.8 (30)
Rock	1,054	38.4 (15)	1,020	22.3 (20)	318	28.9 (15)	911	22.4 (36)	1,021	7.3 (40)
Rusk	106	41.5 (22)	104	30.8 (46)	42	* *	102	24.5 (45)	104	10.6 (56)
Sauk	412	43.7 (26)	380	29.7 (42)	134	24.6 (23)	330	22.1 (35)	380	8.2 (44)
Sawyer	152	50.0 (49)	140	32.9 (53)	61	* *	136	19.1 (24)	140	3.6 (11)
Shawano	215	49.3 (48)	204	27.5 (37)	85	* *	168	24.4 (44)	204	8.8 (47)
Sheboygan	601	33.8 (8)	576	18.2 (7)	152	36.2 (6)	483	14.9 (7)	575	4.5 (23)
St. Croix	314	45.9 (34)	280	25.4 (32)	100	38.0 (4)	248	17.3 (15)	281	3.9 (20)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	120	39.2 (17)	114	20.2 (15)	36	* *	103	20.4 (27)	114	6.1 (34)
Trempealea	199	30.7 (2)	186	15.6 (2)	41	* *	169	14.2 (4)	186	1.6 (1)
Vernon	171	42.1 (23)	165	16.4 (5)	63	* *	151	15.2 (8)	165	3.6 (11)
Vilas	142	57.7 (59)	138	29.7 (42)	71	* *	123	33.3 (55)	139	5.8 (30)
Walworth	530	30.9 (3)	501	18.0 (6)	113	34.5 (8)	411	12.2 (1)	501	8.4 (45)
Washburn	112	47.3 (38)	102	40.2 (59)	45	* *	102	14.7 (6)	102	3.9 (20)
Washington	424	48.1 (43)	406	22.4 (21)	155	51.0 (1)	325	17.8 (17)	406	3.7 (15)
Waukesha	694	43.7 (26)	651	20.9 (16)	215	32.6 (10)	512	18.8 (23)	651	6.0 (32)
Waupaca	264	46.6 (36)	248	31.5 (49)	84	* *	187	36.9 (57)	248	14.1 (58)
Waushara C	133	37.6 (11)	124	22.6 (22)	32	* *	107	27.1 (48)	124	8.9 (48)
Winnebago	818	48.8 (45)	787	24.7 (29)	301	33.2 (9)	660	18.6 (22)	791	3.8 (16)
Wood	425	37.6 (11)	394	23.6 (26)	108	24.1 (24)	343	17.8 (17)	395	7.8 (42)
Wisconsin	34,096	36.1	32,593	19.7	9,317	31.5	28,409	20.4	32,610	6.9
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9	578,852	19.9

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)											
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)		Number	%	Number	%	Number	%	
	Number	% (Rank)	% (Rank)	(Rank)							(Rank)
Adams Cty	128	8.6 (45)	13.3 (54)		132	5.3 (4)		122	5.7 (53)	128	65.6 (45)
Ashland	133	6.0 (19)	3.0 (1)		136	8.8 (27)		121	1.7 (7)	135	63.0 (48)
Barron	256	5.5 (12)	10.2 (40)		265	8.7 (24)		235	2.1 (19)	261	70.5 (32)
Bayfield C	80	*	*	*	84	*	*	75	*	82	*
Brown	1,623	6.3 (22)	8.9 (28)		1,686	9.0 (29)		1,480	3.0 (38)	1,655	70.5 (32)
Buffalo	66	*	*	*	66	*	*	58	*	66	*
Burnett	126	4.0 (4)	7.1 (12)		125	11.2 (54)		109	2.8 (34)	126	70.6 (31)
Calumet	155	5.8 (15)	5.8 (4)		158	8.9 (28)		141	3.5 (47)	156	66.7 (44)
Chippewa	307	6.8 (27)	8.8 (26)		318	7.2 (12)		284	2.5 (29)	313	73.8 (19)
Clark	186	7.0 (32)	7.0 (11)		194	7.7 (16)		172	2.9 (35)	189	78.3 (10)
Columbia	200	6.0 (19)	9.5 (33)		207	8.7 (24)		183	2.2 (22)	201	77.6 (11)
Crawford C	80	*	*	*	84	*	*	76	*	80	*
Dane	1,858	6.4 (23)	8.9 (28)		1,931	8.2 (19)		1,703	2.2 (22)	1,891	80.9 (5)
Dodge	338	4.1 (6)	10.4 (42)		344	9.0 (29)		307	2.3 (25)	339	83.2 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)						
Door	135	7.4 (37)	7.4 (16)	137	5.1 (3)	128	2.3 (25)	136	72.1 (25)
Douglas Ct	277	6.9 (30)	10.1 (38)	285	7.7 (16)	256	2.3 (25)	281	74.4 (16)
Dunn	190	6.8 (27)	9.5 (33)	197	9.1 (32)	174	3.4 (45)	196	64.3 (47)
Eau Claire	560	5.4 (11)	10.4 (42)	571	7.2 (12)	517	1.9 (14)	569	74.9 (15)
Florence	30	*	* * *	32	* *	26	* *	31	* *
Fond du La	505	9.1 (51)	7.3 (14)	523	10.7 (46)	450	3.6 (48)	509	69.5 (35)
Forest	64	*	* * *	71	* *	51	* *	64	* *
Grant Cty	252	6.0 (19)	6.7 (8)	258	10.5 (43)	224	1.8 (11)	253	66.8 (43)
Green Cty	162	6.8 (27)	8.6 (24)	166	8.4 (21)	147	1.4 (4)	165	73.3 (21)
Green Lake	78	*	* * *	82	* *	69	* *	80	* *
Iowa Cty	88	*	* * *	91	* *	78	* *	90	* *
Iron Cty	24	*	* * *	25	* *	21	* *	24	* *
Jackson	145	2.8 (1)	11.0 (48)	156	9.0 (29)	134	0.7 (1)	149	81.2 (4)
Jefferson	411	4.9 (9)	10.5 (46)	434	7.4 (14)	382	1.8 (11)	423	72.8 (23)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾		High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Juneau	179	8.9 (49)	10.1 (38)		181	11.0 (51)	159	1.9 (14)	179	73.2 (22)
Kenosha	1,149	7.7 (40)	7.7 (19)		1,190	8.5 (22)	1,050	3.1 (42)	1,164	60.2 (54)
Kewaunee	83	*	*	*	88	*	73	*	85	*
La Crosse	567	6.5 (24)	10.4 (42)		594	8.2 (19)	522	2.7 (31)	578	73.4 (20)
Lafayette	41	*	*	*	42	*	39	*	42	*
Langlade C	94	*	*	*	101	5.0 (2)	89	*	95	*
Lincoln Ct	126	5.6 (13)	11.1 (49)		130	6.9 (9)	117	1.7 (7)	128	68.0 (40)
Manitowoc	377	7.4 (37)	8.5 (23)		384	9.9 (37)	340	2.1 (19)	378	67.7 (41)
Marathon C	719	5.3 (10)	5.7 (3)		739	6.8 (8)	671	2.4 (28)	727	60.4 (53)
Marinette	222	9.9 (54)	7.7 (19)		224	10.7 (46)	197	3.0 (38)	223	57.0 (56)
Marquette	61	*	*	*	63	*	54	*	62	*
Menominee	103	2.9 (2)	18.4 (56)		107	7.5 (15)	95	*	104	78.8 (8)
Milwaukee	9,401	9.7 (53)	6.2 (7)		9,742	11.1 (52)	8,348	4.1 (51)	9,545	65.1 (46)
Monroe	272	4.0 (4)	8.8 (26)		283	9.2 (33)	247	1.6 (5)	278	72.3 (24)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	Number						
Oconto	174	4.6 (8)	9.2 (32)		181	6.6 (7)	163	2.5 (29)	176	62.5 (49)
Oneida	152	5.9 (18)	9.9 (36)		156	6.4 (6)	142	2.1 (19)	154	72.1 (25)
Outagamie	661	7.3 (35)	5.9 (6)		698	9.2 (33)	597	2.2 (22)	674	70.3 (34)
Ozaukee	125	3.2 (3)	10.4 (42)		128	9.4 (35)	113	1.8 (11)	126	78.6 (9)
Pepin	47	* * *	* *		48	* *	41	* *	47	* *
Pierce	145	7.6 (39)	9.7 (35)		150	10.0 (39)	130	3.1 (42)	148	68.2 (39)
Polk	245	7.3 (35)	10.2 (40)		251	10.8 (48)	221	0.9 (3)	250	79.6 (6)
Portage	313	7.0 (32)	8.9 (28)		324	9.6 (36)	285	3.9 (50)	321	81.6 (3)
Price	95	* * *	* *		97	* *	85	* *	96	* *
Racine Cou	1,307	9.1 (51)	5.8 (4)		1,355	10.0 (39)	1,175	3.1 (42)	1,318	60.6 (52)
Richland C	99	* * *	* *		104	7.7 (16)	91	* *	99	* *
Rock	1,011	8.4 (44)	6.7 (8)		1,040	10.0 (39)	908	3.6 (48)	1,025	61.8 (50)
Rusk	101	6.9 (30)	13.9 (55)		103	8.7 (24)	92	* *	101	74.3 (17)
Sauk	368	4.1 (6)	11.1 (49)		381	5.8 (5)	347	1.7 (7)	377	82.8 (2)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)		Number	%	Number	%	Number	%
	Number	% (Rank)	% (Rank)	(Rank)						
Sawyer	138	5.8 (15)	10.9 (47)		138	10.9 (50)	123	0.8 (2)	138	71.7 (27)
Shawano	197	8.1 (42)	5.6 (2)		203	10.8 (48)	175	2.9 (35)	199	61.3 (51)
Sheboygan	562	7.1 (34)	8.2 (22)		574	8.5 (22)	515	2.7 (31)	567	69.0 (36)
St. Croix	275	5.8 (15)	7.6 (17)		283	9.9 (37)	247	1.6 (5)	280	77.5 (12)
Taylor	108	11.1 (56)	12.0 (53)		112	7.1 (11)	100	7.0 (54)	111	73.9 (18)
Trempealea	181	7.7 (40)	7.2 (13)		182	13.2 (59)	154	1.9 (14)	181	70.7 (30)
Vernon	162	8.6 (45)	8.0 (21)		162	11.1 (52)	143	4.2 (52)	163	75.5 (14)
Vilas	134	6.7 (25)	11.2 (51)		138	11.6 (55)	118	1.7 (7)	132	68.9 (37)
Walworth	488	5.7 (14)	7.6 (17)		497	7.0 (10)	450	2.7 (31)	494	78.9 (7)
Washburn	94	*	*	*	101	4.0 (1)	92	*	95	*
Washington	390	6.7 (25)	6.7 (8)		410	10.5 (43)	351	2.0 (18)	399	76.2 (13)
Waukesha	624	9.0 (50)	8.7 (25)		656	10.2 (42)	559	2.9 (35)	637	71.3 (29)
Waupaca	230	8.3 (43)	10.0 (37)		251	12.4 (58)	203	3.0 (38)	234	68.8 (38)
Waushara C	121	10.7 (55)	11.6 (52)		132	12.1 (57)	106	1.9 (14)	130	66.9 (42)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Winnebago	769	8.7 (48)	7.3 (14)	788	11.7 (56)	679	3.4 (45)	778	71.5 (28)
Wood	372	8.6 (45)	9.1 (31)	391	10.5 (43)	334	3.0 (38)	379	59.6 (55)
Wisconsin	31,539	7.8	7.8	32,660	9.7	28,463	3.1	32,009	69.0
Nation (Prior Year)	1,085,134	8.2	6.6	1,025,278	11.5	890,981	3.6	952,980	65.4

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Adams Cty	424	89.6	0.0	4.7	0.7	0.5	4.5	0.0
Ashland	466	73.0	0.0	1.5	21.2	0.2	4.1	0.0
Barron	925	86.2	5.4	6.8	0.4	0.8	0.4	0.0
Bayfield C	274	65.3	0.0	5.1	26.6	0.7	2.2	0.0
Brown	5,384	51.6	5.3	25.1	9.4	6.6	2.0	0.0
Buffalo	217	93.1	0.0	5.1	0.0	0.0	1.8	0.0
Burnett	374	74.3	0.3	1.9	19.8	0.3	3.5	0.0
Calumet	494	71.9	2.2	17.2	0.0	7.9	0.8	0.0
Chippewa	1,079	95.6	0.2	1.4	0.2	1.4	1.2	0.0
Clark	590	76.6	0.2	20.7	1.0	0.0	1.5	0.0
Columbia	736	84.8	1.4	11.5	0.8	0.3	1.2	0.0
Crawford C	271	96.3	0.4	1.8	0.4	0.4	0.7	0.0
Dane	6,465	41.7	21.2	26.2	0.5	7.9	2.4	0.0
Dodge	1,104	79.1	0.5	18.0	0.3	0.8	1.3	0.0
Door	379	85.2	0.5	11.3	0.8	1.1	1.1	0.0
Douglas Ct	998	88.7	1.1	3.0	1.5	1.0	4.7	0.0
Dunn	703	84.9	0.9	4.0	0.1	8.7	1.4	0.0
Eau Claire	1,873	80.1	1.1	4.7	1.0	10.6	2.5	0.0
Florence	93	*	*	*	*	*	*	*
Fond du La	1,658	74.8	2.7	16.7	0.2	4.4	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Forest	244	77.9	0.0	2.9	14.3	0.4	4.5	0.0
Grant Cty	882	92.2	0.7	5.9	0.0	0.5	0.8	0.0
Green Cty	515	88.0	0.2	10.9	0.0	0.2	0.8	0.0
Green Lake	274	73.4	0.7	23.4	0.7	0.4	1.5	0.0
Iowa Cty	314	93.3	0.3	4.8	0.3	0.6	0.6	0.0
Iron Cty	83	*	*	*	*	*	*	*
Jackson	513	78.4	0.2	4.5	16.0	0.2	0.8	0.0
Jefferson	1,384	63.7	0.7	33.7	0.2	0.7	1.0	0.0
Juneau	574	91.5	0.3	3.5	0.5	0.5	3.7	0.0
Kenosha	3,686	48.5	17.0	31.5	0.3	1.1	1.7	0.0
Kewaunee	259	83.0	0.4	12.7	0.4	0.8	2.7	0.0
La Crosse	2,020	75.8	3.9	3.3	0.5	14.2	2.2	0.0
Lafayette	173	80.3	0.0	19.7	0.0	0.0	0.0	0.0
Langlade C	360	92.2	0.3	6.1	0.3	0.0	1.1	0.0
Lincoln Ct	387	92.0	0.0	6.2	0.5	0.3	1.0	0.0
Manitowoc	1,323	74.5	1.7	12.2	0.8	8.8	2.0	0.0
Marathon C	2,230	66.6	1.5	6.3	0.4	24.3	0.9	0.0
Marinette	763	94.9	0.0	3.9	0.3	0.3	0.7	0.0
Marquette	207	85.5	1.4	11.1	0.0	0.0	1.9	0.0
Menominee	422	4.5	0.0	6.6	33.2	0.0	55.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
		%	%	%	%	%	%	%
Milwaukee	31,638	18.1	49.8	25.9	0.6	4.2	1.5	0.0
Monroe	923	83.5	1.0	13.1	0.5	1.3	0.5	0.0
Oconto	503	96.2	0.2	2.2	1.0	0.2	0.2	0.0
Oneida	546	94.0	0.0	3.8	0.2	0.5	1.5	0.0
Outagamie	2,241	68.7	2.8	15.0	0.9	11.4	1.2	0.0
Ozaukee	424	84.2	3.1	8.0	0.2	2.4	2.1	0.0
Pepin	119	89.1	0.0	8.4	0.0	0.0	2.5	0.0
Pierce	541	90.8	0.7	4.6	1.3	0.7	1.8	0.0
Polk	818	94.5	0.0	3.5	0.5	0.5	1.0	0.0
Portage	1,008	76.9	0.7	9.8	0.3	11.7	0.6	0.0
Price	310	93.9	0.3	2.6	0.6	2.6	0.0	0.0
Racine Cou	4,550	39.9	27.9	29.3	0.3	0.6	2.0	0.0
Richland C	329	89.7	0.0	9.7	0.0	0.6	0.0	0.0
Rock	3,479	68.0	10.5	18.9	0.2	0.9	1.5	0.0
Rusk	331	95.2	0.0	1.8	0.0	0.3	2.7	0.0
Sauk	1,193	80.4	0.6	14.0	4.4	0.1	0.5	0.0
Sawyer	485	55.3	0.2	4.7	37.5	0.6	1.6	0.0
Shawano	692	75.9	0.4	5.2	13.6	0.4	4.5	0.0
Sheboygan	1,833	59.1	2.5	19.5	0.2	18.4	0.3	0.0
St. Croix	985	88.1	0.7	5.9	1.0	3.4	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by County
Combined 3 Years

Comparison of Racial and Ethnic Distribution ⁽²⁾								
County	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Taylor	382	90.8	0.3	6.5	0.5	0.5	1.3	0.0
Trempealea	644	69.7	0.3	28.1	0.2	0.3	1.4	0.0
Vernon	538	92.4	0.4	5.8	0.0	0.9	0.6	0.0
Vilas	436	56.7	0.2	2.3	37.2	0.0	3.7	0.0
Walworth	1,668	56.9	1.3	39.1	1.0	0.7	1.1	0.0
Washburn	346	93.6	0.9	3.8	0.0	0.6	1.2	0.0
Washington	1,301	86.2	1.2	9.2	0.5	1.2	1.6	0.0
Waukesha	2,115	67.8	3.8	24.9	0.3	1.6	1.5	0.0
Waupaca	755	89.7	0.1	8.9	0.0	0.4	0.9	0.0
Waushara C	417	68.8	0.0	30.2	0.0	0.0	1.0	0.0
Winnebago	2,363	76.3	3.7	11.4	0.4	6.7	1.6	0.0
Wood	1,346	83.2	0.6	8.9	1.7	4.3	1.3	0.0
Wisconsin	106,379	53.6	19.2	19.1	1.9	4.5	1.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Adams Cty	424	0.2	9.0	17.9	59.9	12.5	0.5	0.0
Ashland	466	0.2	4.5	12.7	58.6	20.8	3.2	0.0
Barron	925	0.0	3.0	14.6	64.0	16.9	1.5	0.0
Bayfield C	274	0.0	4.7	10.2	58.4	25.2	1.5	0.0
Brown	5,384	0.4	6.4	12.9	61.2	17.6	1.3	0.1
Buffalo	217	0.0	4.1	10.6	62.7	20.7	1.8	0.0
Burnett	374	0.0	5.9	15.0	66.8	12.0	0.3	0.0
Calumet	494	0.0	3.8	13.4	65.2	16.2	1.4	0.0
Chippewa	1,079	0.0	3.6	11.9	66.5	16.5	1.5	0.0
Clark	590	0.2	4.2	14.4	62.5	17.8	0.8	0.0
Columbia	736	0.0	3.9	13.7	63.7	16.6	2.0	0.0
Crawford C	271	0.0	3.3	15.5	64.9	14.4	1.8	0.0
Dane	6,465	0.4	5.1	11.4	61.0	20.8	1.3	0.0
Dodge	1,104	0.2	3.6	12.7	64.8	17.0	1.6	0.1
Door	379	0.0	4.2	9.5	62.3	22.4	1.6	0.0
Douglas Ct	998	0.1	5.0	12.6	67.3	13.8	1.1	0.0
Dunn	703	0.0	6.0	13.9	62.4	16.5	1.1	0.0
Eau Claire	1,873	0.1	4.0	13.3	66.5	15.3	0.9	0.0
Florence	93	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Fond du La	1,658	0.2	5.0	15.3	61.8	16.6	1.1	0.0
Forest	244	0.0	5.7	14.8	63.9	15.2	0.4	0.0
Grant Cty	882	0.0	3.6	13.2	64.7	17.5	1.0	0.0
Green Cty	515	0.6	4.9	15.1	64.7	14.2	0.6	0.0
Green Lake	274	0.4	4.4	16.4	59.9	17.9	1.1	0.0
Iowa Cty	314	0.0	3.5	10.5	65.6	20.1	0.3	0.0
Iron Cty	83	*	*	*	*	*	*	*
Jackson	513	0.2	5.1	14.8	63.5	15.8	0.4	0.2
Jefferson	1,384	0.2	3.5	12.4	62.4	19.7	1.7	0.0
Juneau	574	0.0	6.4	12.0	63.2	16.7	1.6	0.0
Kenosha	3,686	0.4	7.0	15.2	59.8	16.4	1.1	0.0
Kewaunee	259	0.4	3.9	11.2	62.5	20.1	1.5	0.4
La Crosse	2,020	0.1	4.3	13.0	64.4	16.9	1.3	0.0
Lafayette	173	0.0	3.5	10.4	70.5	13.9	1.7	0.0
Langlade C	360	0.0	6.1	12.2	65.0	16.4	0.3	0.0
Lincoln Ct	387	0.5	4.4	16.5	63.0	14.5	1.0	0.0
Manitowoc	1,323	0.2	5.7	13.4	62.9	16.7	1.2	0.0
Marathon C	2,230	0.2	5.1	13.4	63.4	16.8	1.1	0.0
Marinette	763	0.3	4.6	11.9	67.1	15.2	0.9	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Marquette	207	1.0	3.4	14.5	61.8	18.4	1.0	0.0
Menominee	422	0.7	12.6	15.6	53.6	16.8	0.7	0.0
Milwaukee	31,638	0.5	8.2	13.8	58.8	17.4	1.3	0.0
Monroe	923	0.2	5.3	14.8	64.9	14.0	0.8	0.0
Oconto	503	0.2	4.4	17.5	61.2	16.3	0.4	0.0
Oneida	546	0.0	3.5	13.6	63.4	18.1	1.5	0.0
Outagamie	2,241	0.1	5.3	12.5	64.7	16.2	1.2	0.0
Ozaukee	424	0.0	1.7	9.7	66.7	20.3	1.7	0.0
Pepin	119	0.0	0.8	10.1	70.6	17.6	0.8	0.0
Pierce	541	0.2	3.3	12.2	65.4	17.7	1.1	0.0
Polk	818	0.1	5.5	13.4	63.6	15.8	1.6	0.0
Portage	1,008	0.0	5.6	13.5	64.7	15.2	1.1	0.0
Price	310	0.0	2.6	9.0	66.8	21.0	0.6	0.0
Racine Cou	4,550	0.5	7.4	13.8	61.6	15.8	1.0	0.0
Richland C	329	0.0	4.3	11.9	65.3	17.0	1.5	0.0
Rock	3,479	0.4	6.8	15.8	60.5	15.8	0.7	0.0
Rusk	331	0.0	4.2	14.2	61.6	17.8	1.8	0.3
Sauk	1,193	0.0	4.5	14.4	65.0	14.8	1.2	0.1
Sawyer	485	0.4	5.4	15.1	62.5	15.5	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by County
Combined 3 Years

Comparison of Age Distribution ⁽²⁾								
County	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Shawano	692	0.0	4.3	12.4	63.4	18.6	1.0	0.1
Sheboygan	1,833	0.1	6.3	13.1	61.6	17.5	1.1	0.2
St. Croix	985	0.2	3.8	10.3	65.8	19.1	0.9	0.0
Taylor	382	0.3	4.2	10.2	64.4	20.2	0.8	0.0
Trempealea	644	0.3	3.4	12.1	65.2	18.2	0.8	0.0
Vernon	538	0.0	3.7	9.9	63.2	22.5	0.7	0.0
Vilas	436	0.5	4.1	14.7	61.5	17.7	1.6	0.0
Walworth	1,668	0.2	4.2	12.5	62.5	19.6	1.0	0.1
Washburn	346	0.0	4.0	10.1	67.1	18.2	0.6	0.0
Washington	1,301	0.0	4.5	13.8	62.6	18.3	0.7	0.1
Waukesha	2,115	0.2	3.7	11.5	63.7	19.5	1.4	0.0
Waupaca	755	0.3	4.5	15.9	62.4	15.8	1.2	0.0
Waushara C	417	0.0	4.8	15.8	63.1	15.3	0.7	0.2
Winnebago	2,363	0.0	4.7	13.8	65.7	15.0	0.8	0.0
Wood	1,346	0.1	3.7	14.2	66.0	14.7	1.2	0.0
Wisconsin	106,379	0.3	6.1	13.4	61.7	17.3	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	418	4.5 (55)	54.8 (43)	248	24.2 (65)	52.8 (37)	52	* *	338	18.0(32)	387	3.6 (22)	387	9.8(69)
Ashland	462	1.3 (1)	56.1 (53)	393	21.4 (47)	49.9 (15)	69	* *	369	19.2(40)	423	7.8 (70)	423	3.8(17)
Barron	907	3.9 (34)	59.2 (65)	785	17.6 (16)	56.6 (59)	122	38.5 (33)	740	22.0(52)	826	5.8 (63)	826	3.9(20)
Bayfield C	273	1.8 (2)	63.7 (70)	227	20.3 (37)	52.9 (38)	27	* *	186	14.5(12)	243	7.4 (69)	243	5.3(34)
Brown	5,309	5.0 (62)	50.8 (16)	3,482	22.4 (53)	50.9 (21)	833	36.1 (30)	4,617	26.8(64)	4,917	5.1 (54)	4,917	6.3(48)
Buffalo	217	1.8 (2)	61.8 (69)	198	17.2 (11)	56.1 (56)	39	* *	184	16.3(19)	200	3.5 (21)	200	6.0(41)
Burnett	373	3.5 (26)	60.9 (67)	225	16.4 (9)	62.7 (69)	48	* *	305	20.3(44)	337	3.0 (12)	337	0.9(1)
Calumet	488	6.1 (69)	49.0 (3)	434	20.5 (41)	49.8 (14)	85	* *	415	20.5(45)	467	6.6 (65)	467	5.8(38)
Chippewa	1,074	3.1 (18)	54.6 (40)	986	20.2 (36)	52.5 (35)	169	24.3 (12)	943	21.7(49)	997	3.8 (25)	997	5.9(39)
Clark	589	4.2 (42)	55.3 (47)	541	22.7 (54)	48.4 (8)	119	33.6 (27)	515	39.0(70)	554	2.5 (8)	554	4.7(29)
Columbia	734	4.0 (36)	52.3 (27)	556	16.2 (5)	57.7 (61)	94	* *	602	15.1(15)	659	4.7 (49)	659	8.0(63)
Crawford C	269	3.0 (15)	55.8 (50)	185	15.7 (4)	58.9 (64)	51	* *	205	21.0(46)	244	4.9 (53)	244	9.8(69)
Dane	6,376	3.7 (30)	50.7 (13)	4,000	22.1 (51)	49.3 (10)	1,239	27.3 (17)	4,608	19.9(42)	5,753	5.2 (55)	5,753	5.3(34)
Dodge	1,089	5.1 (65)	49.7 (7)	931	19.0 (28)	49.5 (12)	201	29.9 (20)	956	23.6(56)	1,011	4.7 (49)	1,011	4.5(26)
Door	379	4.7 (58)	50.1 (11)	348	17.5 (14)	50.3 (17)	73	* *	327	15.0(14)	350	5.4 (57)	350	8.6(66)
Douglas Ct	992	3.3 (19)	53.4 (31)	749	16.7 (10)	57.1 (60)	151	31.1 (23)	797	29.0(66)	859	6.8 (67)	859	6.9(58)
Dunn	692	4.0 (36)	55.3 (47)	353	19.8 (31)	53.5 (44)	101	15.8 (2)	486	15.6(18)	610	4.4 (42)	610	6.2(46)

- (1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eau Claire	1,869	4.3 (49)	49.9 (10)	125	27.2 (67)	50.4 (19)	270	31.5 (24)	1,590	23.7 (57)	1,727	4.0 (30)	1,727	3.5 (15)
Florence	92	* *	* *	72	* *	* *	14	* *	68	* *	78	* *	78	* *
Fond du La	1,625	4.2 (42)	51.9 (22)	1,447	22.7 (54)	48.1 (7)	262	36.6 (31)	1,364	23.5 (55)	1,487	4.0 (30)	1,487	6.0 (41)
Forest	242	5.0 (62)	58.3 (63)	206	21.4 (47)	58.3 (62)	30	* *	162	19.1 (39)	220	4.5 (46)	220	5.9 (39)
Grant Cty	871	3.8 (31)	56.0 (51)	616	17.5 (14)	55.0 (52)	158	22.2 (8)	732	17.2 (27)	823	4.7 (49)	823	8.1 (65)
Green Cty	512	4.1 (39)	52.9 (28)	453	17.2 (11)	54.5 (48)	90	* *	438	30.4 (68)	452	2.2 (5)	452	2.7 (8)
Green Lake	261	4.2 (42)	54.8 (43)	232	21.1 (43)	58.6 (63)	52	* *	201	14.4 (11)	239	4.2 (37)	239	3.8 (17)
Iowa Cty	312	1.9 (5)	54.2 (35)	250	20.4 (40)	52.4 (33)	55	* *	247	20.2 (43)	290	3.4 (19)	290	7.2 (60)
Iron Cty	79	* *	* *	63	* *	* *	5	* *	53	* *	74	* *	74	* *
Jackson	511	4.1 (39)	53.4 (31)	453	23.8 (63)	46.1 (4)	71	* *	414	24.2 (59)	461	3.0 (12)	461	3.3 (13)
Jefferson	1,363	3.8 (31)	53.0 (29)	856	20.1 (34)	51.6 (26)	256	19.9 (6)	1,134	12.9 (4)	1,250	4.1 (35)	1,250	6.6 (53)
Juneau	571	4.4 (50)	57.1 (59)	326	23.3 (61)	54.6 (49)	73	* *	470	13.8 (9)	524	3.6 (22)	524	8.0 (63)
Kenosha	3,602	3.5 (26)	54.9 (45)	3,329	19.4 (29)	51.3 (25)	668	33.2 (26)	3,079	29.1 (67)	3,278	3.9 (29)	3,278	6.5 (50)
Kewaunee	259	5.0 (62)	57.1 (59)	228	16.2 (5)	62.3 (68)	36	* *	216	22.7 (54)	237	3.0 (12)	237	3.8 (17)
La Crosse	2,015	4.2 (42)	49.7 (7)	753	22.7 (54)	50.3 (17)	291	23.7 (10)	1,600	23.7 (57)	1,811	3.6 (22)	1,811	1.8 (2)
Lafayette	170	2.9 (14)	61.2 (68)	108	22.2 (52)	61.1 (67)	32	* *	123	27.6 (65)	148	1.4 (1)	148	6.8 (55)
Langlade C	358	3.9 (34)	56.1 (53)	318	21.1 (43)	49.4 (11)	53	* *	312	24.7 (61)	329	3.0 (12)	329	4.9 (31)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum						
						Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number
Lincoln Ct	384	1.8 (2)	56.5 (56)	315	18.4 (23)	55.2 (53)	55	* *	337	12.8 (3)	373	5.6 (61)	373	4.6(27)
Manitowoc	1,312	4.4 (50)	49.2 (4)	443	25.5 (66)	52.6 (36)	162	26.5 (16)	1,093	18.9(38)	1,194	4.1 (35)	1,194	6.2(46)
Marathon C	2,211	4.2 (42)	50.7 (13)	1,826	23.3 (61)	46.3 (5)	366	20.2 (7)	1,932	14.0(10)	2,055	3.3 (17)	2,055	2.2(6)
Marinette	747	7.0 (70)	43.5 (1)	560	18.2 (21)	56.4 (58)	108	16.7 (3)	469	18.6(37)	673	6.1 (64)	673	5.1(32)
Marquette	201	3.0 (15)	53.2 (30)	175	24.0 (64)	52.0 (29)	33	* *	150	16.7(23)	183	3.8 (25)	183	6.0(41)
Menominee	421	2.6 (11)	59.9 (66)	337	21.1 (43)	53.4 (43)	42	* *	370	17.0(25)	385	5.5 (60)	385	6.8(55)
Milwaukee	31,313	3.3 (19)	57.0 (58)	21,859	22.7 (54)	49.7 (13)	5,244	43.6 (34)	26,631	34.5(69)	28,663	4.2 (37)	28,663	6.8(55)
Monroe	903	4.4 (50)	51.4 (20)	7	* *	* *	163	37.4 (32)	788	22.3(53)	847	4.0 (30)	847	1.8(2)
Oconto	499	5.6 (68)	51.7 (21)	470	15.5 (2)	60.0 (66)	86	* *	256	16.4(21)	472	4.0 (30)	472	6.4(49)
Oneida	545	4.6 (56)	50.8 (16)	224	20.1 (34)	54.9 (51)	78	* *	448	12.1 (2)	493	3.2 (16)	493	3.9(20)
Outagamie	2,201	4.2 (42)	49.8 (9)	1,801	19.5 (30)	51.9 (28)	360	28.6 (18)	1,896	26.2(62)	2,008	5.2 (55)	2,008	6.6(53)
Ozaukee	416	5.5 (67)	49.5 (5)	365	15.6 (3)	51.0 (23)	84	* *	345	13.6 (8)	366	6.6 (65)	366	7.7(62)
Pepin	118	2.5 (8)	58.5 (64)	109	22.0 (50)	56.0 (55)	13	* *	102	16.7(23)	108	5.6 (61)	108	6.5(50)
Pierce	539	2.4 (7)	50.6 (12)	391	17.9 (17)	55.8 (54)	53	* *	433	17.3(28)	478	4.4 (42)	478	7.3(61)
Polk	811	4.4 (50)	56.5 (56)	543	18.2 (21)	51.7 (27)	104	28.8 (19)	679	19.7(41)	739	5.4 (57)	739	8.9(68)
Portage	998	4.8 (59)	51.2 (19)	518	18.0 (19)	50.6 (20)	157	26.1 (15)	830	13.5 (6)	908	2.3 (6)	908	4.6(27)
Price	308	1.9 (5)	57.5 (61)	281	14.9 (1)	54.8 (50)	39	* *	271	12.9 (4)	283	2.8 (11)	283	5.7(36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI ⁽⁴⁾			Weight Gain ⁽⁵⁾			Anemia (Low Hb/Hct) ⁽⁶⁾				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Racine Cou	4,503	3.4 (23)	58.0 (62)	3,987	21.6 (49)	52.3 (31)	756	34.1 (28)	3,804	24.6(60)	4,037	4.4 (42)	4,037	6.5(50)
Richland C	327	2.8 (12)	56.3 (55)	252	17.9 (17)	59.1 (65)	42	* *	249	21.7(49)	297	3.4 (19)	297	6.1(44)
Rock	3,389	4.2 (42)	55.3 (47)	1,394	18.7 (24)	47.0 (6)	625	34.6 (29)	2,888	26.3(63)	3,103	1.9 (3)	3,103	2.8(9)
Rusk	331	4.8 (59)	52.0 (23)	276	22.8 (58)	48.6 (9)	38	* *	267	13.5 (6)	310	2.6 (9)	310	2.3(7)
Sauk	1,161	4.1 (39)	55.0 (46)	1,020	18.7 (24)	52.4 (33)	197	18.8 (5)	997	15.1(15)	1,068	3.8 (25)	1,068	4.2(24)
Sawyer	485	3.3 (19)	52.2 (26)	374	19.8 (31)	53.2 (42)	69	* *	355	21.4(48)	423	1.7 (2)	423	1.9(4)
Shawano	686	3.8 (31)	54.4 (37)	541	20.3 (37)	53.0 (39)	94	* *	342	18.1(33)	638	4.7 (49)	638	5.2(33)
Sheboygan	1,829	3.4 (23)	51.1 (18)	1,533	23.2 (60)	44.0 (2)	354	23.4 (9)	1,616	17.1(26)	1,693	4.5 (46)	1,693	3.1(12)
St. Croix	956	2.8 (12)	52.0 (23)	850	18.0 (19)	53.5 (44)	134	14.2 (1)	658	10.9 (1)	828	5.4 (57)	828	8.6(66)
Taylor	381	5.2 (66)	54.6 (40)	350	20.0 (33)	51.1 (24)	54	* *	336	18.2(34)	356	4.2 (37)	356	3.7(16)
Trempealea	638	3.3 (19)	54.2 (35)	245	31.0 (68)	44.5 (3)	107	30.8 (22)	526	21.3(47)	579	4.0 (30)	579	4.0(22)
Vernon	536	3.5 (26)	54.7 (42)	352	16.2 (5)	56.3 (57)	73	* *	431	14.8(13)	499	3.8 (25)	499	4.2(24)
Vilas	436	2.5 (8)	56.0 (51)	260	33.1 (69)	42.7 (1)	69	* *	327	18.3(35)	390	2.6 (9)	390	3.3(13)
Walworth	1,646	3.4 (23)	53.5 (33)	1,278	21.0 (42)	52.3 (31)	259	25.1 (14)	1,414	16.3(19)	1,499	2.1 (4)	1,499	2.0(5)
Washburn	345	3.5 (26)	54.5 (38)	282	16.3 (8)	53.9 (46)	43	* *	290	15.2(17)	303	3.3 (17)	303	3.0(11)
Washington	1,288	3.0 (15)	49.5 (5)	1,134	18.8 (27)	53.0 (39)	197	17.8 (4)	1,150	16.6(22)	1,190	4.2 (37)	1,190	7.1(59)
Waukesha	2,096	4.0 (36)	48.5 (2)	1,867	18.7 (24)	53.0 (39)	371	24.5 (13)	1,584	17.6(30)	1,927	4.6 (48)	1,927	5.7(36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by County
Combined 3 Years

Comparison of Maternal Health Indicators ^(2, 3)														
County	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Waupaca	746	4.6 (56)	50.7 (13)	669	23.0 (59)	50.2 (16)	103	30.1 (21)	558	18.5(36)	677	4.3 (41)	677	6.1(44)
Waushara C	393	2.5 (8)	54.5 (38)	349	21.2 (46)	52.1 (30)	67	* *	313	17.3(28)	351	7.1 (68)	351	4.0(22)
Winnebago	2,337	4.4 (50)	52.1 (25)	1,875	17.4 (13)	54.0 (47)	485	24.1 (11)	1,998	17.9(31)	2,168	4.4 (42)	2,168	4.7(29)
Wood	1,334	4.8 (59)	53.7 (34)	1,170	20.3 (37)	50.9 (21)	317	33.1 (25)	1,035	21.7(49)	1,224	2.3 (6)	1,224	2.9(10)
Wisconsin	105,123	3.8	54.0	74,758	21.1	51.0	17,490	32.5	86,964	25.2	96,053	4.2	96,053	5.7

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	394	89.6 (27)	3.0 (57)	410	43.7 (11)	397	23.2 (27)	342	87.4 (35)
Ashland	428	86.4 (54)	1.6 (32)	458	42.1 (14)	440	26.8 (12)	401	89.0 (27)
Barron	855	92.5 (5)	0.9 (13)	915	46.1 (9)	857	23.9 (22)	736	86.4 (38)
Bayfield C	248	94.8 (3)	1.6 (32)	267	48.7 (4)	254	24.4 (19)	225	94.2 (1)
Brown	4,869	80.1 (67)	3.2 (60)	5,226	36.4 (31)	4,947	16.8 (58)	4,320	78.8 (64)
Buffalo	205	88.3 (39)	1.0 (16)	213	32.4 (43)	207	30.4 (3)	157	84.7 (50)
Burnett	335	89.0 (32)	1.5 (28)	360	45.3 (10)	357	18.2 (47)	313	83.7 (56)
Calumet	460	86.3 (55)	2.2 (46)	484	28.5 (60)	478	22.8 (31)	368	86.4 (38)
Chippewa	1,009	92.5 (5)	0.5 (5)	1,071	36.3 (32)	1,014	25.9 (14)	869	84.2 (54)
Clark	559	88.6 (37)	0.9 (13)	569	24.1 (67)	564	29.3 (8)	464	90.7 (14)
Columbia	660	83.3 (62)	1.5 (28)	723	34.0 (37)	694	39.0 (1)	565	89.2 (26)
Crawford C	247	92.3 (9)	0.4 (4)	262	30.9 (54)	256	28.1 (10)	198	84.8 (49)
Dane	5,772	86.2 (56)	1.5 (28)	6,295	26.7 (63)	5,848	16.7 (60)	4,878	84.5 (51)
Dodge	1,025	90.0 (23)	1.3 (22)	1,085	31.2 (51)	1,045	20.8 (37)	815	86.4 (38)
Door	349	89.1 (31)	0.9 (13)	376	34.8 (36)	363	31.1 (2)	313	93.0 (4)
Douglas Ct	891	88.4 (38)	1.2 (18)	981	43.6 (12)	923	22.2 (33)	803	88.9 (29)
Dunn	631	89.4 (28)	0.8 (9)	694	42.2 (13)	659	23.7 (23)	567	89.6 (22)
Eau Claire	1,730	87.2 (48)	0.8 (9)	1,846	38.0 (26)	1,770	16.4 (61)	1,515	84.4 (53)
Florence	79	* *	* *	92	* *	84	* *	73	* *
Fond du La	1,514	86.5 (53)	3.4 (62)	1,633	38.5 (25)	1,550	17.4 (53)	1,305	81.6 (60)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Forest	221	87.3 (47)	2.3 (49)	236	48.3 (5)	224	24.6 (18)	194	92.8 (5)	
Grant Cty	830	91.1 (13)	1.8 (38)	864	31.4 (49)	833	23.2 (27)	667	89.7 (21)	
Green Cty	457	89.3 (30)	2.0 (43)	507	38.9 (21)	490	19.6 (42)	432	90.0 (19)	
Green Lake	249	88.0 (42)	0.8 (9)	271	26.6 (64)	248	24.2 (20)	189	93.1 (3)	
Iowa Cty	291	92.4 (7)	1.7 (34)	311	29.3 (59)	298	22.5 (32)	223	88.3 (33)	
Iron Cty	75	* *	* *	80	* *	75	* *	62	* *	
Jackson	472	90.5 (18)	1.3 (22)	496	39.3 (19)	485	17.5 (52)	404	76.2 (67)	
Jefferson	1,209	84.9 (58)	4.0 (64)	1,353	31.9 (44)	1,145	18.7 (44)	1,087	84.5 (51)	
Juneau	536	90.7 (15)	2.8 (55)	562	40.2 (17)	535	25.0 (16)	458	87.6 (34)	
Kenosha	3,403	77.8 (70)	5.4 (69)	3,618	30.0 (55)	3,388	17.9 (49)	2,640	83.6 (57)	
Kewaunee	237	85.2 (57)	1.7 (34)	254	33.9 (38)	247	30.4 (3)	180	88.9 (29)	
La Crosse	1,813	92.1 (10)	1.2 (18)	1,964	38.6 (23)	1,855	20.0 (39)	1,527	57.6 (69)	
Lafayette	144	92.4 (7)	0.0 (1)	173	22.5 (68)	152	17.8 (50)	120	92.5 (9)	
Langlade C	334	88.9 (34)	3.3 (61)	354	31.9 (44)	339	18.0 (48)	291	92.8 (5)	
Lincoln Ct	365	90.1 (21)	3.0 (57)	383	33.7 (40)	372	17.2 (54)	303	92.7 (8)	
Manitowoc	1,185	83.7 (60)	4.5 (67)	1,302	47.1 (7)	1,242	20.0 (39)	1,103	83.8 (55)	
Marathon C	2,066	83.4 (61)	5.5 (70)	2,197	31.2 (51)	2,037	15.1 (63)	1,766	85.6 (46)	
Marinette	688	89.0 (32)	3.6 (63)	741	39.4 (18)	701	25.0 (16)	591	86.6 (37)	
Marquette	186	88.2 (41)	2.7 (52)	200	31.5 (47)	191	29.8 (7)	146	91.8 (11)	
Menominee	389	78.1 (69)	1.8 (38)	409	50.9 (2)	402	8.7 (70)	361	92.8 (5)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)									
County	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Milwaukee	29,055	79.7 (68)	2.7 (52)	30,799	33.8 (39)	29,879	12.4 (68)	24,806	80.2 (63)
Monroe	856	87.6 (45)	0.8 (9)	905	27.8 (62)	843	18.3 (46)	676	78.3 (65)
Oconto	477	88.9 (34)	1.7 (34)	489	36.2 (33)	484	23.1 (29)	395	89.6 (22)
Oneida	503	89.9 (24)	2.2 (46)	530	39.2 (20)	479	26.7 (13)	413	90.3 (16)
Outagamie	1,874	82.0 (65)	1.5 (28)	2,214	31.4 (49)	2,097	20.9 (36)	1,697	81.6 (60)
Ozaukee	377	89.4 (28)	0.5 (5)	417	28.1 (61)	398	23.6 (24)	302	91.1 (13)
Pepin	111	90.1 (21)	2.7 (52)	116	38.8 (22)	109	28.4 (9)	94	* *
Pierce	482	90.7 (15)	1.2 (18)	526	38.6 (23)	484	22.9 (30)	411	85.6 (46)
Polk	743	87.5 (46)	4.3 (66)	811	41.2 (15)	761	23.3 (26)	627	88.8 (31)
Portage	931	87.2 (48)	1.3 (22)	991	37.4 (27)	565	17.0 (57)	781	86.0 (43)
Price	281	91.5 (11)	2.8 (55)	306	57.8 (1)	293	20.1 (38)	257	81.7 (59)
Racine Cou	4,082	81.7 (66)	2.2 (46)	4,462	37.0 (28)	4,263	13.0 (66)	3,621	80.3 (62)
Richland C	298	93.3 (4)	0.0 (1)	324	31.2 (51)	308	23.4 (25)	249	92.4 (10)
Rock	3,189	88.7 (36)	5.1 (68)	3,427	29.6 (58)	3,280	12.6 (67)	2,749	86.9 (36)
Rusk	312	91.3 (12)	1.9 (41)	319	47.3 (6)	319	30.1 (6)	278	91.7 (12)
Sauk	1,084	88.3 (39)	1.8 (38)	1,164	33.2 (41)	1,101	15.5 (62)	820	88.4 (32)
Sawyer	430	87.2 (48)	4.2 (65)	475	46.9 (8)	440	17.7 (51)	401	86.3 (42)
Shawano	631	89.9 (24)	3.0 (57)	668	36.7 (29)	640	27.2 (11)	545	93.2 (2)
Sheboygan	1,712	82.5 (64)	2.1 (44)	1,811	20.7 (70)	1,777	17.1 (55)	1,403	85.7 (45)
St. Croix	876	90.2 (20)	1.0 (16)	969	36.1 (34)	903	25.2 (15)	697	90.2 (17)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Behavioral Indicators by County
Combined 3 Years

Comparison of Maternal Behavioral Indicators ^(2, 3)										
County	Medical Care				WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Taylor	360	91.1 (13)	0.3 (3)	375	35.7 (35)	364	30.2 (5)	308	90.6 (15)	
Trempealea	585	87.7 (44)	2.1 (44)	615	33.0 (42)	606	13.2 (65)	500	68.6 (68)	
Vernon	463	95.2 (2)	0.6 (8)	525	40.8 (16)	455	14.9 (64)	420	77.6 (66)	
Vilas	385	83.1 (63)	1.3 (22)	426	31.7 (46)	400	19.0 (43)	353	89.0 (27)	
Walworth	1,530	84.1 (59)	2.5 (51)	1,633	29.8 (57)	1,545	22.2 (33)	1,223	85.4 (48)	
Washburn	310	95.5 (1)	1.3 (22)	338	50.9 (2)	328	24.1 (21)	288	89.9 (20)	
Washington	1,207	90.7 (15)	1.7 (34)	1,274	36.7 (29)	1,230	19.9 (41)	974	89.4 (25)	
Waukesha	1,935	87.2 (48)	0.5 (5)	2,082	24.7 (65)	2,002	18.6 (45)	1,473	89.5 (24)	
Waupaca	684	90.5 (18)	1.3 (22)	736	31.5 (47)	690	10.3 (69)	537	86.4 (38)	
Waushara C	374	88.0 (42)	1.9 (41)	411	29.9 (56)	376	21.3 (35)	294	90.1 (18)	
Winnebago	2,174	86.8 (52)	2.3 (49)	2,323	24.4 (66)	2,265	16.8 (58)	1,831	83.3 (58)	
Wood	1,211	89.9 (24)	1.2 (18)	1,290	21.0 (69)	1,302	17.1 (55)	1,040	86.0 (43)	
Wisconsin	96,932	84.3	2.5	103,986	33.6	98,952	17.0	82,434	83.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Adams Cty	409	51.1 (60)	389	33.7 (63)	171	22.2 (49)	367	37.6 (69)	387	18.9 (70)
Ashland	452	52.9 (64)	423	33.8 (64)	202	30.2 (23)	412	25.2 (47)	424	5.4 (19)
Barron	887	46.4 (42)	837	28.4 (40)	342	19.6 (55)	773	21.3 (30)	837	4.5 (14)
Bayfield C	261	52.5 (63)	243	30.0 (48)	112	28.6 (26)	230	23.0 (38)	243	4.1 (9)
Brown	5,129	38.5 (18)	4,904	20.8 (12)	1,550	32.3 (13)	4,409	20.0 (23)	4,902	8.8 (57)
Buffalo	203	43.8 (30)	200	32.0 (55)	71	* *	162	17.9 (16)	202	5.4 (19)
Burnett	360	54.7 (67)	337	39.2 (70)	158	12.7 (61)	319	31.7 (64)	336	7.7 (48)
Calumet	478	38.5 (18)	466	23.8 (23)	137	30.7 (19)	377	18.0 (18)	467	4.3 (13)
Chippewa	1,056	53.7 (65)	1,003	31.1 (53)	458	33.6 (10)	889	21.1 (28)	1,002	5.0 (17)
Clark	580	38.4 (17)	554	22.2 (17)	176	30.7 (19)	479	19.6 (22)	554	6.9 (41)
Columbia	718	51.4 (62)	656	28.7 (41)	258	21.7 (50)	587	23.5 (40)	654	7.2 (44)
Crawford C	255	42.4 (22)	243	25.1 (30)	76	* *	205	16.1 (9)	244	4.1 (9)
Dane	6,183	29.7 (2)	5,755	16.3 (2)	1,236	35.4 (7)	5,056	15.2 (3)	5,748	4.6 (15)
Dodge	1,067	44.2 (33)	1,010	27.0 (37)	353	28.6 (26)	845	21.4 (32)	1,010	6.2 (34)
Door	368	44.8 (37)	349	19.8 (9)	135	43.0 (2)	307	15.3 (4)	349	3.4 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Douglas Ct	938	55.4 (68)	865	33.1 (59)	419	23.6 (45)	848	27.7 (56)	866	6.5 (36)
Dunn	683	44.4 (35)	614	23.3 (22)	231	30.3 (22)	578	16.1 (9)	614	4.1 (9)
Eau Claire	1,808	44.7 (36)	1,725	26.8 (35)	645	27.9 (33)	1,554	17.5 (14)	1,726	3.1 (2)
Florence	84	* *	78	* *	27	* *	73	* *	78	* *
Fond du La	1,592	45.4 (40)	1,502	25.1 (30)	561	28.0 (31)	1,352	21.2 (29)	1,498	5.9 (27)
Forest	237	50.2 (58)	220	32.7 (56)	83	* *	201	35.3 (68)	220	8.2 (52)
Grant Cty	838	36.9 (14)	826	22.0 (16)	254	28.0 (31)	687	18.6 (20)	828	6.3 (35)
Green Cty	502	49.2 (52)	453	28.7 (41)	194	31.4 (15)	445	26.7 (52)	453	9.5 (60)
Green Lake	246	41.5 (21)	239	29.7 (47)	68	* *	193	27.5 (55)	239	9.6 (61)
Iowa Cty	297	49.2 (52)	289	33.2 (61)	110	14.5 (58)	227	22.0 (35)	289	5.9 (27)
Iron Cty	76	* *	72	* *	31	* *	62	* *	73	* *
Jackson	493	49.9 (56)	465	33.3 (62)	200	20.5 (54)	426	25.6 (49)	467	6.0 (31)
Jefferson	1,323	35.9 (11)	1,248	21.6 (14)	363	28.4 (29)	1,109	17.9 (16)	1,248	7.0 (42)
Juneau	560	46.4 (42)	528	33.9 (65)	207	15.5 (57)	476	31.3 (62)	529	15.1 (68)
Kenosha	3,290	33.6 (7)	3,321	19.8 (9)	819	33.0 (12)	2,687	24.0 (43)	3,322	7.2 (44)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Kewaunee	252	43.3 (28)	241	22.4 (18)	86	* *	194	24.7 (46)	241	5.4 (19)
La Crosse	1,950	36.5 (13)	1,797	19.4 (8)	536	25.0 (43)	1,648	15.4 (5)	1,797	2.7 (1)
Lafayette	156	44.9 (38)	147	23.1 (20)	44	* *	120	15.0 (2)	147	6.8 (40)
Langlade C	350	47.7 (49)	334	32.9 (58)	130	20.8 (53)	297	30.3 (60)	335	9.6 (61)
Lincoln Ct	381	46.5 (44)	372	29.6 (45)	136	26.5 (39)	309	32.7 (65)	373	8.3 (54)
Manitowoc	1,279	46.8 (45)	1,194	29.6 (45)	484	22.3 (48)	1,113	24.3 (44)	1,196	5.9 (27)
Marathon C	2,160	36.4 (12)	2,060	24.6 (26)	602	23.6 (45)	1,790	18.8 (21)	2,063	5.2 (18)
Marinette	714	50.0 (57)	673	32.7 (56)	289	24.6 (44)	614	27.4 (54)	673	7.4 (47)
Marquette	200	49.5 (55)	183	35.5 (69)	67	* *	151	33.1 (66)	183	10.4 (65)
Menominee	409	66.3 (70)	387	33.1 (59)	232	40.9 (3)	375	28.5 (57)	386	4.9 (16)
Milwaukee	29,779	24.3 (1)	28,734	12.8 (1)	5,396	35.9 (6)	25,551	22.4 (36)	28,721	10.7 (66)
Monroe	851	45.4 (40)	849	31.7 (54)	308	14.0 (59)	705	21.8 (33)	847	7.9 (50)
Oconto	483	48.7 (51)	475	34.1 (66)	199	19.1 (56)	399	28.6 (58)	475	9.9 (63)
Oneida	525	59.0 (69)	488	34.8 (68)	232	26.3 (40)	428	29.9 (59)	489	6.1 (33)
Outagamie	2,091	44.3 (34)	2,011	22.4 (18)	689	25.7 (41)	1,734	15.9 (8)	2,012	3.8 (8)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ozaukee	403	42.9 (25)	366	18.9 (7)	123	43.9 (1)	310	17.7 (15)	366	3.3 (4)
Pepin	117	30.8 (3)	110	21.8 (15)	31	* *	97	* *	110	7.3 (46)
Pierce	506	46.8 (45)	474	27.6 (38)	179	29.6 (24)	424	31.4 (63)	473	7.0 (42)
Polk	784	50.5 (59)	746	28.0 (39)	317	27.8 (35)	644	26.1 (50)	745	5.8 (24)
Portage	972	34.0 (8)	911	17.9 (6)	262	30.9 (17)	829	15.6 (6)	912	3.7 (6)
Price	302	42.7 (23)	283	25.1 (30)	106	22.6 (47)	276	20.3 (25)	283	6.0 (31)
Racine Cou	4,347	33.0 (5)	4,105	16.9 (3)	1,092	30.9 (17)	3,728	22.5 (37)	4,104	8.2 (52)
Richland C	320	43.8 (30)	296	25.0 (29)	103	27.2 (37)	260	21.9 (34)	296	3.7 (6)
Rock	3,286	40.7 (20)	3,121	25.1 (30)	1,046	27.9 (33)	2,806	25.5 (48)	3,118	9.1 (58)
Rusk	321	43.0 (26)	310	30.3 (50)	119	21.0 (52)	284	23.9 (41)	310	8.7 (56)
Sauk	1,154	43.2 (27)	1,076	30.8 (51)	368	21.2 (51)	940	27.3 (53)	1,077	11.6 (67)
Sawyer	470	51.1 (60)	433	29.1 (44)	196	30.6 (21)	419	23.9 (41)	431	6.7 (39)
Shawano	673	49.3 (54)	640	28.8 (43)	270	34.4 (9)	555	23.2 (39)	639	7.7 (48)
Sheboygan	1,788	34.7 (10)	1,700	19.9 (11)	452	28.8 (25)	1,446	17.2 (12)	1,698	6.5 (36)
St. Croix	925	45.1 (39)	837	24.1 (25)	292	33.2 (11)	733	17.3 (13)	838	5.8 (24)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7B
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by County
Combined 3 Years

Comparison of Smoking Indicators (2, 3)										
County	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Taylor	376	42.8 (24)	361	26.6 (34)	124	27.4 (36)	313	24.6 (45)	362	9.1 (58)
Trempealea	612	31.9 (4)	577	17.3 (4)	149	25.5 (42)	524	15.8 (7)	578	3.1 (2)
Vernon	523	38.0 (16)	498	23.1 (20)	167	31.1 (16)	447	18.1 (19)	500	4.2 (12)
Vilas	411	54.3 (66)	384	30.2 (49)	181	27.1 (38)	361	31.0 (61)	385	8.6 (55)
Walworth	1,587	33.3 (6)	1,520	17.6 (5)	375	40.3 (4)	1,244	14.6 (1)	1,519	5.9 (27)
Washburn	335	43.9 (32)	304	34.2 (67)	125	13.6 (60)	301	16.6 (11)	304	6.6 (38)
Washington	1,259	48.5 (50)	1,198	23.9 (24)	465	39.6 (5)	999	20.5 (26)	1,198	5.8 (24)
Waukesha	2,049	43.7 (29)	1,930	20.9 (13)	605	34.7 (8)	1,520	20.0 (23)	1,930	5.6 (23)
Waupaca	736	47.0 (47)	690	31.0 (52)	248	28.6 (26)	554	34.7 (67)	688	16.0 (69)
Waushara C	372	34.4 (9)	353	24.6 (26)	87	* *	301	26.6 (51)	353	10.2 (64)
Winnebago	2,275	47.2 (48)	2,151	26.9 (36)	781	31.5 (14)	1,853	20.5 (26)	2,159	5.4 (19)
Wood	1,303	37.8 (15)	1,225	24.7 (28)	352	28.4 (29)	1,053	21.3 (30)	1,225	8.0 (51)
Wisconsin	101,159	36.1	96,358	20.6	27,692	30.2	84,984	21.6	96,348	7.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one county's rate to other counties. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)										
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
Adams Cty	367	9.3 (65)	12.3 (66)	368	10.1 (46)	322	4.3 (68)	369	65.9 (46)	
Ashland	411	4.9 (2)	9.7 (51)	418	7.7 (7)	374	2.4 (21)	420	66.2 (44)	
Barron	801	6.9 (41)	9.6 (49)	839	9.7 (42)	717	2.5 (24)	826	71.3 (32)	
Bayfield C	228	5.3 (9)	14.0 (67)	234	6.4 (2)	210	2.9 (42)	234	73.9 (17)	
Brown	4,718	6.8 (34)	9.1 (43)	4,781	10.8 (58)	4,106	2.8 (39)	4,795	65.3 (50)	
Buffalo	197	9.6 (67)	7.1 (11)	199	11.6 (64)	172	1.7 (10)	199	75.4 (14)	
Burnett	326	4.9 (2)	8.3 (24)	325	9.8 (44)	283	2.5 (24)	325	71.4 (30)	
Calumet	456	7.7 (51)	7.2 (13)	455	8.4 (16)	410	2.9 (42)	461	64.6 (54)	
Chippewa	974	7.2 (47)	7.4 (15)	998	8.8 (19)	883	2.4 (21)	988	69.6 (35)	
Clark	534	7.9 (57)	8.1 (21)	528	9.5 (36)	467	3.2 (52)	538	76.8 (9)	
Columbia	625	6.6 (30)	10.7 (56)	641	7.5 (5)	571	3.9 (64)	637	72.2 (23)	
Crawford C	239	6.3 (22)	10.0 (52)	237	8.0 (10)	213	1.9 (11)	240	66.3 (43)	
Dane	5,538	6.8 (34)	9.0 (38)	5,652	8.9 (21)	4,911	2.6 (28)	5,646	79.1 (5)	
Dodge	974	5.9 (16)	8.3 (24)	997	9.6 (38)	871	2.3 (18)	990	79.7 (3)	

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm ⁽⁶⁾		Full Term LBW ⁽⁷⁾		Ever Breastfed	
	Low ⁽⁴⁾	High ⁽⁵⁾		Number	%	Number	%	Number	%
	Number	% ^(Rank)	% ^(Rank)						
Door	339	6.8 (34)	8.6 (28)	343	9.0 (23)	307	2.9 (42)	341	74.8 (16)
Douglas Ct	853	4.9 (2)	11.1 (61)	864	9.1 (26)	768	1.3 (2)	859	72.1 (25)
Dunn	601	5.7 (13)	10.3 (54)	616	8.3 (15)	544	2.6 (28)	613	69.2 (36)
Eau Claire	1,687	5.9 (16)	8.8 (34)	1,696	7.8 (8)	1,531	2.6 (28)	1,705	72.7 (20)
Florence	75	* * *	* *	77	* *	66	* *	76	* *
Fond du La	1,438	7.8 (54)	7.5 (18)	1,490	10.5 (53)	1,275	3.1 (51)	1,460	68.6 (39)
Forest	202	9.9 (70)	7.4 (15)	211	14.7 (70)	166	3.6 (59)	204	53.9 (69)
Grant Cty	791	6.2 (21)	8.7 (30)	810	9.6 (38)	697	2.0 (13)	808	65.6 (48)
Green Cty	442	6.6 (30)	9.5 (46)	445	10.3 (50)	390	1.5 (4)	446	65.7 (47)
Green Lake	234	7.7 (51)	9.4 (45)	243	9.1 (26)	211	3.3 (54)	239	58.6 (66)
Iowa Cty	281	7.8 (54)	6.8 (8)	286	8.7 (18)	253	3.6 (59)	284	69.0 (38)
Iron Cty	70	* * *	* *	71	* *	59	* *	72	* *
Jackson	442	5.2 (8)	12.2 (65)	448	8.9 (21)	389	1.5 (4)	452	76.3 (11)
Jefferson	1,190	5.0 (7)	9.6 (49)	1,228	9.4 (33)	1,057	1.5 (4)	1,213	72.0 (26)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Juneau	516	5.8 (14)	9.5 (46)	518	9.3 (30)	462	1.5 (4)	518	71.4 (30)
Kenosha	3,265	8.4 (62)	6.6 (5)	3,328	9.9 (45)	2,897	3.6 (59)	3,294	62.2 (59)
Kewaunee	228	7.5 (50)	8.8 (34)	234	9.0 (23)	204	2.9 (42)	232	72.4 (21)
La Crosse	1,718	6.8 (34)	10.2 (53)	1,746	9.5 (36)	1,514	2.7 (37)	1,742	72.3 (22)
Lafayette	144	6.3 (22)	6.3 (3)	149	5.4 (1)	136	5.1 (69)	145	66.2 (44)
Langlade C	312	7.1 (44)	8.0 (20)	329	7.0 (4)	284	3.9 (64)	317	61.8 (61)
Lincoln Ct	355	5.9 (16)	9.3 (44)	368	6.8 (3)	329	2.7 (37)	361	64.0 (56)
Manitowoc	1,175	7.7 (51)	7.8 (19)	1,192	10.6 (55)	1,039	3.7 (63)	1,189	62.1 (60)
Marathon C	2,002	6.5 (29)	6.0 (1)	2,046	8.2 (13)	1,814	2.5 (24)	2,030	60.0 (64)
Marinette	665	8.4 (62)	8.6 (28)	663	10.4 (52)	577	2.9 (42)	669	52.9 (70)
Marquette	177	6.8 (34)	6.8 (8)	179	10.6 (55)	155	2.6 (28)	181	63.5 (58)
Menominee	371	4.9 (2)	14.8 (69)	368	9.2 (28)	325	1.5 (4)	373	71.6 (28)
Milwaukee	27,456	9.4 (66)	6.1 (2)	28,049	11.1 (59)	23,823	3.9 (64)	27,867	64.5 (55)
Monroe	818	5.6 (11)	9.0 (38)	847	9.0 (23)	733	2.5 (24)	839	69.7 (34)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Oconto	458	6.3 (22)	8.7 (30)	461	8.0 (10)	412	2.9 (42)	462	58.2 (67)
Oneida	472	6.6 (30)	8.7 (30)	477	8.0 (10)	428	2.6 (28)	477	65.2 (52)
Outagamie	1,917	8.1 (59)	6.3 (3)	2,004	9.6 (38)	1,710	2.9 (42)	1,963	67.3 (41)
Ozaukee	364	4.9 (2)	10.7 (56)	367	11.2 (61)	317	1.3 (2)	368	77.2 (8)
Pepin	107	8.4 (62)	14.0 (67)	108	11.1 (59)	94	* *	109	79.8 (2)
Pierce	467	6.0 (19)	10.3 (54)	469	10.2 (48)	408	2.2 (16)	476	70.0 (33)
Polk	722	6.4 (25)	8.7 (30)	740	8.2 (13)	661	2.1 (14)	736	78.8 (7)
Portage	886	7.2 (47)	9.0 (38)	904	10.2 (48)	785	3.2 (52)	904	75.8 (13)
Price	270	5.6 (11)	8.5 (27)	277	10.1 (46)	240	2.1 (14)	276	75.0 (15)
Racine Cou	3,988	9.6 (67)	6.6 (5)	4,042	11.6 (64)	3,457	3.5 (58)	4,019	60.3 (63)
Richland C	283	5.3 (9)	15.2 (70)	291	9.3 (30)	253	2.4 (21)	288	80.9 (1)
Rock	3,066	7.2 (47)	8.1 (21)	3,130	9.4 (33)	2,736	3.0 (49)	3,114	54.1 (68)
Rusk	299	6.4 (25)	12.0 (64)	295	8.5 (17)	264	3.4 (56)	300	64.7 (53)
Sauk	1,045	4.4 (1)	10.8 (58)	1,057	7.9 (9)	946	1.9 (11)	1,067	79.7 (3)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sawyer	420	6.4 (25)	9.0 (38)	421	10.7 (57)	368	1.6 (9)	421	69.1 (37)
Shawano	617	6.8 (34)	7.3 (14)	620	11.9 (68)	527	2.3 (18)	625	65.6 (48)
Sheboygan	1,641	7.1 (44)	7.4 (15)	1,668	9.4 (33)	1,475	3.0 (49)	1,656	66.9 (42)
St. Croix	831	6.6 (30)	8.9 (36)	844	8.8 (19)	743	2.6 (28)	851	76.3 (11)
Taylor	342	8.2 (60)	10.8 (58)	351	9.7 (42)	304	3.6 (59)	348	71.6 (28)
Trempealea	544	6.4 (25)	9.0 (38)	550	11.8 (67)	459	2.2 (16)	555	72.8 (18)
Vernon	486	7.0 (43)	11.1 (61)	486	9.3 (30)	430	2.6 (28)	492	76.4 (10)
Vilas	368	6.0 (19)	11.4 (63)	375	11.7 (66)	318	0.9 (1)	367	61.3 (62)
Walworth	1,490	5.8 (14)	8.9 (36)	1,509	7.6 (6)	1,351	2.8 (39)	1,512	79.0 (6)
Washburn	289	6.9 (41)	6.6 (5)	294	10.5 (53)	254	3.9 (64)	292	71.9 (27)
Washington	1,135	6.8 (34)	7.0 (10)	1,177	10.3 (50)	1,008	2.6 (28)	1,167	72.8 (18)
Waukesha	1,868	7.1 (44)	8.3 (24)	1,914	9.2 (28)	1,668	2.6 (28)	1,901	72.2 (23)
Waupaca	631	9.7 (69)	9.5 (46)	679	12.7 (69)	539	3.3 (54)	649	65.3 (50)
Waushara C	344	7.8 (54)	10.8 (58)	369	11.4 (63)	301	2.3 (18)	364	64.0 (56)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one county's rate to other counties. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- * Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8B
2007 - 2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by County
Combined 3 Years

Comparison of Infant Health Indicators ^(2, 3)									
County	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Winnebago	2,101	8.3 (61)	7.1 (11)	2,140	11.2 (61)	1,840	3.4 (56)	2,134	68.5 (40)
Wood	1,141	7.9 (57)	8.1 (21)	1,166	9.6 (38)	986	2.8 (39)	1,165	58.7 (65)
Wisconsin	92,827	7.7	7.8	94,701	10.1	81,797	3.1	94,255	67.3

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one county's rate to other counties. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

