

**Table 3E**  
**2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Region**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Region</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic</b>	<b>Black, Not Hispanic</b>	<b>Hispanic</b>	<b>American Indian/Alaska Native</b>	<b>Asian/Pacific Islander</b>	<b>Multiple Races</b>	<b>All Other/Unknown</b>
		%	%	%	%	%	%	%
NERO	6,876	67.5	2.9	16.3	4.0	6.5	2.6	0.0
NRO	2,916	76.3	0.5	6.7	6.1	8.8	1.5	0.0
SERO	15,338	29.3	39.0	26.1	0.6	3.3	1.7	0.0
SRO	6,123	65.7	9.7	19.1	0.8	3.2	1.6	0.0
WRO	4,643	83.6	1.4	6.6	1.8	4.7	1.8	0.0
<b>Wisconsin</b>	<b>35,896</b>	<b>53.7</b>	<b>19.1</b>	<b>18.9</b>	<b>1.9</b>	<b>4.5</b>	<b>1.9</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,307,887</b>	<b>42.0</b>	<b>24.2</b>	<b>28.1</b>	<b>1.0</b>	<b>3.1</b>	<b>1.1</b>	<b>0.5</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4E**  
**2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Region**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Region</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
NERO	6,876	0.3	5.0	13.4	62.6	17.6	1.1	0.1
NRO	2,916	0.1	4.0	12.6	64.5	17.7	1.2	0.0
SERO	15,338	0.4	6.9	13.0	59.8	18.6	1.3	0.0
SRO	6,123	0.2	4.2	13.2	61.8	19.2	1.4	0.0
WRO	4,643	0.1	4.0	12.4	65.2	17.1	1.2	0.0
<b>Wisconsin</b>	<b>35,896</b>	<b>0.3</b>	<b>5.5</b>	<b>13.0</b>	<b>61.8</b>	<b>18.3</b>	<b>1.2</b>	<b>0.0</b>
<b>Nation (Prior Year)</b>	<b>1,307,887</b>	<b>0.3</b>	<b>6.3</b>	<b>12.6</b>	<b>59.7</b>	<b>19.4</b>	<b>1.4</b>	<b>0.3</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5E**  
**2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Region**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Region</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia (Low Hb/Hct)</b> (6)				<b>Gestational Diabetes During Pregnancy</b>		<b>Hypertension During Pregnancy</b>	
	<b>Under-weight</b>		<b>Over-weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	6,783	4.7 (5)	52.1 (1)	5,395	21.0 (2)	51.2 (4)	1,079	28.2 (4)	5,317	20.4 (2)	6,300	5.0 (5)	6,300	5.8(4)
NRO	2,894	3.7 (3)	55.5 (4)	2,167	22.1 (3)	50.9 (3)	438	26.7 (2)	2,299	17.9 (1)	2,670	3.8 (1)	2,670	3.7(1)
SERO	15,168	3.2 (1)	56.7 (5)	11,780	22.4 (4)	50.1 (1)	2,460	38.5 (5)	12,535	30.9 (5)	13,964	4.6 (3)	13,964	6.7(5)
SRO	6,023	3.6 (2)	54.4 (2)	4,577	19.4 (1)	52.7 (5)	1,076	25.4 (1)	4,720	20.6 (3)	5,525	4.4 (2)	5,525	5.6(3)
WRO	4,614	3.9 (4)	54.8 (3)	2,635	22.5 (5)	50.6 (2)	644	27.5 (3)	3,598	21.6 (4)	4,183	4.9 (4)	4,183	4.1(2)
<b>Wisconsin</b>	<b>35,482</b>	<b>3.7</b>	<b>55.1</b>	<b>26,554</b>	<b>21.6</b>	<b>50.9</b>	<b>5,697</b>	<b>31.9</b>	<b>28,469</b>	<b>25.0</b>	<b>32,642</b>	<b>4.6</b>	<b>32,642</b>	<b>5.8</b>
<b>Nation (Prior Year)</b>	<b>1,224,766</b>	<b>4.7</b>	<b>52.1</b>	<b>963,624</b>	<b>21.4</b>	<b>48.2</b>	<b>188,530</b>	<b>33.8</b>	<b>416,710</b>	<b>29.6</b>	<b>346,704</b>	<b>4.4</b>	<b>395,960</b>	<b>5.2</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6E**  
**2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Region**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>													
<b>Region</b>	Medical Care				WIC Enrollment			Multivitamin Consumption					
	1st Trimester		No Care		1st Trimester			Pregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	%	% (Rank)	Number	%	% (Rank)	Number	%	% (Rank)	%		
NERO	6,311	85.1 (4)	3.0	(4)	6,853	33.6	(3)	6,643	19.4	(2)	5,358	84.0	(3)
NRO	2,701	89.4 (3)	2.4	(3)	2,909	36.0	(2)	2,722	19.2	(3)	2,353	87.8	(1)
SERO	14,284	81.8 (5)	3.4	(5)	15,276	33.6	(3)	15,008	14.3	(5)	11,902	82.5	(4)
SRO	5,609	89.9 (2)	1.9	(2)	6,112	31.4	(5)	5,842	18.3	(4)	4,792	86.8	(2)
WRO	4,305	90.7 (1)	1.6	(1)	4,628	39.1	(1)	4,541	21.5	(1)	3,662	81.6	(5)
<b>Wisconsin</b>	<b>33,210</b>	<b>85.6</b>	<b>2.8</b>		<b>35,778</b>	<b>34.1</b>		<b>34,756</b>	<b>17.3</b>		<b>28,067</b>	<b>83.8</b>	
<b>Nation (Prior Year)</b>	<b>580,485</b>	<b>80.1</b>	<b>3.1</b>		<b>1,187,387</b>	<b>31.7</b>		<b>266,567</b>	<b>19.3</b>		<b>269,548</b>	<b>80.6</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7E**  
**2009 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Region**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Region</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	6,581	43.8 (4)	6,282	23.6 (3)	2,220	29.9 (3)	5,417	21.0 (5)	6,286	6.5 (4)
NRO	2,822	41.7 (3)	2,659	25.5 (4)	896	27.5 (4)	2,380	20.7 (3)	2,664	5.1 (2)
SERO	14,382	27.6 (1)	13,959	14.1 (1)	2,946	36.8 (1)	12,009	20.8 (4)	13,961	8.4 (5)
SRO	5,859	38.6 (2)	5,520	21.9 (2)	1,706	30.8 (2)	4,855	19.1 (1)	5,522	6.1 (3)
WRO	4,452	45.0 (5)	4,173	25.9 (5)	1,549	26.8 (5)	3,748	19.5 (2)	4,177	4.5 (1)
<b>Wisconsin</b>	<b>34,096</b>	<b>36.1</b>	<b>32,593</b>	<b>19.7</b>	<b>9,317</b>	<b>31.5</b>	<b>28,409</b>	<b>20.4</b>	<b>32,610</b>	<b>6.9</b>
<b>Nation (Prior Year)</b>	<b>999,199</b>	<b>23.1</b>	<b>856,953</b>	<b>14.2</b>	<b>146,933</b>	<b>35.5</b>	<b>655,095</b>	<b>18.9</b>	<b>578,852</b>	<b>19.9</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8E**  
**2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Region**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Region</b>	<b>Birthweight</b>				<b>Preterm</b> (6)		<b>Full Term LBW</b> (7)		<b>Ever Breastfed</b>	
	Low (4)		High (5)							
	Number	% (Rank)	% (Rank)		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
NERO	6,076	7.4 (4)	8.0 (2)		6,300	9.7 (4)	5,485	2.8 (3)	6,166	69.0 (3)
NRO	2,562	6.6 (3)	8.4 (3)		2,653	8.7 (2)	2,338	2.8 (3)	2,597	67.3 (4)
SERO	13,484	9.2 (5)	6.5 (1)		13,978	10.6 (5)	12,046	3.8 (5)	13,683	65.5 (5)
SRO	5,377	6.5 (2)	9.0 (4)		5,557	8.5 (1)	4,909	2.5 (2)	5,455	74.8 (1)
WRO	4,040	6.2 (1)	9.5 (5)		4,172	8.8 (3)	3,685	2.2 (1)	4,108	73.9 (2)
<b>Wisconsin</b>	<b>31,539</b>	<b>7.8</b>	<b>7.8</b>		<b>32,660</b>	<b>9.7</b>	<b>28,463</b>	<b>3.1</b>	<b>32,009</b>	<b>69.0</b>
<b>Nation (Prior Year)</b>	<b>1,085,134</b>	<b>8.2</b>	<b>6.6</b>		<b>1,025,278</b>	<b>11.5</b>	<b>890,981</b>	<b>3.6</b>	<b>952,980</b>	<b>65.4</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Racial and Ethnic Distribution by Region**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Region</b>	Number Included in Analysis	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
NERO	20,040	67.0	2.9	16.5	4.3	6.8	2.6	0.0
NRO	8,577	77.0	0.6	6.1	6.1	8.6	1.5	0.0
SERO	45,382	29.1	39.2	26.5	0.5	3.2	1.5	0.0
SRO	18,381	65.6	9.7	19.2	0.6	3.2	1.7	0.0
WRO	13,999	84.1	1.4	6.4	1.7	4.6	1.8	0.0
<b>Wisconsin</b>	<b>106,379</b>	<b>53.6</b>	<b>19.2</b>	<b>19.1</b>	<b>1.9</b>	<b>4.5</b>	<b>1.8</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Age Distribution by Region**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Region</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
NERO	20,040	0.2	5.5	13.5	62.6	16.9	1.1	0.1
NRO	8,577	0.2	4.6	13.3	63.8	16.9	1.2	0.0
SERO	45,382	0.5	7.5	13.7	59.7	17.4	1.2	0.0
SRO	18,381	0.2	5.1	13.0	62.3	18.1	1.2	0.0
WRO	13,999	0.1	4.3	13.0	65.1	16.4	1.1	0.0
<b>Wisconsin</b>	<b>106,379</b>	<b>0.3</b>	<b>6.1</b>	<b>13.4</b>	<b>61.7</b>	<b>17.3</b>	<b>1.2</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Health Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Region</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia (Low Hb/Hct)</b> (6)				<b>Gestational Diabetes During Pregnancy</b>		<b>Hypertension During Pregnancy</b>	
	<b>Under- weight</b>		<b>Over- weight</b>	<b>&lt; Ideal</b>		<b>&gt; Ideal</b>	<b>3rd Trimester</b>		<b>Postpartum</b>					
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NERO	19,766	4.5 (5)	51.1 (1)	14,990	20.9 (3)	51.3 (3)	3,253	28.3 (4)	16,237	21.9 (3)	18,210	4.9 (5)	18,210	5.6(4)
NRO	8,515	3.9 (3)	53.2 (3)	6,531	21.1 (4)	50.2 (1)	1,384	24.6 (1)	6,985	16.6 (1)	7,801	3.3 (1)	7,801	3.5(1)
SERO	44,864	3.4 (1)	56.1 (5)	33,819	21.8 (5)	50.5 (2)	7,579	39.3 (5)	38,007	31.0 (5)	40,960	4.2 (3)	40,960	6.6(5)
SRO	18,098	3.9 (3)	53.1 (2)	11,547	19.9 (1)	51.4 (4)	3,238	27.1 (2)	14,418	20.3 (2)	16,508	4.0 (2)	16,508	5.3(3)
WRO	13,880	3.8 (2)	53.3 (4)	7,871	20.0 (2)	52.8 (5)	2,036	28.2 (3)	11,317	22.0 (4)	12,574	4.3 (4)	12,574	4.5(2)
<b>Wisconsin</b>	<b>105,123</b>	<b>3.8</b>	<b>54.0</b>	<b>74,758</b>	<b>21.1</b>	<b>51.0</b>	<b>17,490</b>	<b>32.5</b>	<b>86,964</b>	<b>25.2</b>	<b>96,053</b>	<b>4.2</b>	<b>96,053</b>	<b>5.7</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Maternal Behavioral Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>												
<b>Region</b>	Medical Care					WIC Enrollment			Multivitamin Consumption			
	1st Trimester		No Care			1st Trimester			Pregnancy (4)		During Pregnancy (5)	
	Number	% (Rank)	%	% (Rank)	%	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
NERO	18,123	84.1 (4)	2.6	(3)	19,619	33.5 (3)	18,769	18.8 (3)	15,646	83.6 (3)		
NRO	7,846	87.4 (3)	2.8	(5)	8,389	34.9 (2)	7,597	19.5 (2)	6,800	87.7 (1)		
SERO	41,589	80.6 (5)	2.7	(4)	44,285	33.3 (4)	42,705	13.9 (5)	35,039	81.4 (4)		
SRO	16,599	88.0 (2)	2.4	(2)	17,985	30.5 (5)	16,837	18.3 (4)	14,023	86.2 (2)		
WRO	12,775	89.8 (1)	1.3	(1)	13,708	38.4 (1)	13,044	21.7 (1)	10,926	81.3 (5)		
<b>Wisconsin</b>	<b>96,932</b>	<b>84.3</b>	<b>2.5</b>		<b>103,986</b>	<b>33.6</b>	<b>98,952</b>	<b>17.0</b>	<b>82,434</b>	<b>83.1</b>		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** (1)  
**Wisconsin**  
**Comparison of Smoking Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Region</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	Number	% (Rank)	Number	% (Rank)			Number	% (Rank)	Number	% (Rank)
NERO	19,158	43.3 (4)	18,229	24.8 (3)	6,381	29.4 (2)	15,995	21.4 (3)	18,229	7.1 (3)
NRO	8,287	42.6 (3)	7,816	26.4 (4)	2,730	27.0 (4)	6,985	22.6 (5)	7,825	6.4 (2)
SERO	42,714	28.1 (1)	41,174	14.7 (1)	8,875	35.4 (1)	36,039	22.1 (4)	41,160	9.5 (5)
SRO	17,591	38.0 (2)	16,535	22.9 (2)	4,950	28.4 (3)	14,577	20.8 (2)	16,528	7.2 (4)
WRO	13,409	45.0 (5)	12,604	27.0 (5)	4,756	25.1 (5)	11,388	20.5 (1)	12,606	5.1 (1)
<b>Wisconsin</b>	<b>101,159</b>	<b>36.1</b>	<b>96,358</b>	<b>20.6</b>	<b>27,692</b>	<b>30.2</b>	<b>84,984</b>	<b>21.6</b>	<b>96,348</b>	<b>7.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one region's rate to other regions. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8E**  
**2007 - 2009 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Wisconsin**  
**Comparison of Infant Health Indicators by Region**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Region</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
NERO	17,577	7.5 (4)	8.1 (2)	17,958	10.4 (4)	15,476	3.0 (4)	17,861	65.6 (3)
NRO	7,487	6.8 (3)	8.3 (3)	7,656	9.2 (3)	6,668	2.7 (3)	7,604	64.7 (5)
SERO	39,566	9.0 (5)	6.5 (1)	40,386	10.8 (5)	34,521	3.7 (5)	40,128	65.2 (4)
SRO	15,987	6.5 (2)	9.2 (4)	16,295	9.0 (1)	14,248	2.6 (2)	16,257	71.5 (2)
WRO	12,210	6.4 (1)	9.4 (5)	12,406	9.1 (2)	10,884	2.4 (1)	12,405	72.6 (1)
<b>Wisconsin</b>	<b>92,827</b>	<b>7.7</b>	<b>7.8</b>	<b>94,701</b>	<b>10.1</b>	<b>81,797</b>	<b>3.1</b>	<b>94,255</b>	<b>67.3</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one region's rate to other regions. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

