

Table 4E
2007 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Region
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Region								
Region	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
NERO	23,533	56.5	2.8	23.6	4.6	7.0	5.4	0.0
NRO	10,041	70.1	0.9	7.6	7.1	9.2	5.0	0.0
SERO	53,160	20.9	37.6	33.8	0.3	2.8	4.6	0.0
SRO	19,057	57.0	10.2	24.3	0.4	2.9	5.1	0.0
WRO	16,866	77.8	1.4	8.5	1.6	4.5	6.2	0.0
Unknown Regions	142	6.3	0.0	6.3	56.3	0.0	30.3	0.7
Wisconsin	122,799	45.2	18.6	24.7	1.9	4.4	5.1	0.0
Nation (Prior Year)	7,599,318	31.5	18.5	41.1	1.0	2.5	1.5	3.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5E
2007 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Age Distribution by Region
Children Aged < 5 Years (2)

Comparison of Age Distribution by Region				
Region	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
NERO	23,533	34.6	21.5	43.9
NRO	10,041	36.7	20.2	43.1
SERO	53,160	36.5	21.5	42.0
SRO	19,057	37.5	20.8	41.7
WRO	16,866	36.3	20.4	43.3
Unknown Regions	142	26.1	26.8	47.2
Wisconsin	122,799	36.3	21.1	42.6
Nation (Prior Year)	7,599,318	37.6	21.8	40.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6E
2007 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Region (3,4)												
Region	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
NERO	5,856	8.1 (3)	7.9 (2)	22,979	4.9 (2)	4.0 (6)	12.1 (3)	10,167	15.8 (2)	12.9 (3)	16,430	7.3 (2)
NRO	2,513	7.1 (1)	8.4 (3)	9,840	6.1 (6)	3.0 (2)	11.9 (2)	4,234	17.8 (5)	12.6 (2)	6,865	5.5 (1)
SERO	13,794	9.9 (5)	6.3 (1)	51,980	5.6 (5)	3.9 (5)	12.3 (4)	22,036	15.9 (3)	13.5 (4)	35,956	15.8 (6)
SRO	4,928	8.2 (4)	8.7 (4)	18,622	5.4 (4)	3.2 (3)	12.8 (5)	7,888	15.3 (1)	13.5 (4)	12,917	9.1 (4)
WRO	4,045	7.2 (2)	8.9 (5)	16,513	5.3 (3)	3.3 (4)	11.6 (1)	7,249	16.9 (4)	12.0 (1)	11,724	8.6 (3)
Unknown Regions	26	* *	* *	137	2.2 (1)	2.2 (1)	25.5 (6)	62	* *	* *	114	9.6 (5)
Wisconsin	31,162	8.7	7.5	120,071	5.4	3.7	12.2	51,636	16.1	13.1	84,006	11.3
Nation (Prior Year)	1,842,970	9.2	6.6	7,192,043	6.4	4.7	13.8	2,945,212	16.4	14.8	4,521,591	14.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares this region's rate to other regions. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.
(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7E
2007 Pediatric Nutrition Surveillance (1)
Wisconsin

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region
(4,5)

Region	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
NERO	5,877	60.8 (5)	2,499	24.8 (4)	3,606	11.4 (3)	1,805	16.7 (4)	2,290	6.1 (4)	11,006	82.1 (4)	23,252	9.6 (2)
NRO	2,517	61.4 (3)	1,131	24.5 (5)	1,456	11.1 (4)	788	20.2 (1)	1,037	8.0 (2)	4,603	83.3 (3)	9,946	10.0 (3)
SERO	13,771	61.3 (4)	5,806	26.1 (2)	8,385	12.0 (2)	3,969	13.0 (5)	5,425	4.7 (5)	23,836	79.3 (5)	52,567	12.3 (5)
SRO	4,933	67.0 (2)	2,194	32.5 (1)	2,915	16.4 (1)	1,683	19.1 (3)	1,947	8.8 (1)	8,479	84.5 (2)	18,705	10.7 (4)
WRO	4,052	70.0 (1)	1,968	25.4 (3)	2,566	10.5 (5)	1,429	19.4 (2)	1,810	7.3 (3)	7,812	85.3 (1)	16,732	8.9 (1)
Unknown Regions	21	* *	18	* *	24	* *	12	* *	18	* *	73	* *	138	13.8 (6)
Wisconsin	31,171	63.2	13,616	26.6	18,952	12.3	9,686	16.3	12,527	6.2	55,809	81.8	121,340	10.9
Nation (Prior Year)	1,577,832	60.1	790,341	25.2	875,205	18.1	104,492	10.3	143,261	5.6	584,732	73.3	2,855,911	10.6

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6E
2005 - 2007 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Region (3,4)												
Region	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
NERO	16,747	8.1 (3)	8.0 (2)	66,882	5.2 (1)	4.1 (4)	12.3 (3)	30,298	16.0 (2)	13.0 (3)	47,656	8.0 (2)
NRO	7,196	7.3 (1)	8.9 (3)	28,999	6.0 (5)	3.2 (1)	12.1 (2)	12,930	16.4 (4)	12.1 (1)	20,555	5.4 (1)
SERO	40,260	9.6 (6)	6.6 (1)	152,211	5.7 (4)	4.4 (5)	12.3 (3)	66,382	16.2 (3)	13.4 (5)	105,694	15.5 (6)
SRO	14,278	8.3 (4)	8.9 (3)	54,540	5.4 (2)	3.3 (2)	12.7 (5)	23,391	15.6 (1)	13.1 (4)	37,951	8.9 (4)
WRO	11,725	7.4 (2)	9.0 (5)	47,841	5.6 (3)	3.5 (3)	11.7 (1)	21,463	16.7 (5)	12.6 (2)	33,798	8.7 (3)
Unknown Regions	446	8.7 (5)	9.6 (6)	899	6.5 (6)	6.5 (6)	16.1 (6)	249	18.9 (6)	22.5 (6)	434	12.7 (5)
Wisconsin	90,652	8.6	7.7	351,372	5.6	4.0	12.3	154,713	16.2	13.0	246,088	11.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this region's rate to other regions. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7E
2005 - 2007 Pediatric Nutrition Surveillance (1)
Wisconsin

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region
(4,5)

Region	Breastfeeding (5)				Exclusive Breastfeeding (6)				≤2 Hours TV Viewing/Day (7)		Smoking in Household			
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% (Rank)	Number	% (Rank)		
NERO	18,398	61.2 (3)	11,544	22.2 (5)	12,034	13.1 (3)	9,019	12.1 (4)	8,766	5.5 (4)	24,612	79.8 (5)	37,277	17.6 (2)
NRO	7,866	59.8 (6)	5,060	22.5 (4)	4,925	14.1 (2)	3,696	16.3 (3)	3,595	7.6 (3)	10,562	82.1 (4)	16,002	19.8 (5)
SERO	43,338	60.5 (4)	26,477	22.6 (3)	27,050	12.8 (4)	20,042	10.0 (5)	19,684	4.5 (5)	54,138	77.4 (6)	84,667	18.2 (3)
SRO	15,738	67.1 (2)	9,992	29.1 (1)	9,869	16.7 (1)	7,408	16.4 (2)	7,049	8.8 (1)	18,977	84.6 (2)	30,914	18.9 (4)
WRO	12,875	68.5 (1)	8,415	23.7 (2)	8,264	12.8 (4)	6,435	16.9 (1)	6,197	7.9 (2)	17,522	84.0 (3)	26,373	17.2 (1)
Unknown Regions	272	59.9 (5)	206	15.0 (6)	209	6.2 (6)	164	5.5 (6)	161	3.1 (6)	199	94.0 (1)	450	26.7 (6)
Wisconsin	98,487	62.7	61,694	23.7	62,351	13.5	46,764	12.8	45,452	6.1	126,010	80.3	195,683	18.2

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in

(6) the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) Children ages 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

