

Table 4E
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Racial and Ethnic Distribution by Region
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution								
Region	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other / Unknown %
NERO	26,039	57.0	3.0	23.3	3.3	6.8	6.6	0.0
NRO	11,076	68.9	0.8	8.5	7.0	9.2	5.6	0.0
SERO	57,496	20.8	37.5	33.6	0.3	3.1	4.7	0.0
SRO	23,200	57.6	9.1	25.6	0.5	2.6	4.6	0.0
WRO	18,245	76.5	1.4	9.4	1.9	4.6	6.1	0.0
Unknown Regions	1	*	*	*	*	*	*	*
Wisconsin	136,057	45.4	18.2	25.0	1.6	4.4	5.3	0.0
Nation (Prior Year)	8,164,612	31.7	18.8	41.0	1.0	2.6	2.1	2.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 5E
2009 Pediatric Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Region
Children Aged < 5 Years ⁽²⁾

Comparison of Age Distribution				
Region	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
NERO	26,039	32.9	21.6	45.6
NRO	11,076	34.5	20.1	45.4
SERO	57,496	34.5	21.0	44.5
SRO	23,200	35.3	20.0	44.7
WRO	18,245	33.6	20.5	45.8
Unknown Regions	1	*	*	*
Wisconsin	136,057	34.2	20.8	45.0
Nation (Prior Year)	8,164,612	35.8	23.0	41.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 6E
2009 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Region	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Obese (7)		≥ 2 Yrs Over-weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
NERO	6,120	8.4 (4)	7.9 (2)	25,462	5.1 (1)	3.9 (5)	12.8 (2)	11,682	16.2 (1)	13.6 (3)	18,737	6.5 (1)
NRO	2,581	7.6 (3)	8.3 (3)	10,843	6.0 (4)	2.8 (2)	13.1 (3)	4,945	16.2 (1)	13.9 (4)	8,059	7.7 (2)
SERO	13,856	9.9 (5)	6.4 (1)	56,311	6.0 (4)	3.2 (4)	13.6 (5)	25,132	16.9 (3)	13.9 (4)	41,031	15.6 (5)
SRO	5,347	6.9 (1)	8.9 (4)	22,823	5.2 (2)	2.6 (1)	13.1 (3)	10,269	16.9 (3)	13.5 (2)	16,764	8.5 (3)
WRO	4,070	7.0 (2)	9.6 (5)	17,950	5.4 (3)	3.0 (3)	12.7 (1)	8,252	17.8 (5)	13.3 (1)	13,241	9.0 (4)
Wisconsin	31,974	8.6	7.7	133,389	5.6	3.2	13.2	60,280	16.8	13.7	97,832	11.1
Nation (Prior Year)	1,909,620	9.0	6.4	7,107,000	6.0	4.5	14.1	2,943,996	16.5	14.8	4,916,523	14.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this region's rate to other regions. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



**Table 7E
2009 Pediatric Nutrition Surveillance (1)
Wisconsin**

**Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)**

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)				Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household			
Region	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)		
NERO	6,137	67.2 (3)	2,543	25.5 (5)	4,011	11.0 (5)	2,020	20.3 (4)	2,467	8.7 (4)	12,583	82.3 (4)	25,898	8.0 (3)
NRO	2,579	65.2 (4)	1,145	25.8 (4)	1,486	11.8 (4)	919	21.5 (3)	1,084	11.1 (1)	5,351	85.6 (2)	11,030	8.0 (3)
SERO	13,885	63.1 (5)	5,970	26.2 (3)	8,739	12.7 (2)	4,323	14.3 (5)	5,812	6.6 (5)	27,215	79.8 (5)	57,152	9.8 (5)
SRO	5,364	73.1 (1)	2,595	34.1 (1)	3,417	18.1 (1)	2,081	22.1 (2)	2,519	10.2 (3)	11,030	83.5 (3)	23,067	7.4 (2)
WRO	4,073	72.4 (2)	1,932	26.7 (2)	2,721	12.5 (3)	1,502	23.6 (1)	1,834	10.6 (2)	8,875	86.0 (1)	18,172	6.6 (1)
Wisconsin	32,038	66.9	14,185	27.5	20,374	13.2	10,845	18.8	13,716	8.6	65,054	82.2	135,319	8.5
Nation (Prior Year)	1,719,083	62.0	848,761	26.9	1,038,203	19.1	144,421	12.9	206,416	7.2	744,697	74.8	3,277,352	10.9

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 6E
2007 - 2009 Pediatric Nutrition Surveillance (1)
Wisconsin
Comparison of Growth and Anemia Indicators by Region
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)												
Region	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Obese (7)		≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number
NERO	17,914	8.3 (4)	8.0 (2)	72,510	5.0 (1)	4.0 (5)	12.8 (3)	32,730	15.9 (1)	13.4 (2)	52,706	7.0 (2)
NRO	7,604	7.5 (3)	8.2 (3)	31,145	6.0 (5)	2.9 (1)	12.7 (2)	13,837	16.8 (4)	13.5 (4)	22,624	6.5 (1)
SERO	41,053	9.8 (5)	6.3 (1)	159,808	5.9 (4)	3.6 (4)	13.2 (5)	69,573	16.4 (3)	13.8 (5)	113,899	15.7 (5)
SRO	16,199	7.4 (2)	8.8 (4)	64,785	5.1 (2)	3.0 (2)	12.9 (4)	28,249	15.9 (1)	13.4 (2)	46,429	8.5 (3)
WRO	12,306	7.1 (1)	9.3 (5)	51,980	5.3 (3)	3.1 (3)	12.3 (1)	23,401	17.6 (5)	12.9 (1)	37,798	9.0 (4)
Unknown Regions	0	* *	* *	1	* *	* *	* *	1	* *	* *	1	* *
Wisconsin	95,076	8.6	7.6	380,229	5.5	3.4	12.9	167,791	16.4	13.5	273,457	11.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this region's rate to other regions. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies overweight children and ≥ 95th percentile category identifies obese children.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.



Table 7E
2007 - 2009 Pediatric Nutrition Surveillance (1)
Wisconsin

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Region (2)
Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)														
	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤2 Hours TV Viewing/Day (8)		Smoking in Household	
Region	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
NERO	19,377	64.3 (3)	12,965	20.7 (5)	13,283	10.5 (4)	12,145	12.6 (4)	12,180	5.9 (4)	35,294	82.0 (4)	40,688	8.7 (3)
NRO	8,224	63.3 (4)	5,652	20.9 (4)	5,391	10.5 (4)	5,164	14.3 (3)	5,211	7.4 (3)	15,003	84.6 (2)	17,250	8.6 (2)
SERO	43,757	63.2 (5)	29,023	21.9 (3)	29,745	12.0 (2)	26,835	8.6 (5)	27,555	4.0 (5)	75,353	79.4 (5)	88,664	10.8 (5)
SRO	17,753	70.0 (2)	12,117	29.0 (1)	11,746	16.4 (1)	11,267	15.0 (2)	11,207	7.6 (2)	30,424	83.9 (3)	36,682	8.7 (3)
WRO	13,386	71.1 (1)	9,328	23.0 (2)	9,335	11.3 (3)	8,521	15.9 (1)	8,624	7.7 (1)	25,207	85.6 (1)	28,950	7.2 (1)
Unknown Regions	0	* *	0	* *	0	* *	0	* *	0	* *	1	* *	1	* *
Wisconsin	102,497	65.6	69,085	23.0	69,500	12.2	63,932	11.9	64,777	5.7	181,282	82.0	212,235	9.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this region's rate to other regions. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.

(8) Children aged 2 years and older included in the analyses.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

