# **Technical Notes**

# Annual Wisconsin Births and Infant Mortality Report, 2018 (P-01161-22-TN)

Calculations	
Statistical Tests for Trend	2
Population Estimates	3
Race/Ethnicity Classification	s3
Teens	4
Prematurity	4
Prenatal Care Utilization	5
Mortality	5
Perinatal Regions	6
Neighborhood Poverty	6
Urban-Rural Classification	6
Birth Certificate Checklists	7
Limitations of the Data	13
Changes Beginning with 2011	Births
Survey Instrument (Birth Cert	tificate Worksheet) Data Elements 16
Regions	23
Calculations	
Crude Birth Rate =	$\left(\frac{Number\ of\ live\ births}{Total\ population}\right) \times 1,000$
	( Total population )
General Fertility Rate =	$\left(\frac{Number\ of\ live\ births}{Number\ of\ female\ population\ aged\ 15-44}\right)  imes 1,000$
Pregnancy Rate = $\left(\frac{1}{2}\right)$	Number of live births+number of fetal deaths+number of induced abortions $\times 1,000$ Number female population in age group
Age-Specific Birth Rate =	$\left(\frac{\text{Number of live births to mothers in age group}}{\text{Number of female population in age group}}\right) \times 1,000$
D 6 16 E 1111 D 1	/ Number of live births to mothers in race group
Race-Specific Fertility Rate =	$\left(\frac{\textit{Number of live births to mothers in race group}}{\textit{Number of female population aged 15-44 in race group}}\right)  imes 1,000$
Infant Mortality Rate =	(Number of infant deaths) v 1 000
mant mortatity Rate -	$\left(\frac{Number\ of\ inf\ ant\ deaths}{Number\ of\ live\ births}\right) \times 1,000$

 $(Number\ of\ neonatal\ deaths) \times 1,000$ Neonatal Mortality Rate = Number of live births (Number of postneonatal deaths)  $\times 1,000$ Postneonatal Mortality Rate = Number of live births Number of fetal deaths Fetal Death Rate = (Number of fetal deaths + number of neonatal deaths)  $\times 1,000$ Perinatal Mortality Rate = Number of fetal deaths+number of live births (Race 1 infant mortality rate) Racial Infant Mortality Disparity Ratio = Race 2 in fant mortality rate.  $\left(\frac{\text{Mother's prepregnancy weight in pounds}}{1}\right) * 703$ Body Mass Index = (Mother's height in inches)<sup>2</sup>

Before 2006, Wisconsin teen birth rates (the rate of births to mothers aged 15 - 19, and the rate of births to mothers aged 15 - 17) included births to mothers under age 15 in the numerator. The rate was calculated as the number of births to mothers less than 20 (or less than 18) divided by the number of females aged 15 - 19 (or 15 - 17), multiplied by 1,000. Beginning with the 2006 data report, births to mothers aged 10 - 14 were placed in a separate category whenever birth rates were shown.

Similarly, births to mothers aged 45 and older were included in birth rates for women aged 40 - 44 prior to 2006. The result was then divided by the number of females aged 40 - 44 and multiplied by 1,000. Beginning with the 2006 data report, births to women 45 years of age and older were removed from the 40 - 44 birth rate calculations.

In the 2017 report, the teen pregnancy rate calculation underwent a change similar to the teen birth rate change in 2006. The numerator used to include all births, fetal deaths, and induced abortions among females under 20 years. Now, the numerator is limited to females aged 15 - 19 years to match the denominator specifications. The reported induced abortions are defined as abortions that occurred in Wisconsin (residency not confirmed).

National statistics for the general fertility rate<sup>1</sup>, low birthweight<sup>1</sup>, and infant mortality rate<sup>2</sup> were obtained from the Centers for Disease Control and Prevention's (CDC) National Center for Health Statistics.

# Statistical Tests for Trend

The Cochran-Armitage trend test was used for all trend analyses. This modified chi squared test is appropriate for testing the relationship between an ordinal variable (e.g., year) and a dichotomous variable (e.g., infant death) and is robust for testing whether a trend is monotonically increasing or decreasing. Small p-values using this test suggest that the population slope of the linear trend is nonzero.

<sup>1</sup> https://www.cdc.gov/nchs/data/vsrr/report004.pdf

https://wonder.cdc.gov/ucd-icd10.html

All trend tests included data from each individual year within the range indicated. Although Figure 7 displays three-year rolling averages from 2011-2013 to 2015-2017 (5 data points) to reduce visual noise, the statistical tests used the data from each independent year 2011-2017 (7 data points).

# **Population Estimates**

The Department of Health Services, Division of Public Health, Office of Health Informatics produces mid-year population estimates for the counties and state of Wisconsin by age, sex, race, and Hispanic ethnicity for non-census years. These estimates are used to calculate population-based health statistics, such as teen birth rates and other age-specific fertility rates. Estimates from the previous year are used to calculate rates. For example, population estimates from 2016 were used to calculate 2017 teen birth rates.

# **Race/Ethnicity Classifications**

Prior to 2011, only one race and one ethnicity could be reported for each mother. Wisconsin birth reports included race/ethnicity categories of non-Hispanic white, non-Hispanic Black, non-Hispanic American Indian, Hispanic, non-Hispanic Laotian or Hmong, and non-Hispanic other Asian. For the 2011 infant death data, grouping mother's race/ethnicity the same way as in 2010 was attempted. In this report, the race or ethnicity of the mother, rather than the infant, is always used when describing births, birth outcomes, and infant mortality.

Beginning in 2011, the mother could indicate multiple races and multiple ethnicities. The race categories are listed below.

- White
- Black/African American
- American Indian/Alaska Native
- Asian Indian
- Chinese
- Filipino
- Japanese
- Korean
- Vietnamese
- Laotian
- Hmong
- Other Asian
- Native Hawaiian
- Guamanian or Chamorro
- Samoan
- Other Pacific Islander
- Other

The Hispanic ethnicity categories are:

- Mexican, Mexican American, Chicana
- Puerto Rican
- Cuban
- Other

This changed how race and ethnicity were classified for 2011 birth and fetal death data. All live births and fetal deaths were classified by race and Hispanic origin of mother into one of nine categories for reporting purposes:

- Non-Hispanic white alone (shortened to 'White' in the report)
- Non-Hispanic Black/African American alone (shortened to 'Black' in the report)
- Non-Hispanic American Indian/Alaska Native alone (shortened to 'American Indian' in the report)
- Hispanic or Latina (Hispanics/Latinas may be of any race)
- Non-Hispanic Laotian or Hmong alone (shortened to 'Laotian or Hmong' in the report)
- Non-Hispanic other Asian/Pacific Islander alone (includes Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese, Other Asian, Native Hawaiian, Guamanian or Chamorro, Samoan, and Other Pacific Islander; shortened to 'Other Asian' in the report)
- Non-Hispanic Other race alone
- Non-Hispanic Unknown alone
- Non-Hispanic two or more races (shortened to 'Two or more races' in the report)

When calculating birth rates or fertility rates by race/ethnicity, a different classification is used to match the categories available in the bridged-race population estimates published by the <a href="National Center for Health Statistics">National Center for Health Statistics</a> and modified by the <a href="Wisconsin Department of Health Services">Wisconsin Department of Health Services</a>. Population estimates are used to calculate rates (see calculations above). The categories are:

- Non-Hispanic white alone
- Non-Hispanic Black/African American alone
- Non-Hispanic American Indian/Alaska Native alone
- Non-Hispanic Asian alone (includes Asian Indian, Chinese, Filipino, Japanese, Korean, Vietnamese, Laotian, Hmong, Other Asian, Native Hawaiian, Guamanian or Chamorro, Samoan, and Other Pacific Islander)
- Hispanic
- Other/unknown

#### **Teens**

All mothers under 20 years of age are included in teen birth counts. The teen birth rate calculation only includes mothers aged 15 - 19 years. Age-specific birth counts and rates for mothers aged 10 - 14 years, 15 - 17 years, and 18 - 19 years can be found in Table 3-2 of the separate supplemental Birth Tables 2017.

#### **Prematurity**

Births were classified as premature (less than full term) if the gestational age was less than 37 weeks. However, the method for calculating gestational age has changed in recent years. Prior to the 2014 data year, the Wisconsin Interactive Statistics on Health and other DHS birth data reports used an estimate based on the computed difference between the date of reported last normal menses and the date of the infant's birth. Beginning with the 2015 data year, WISH and other DHS birth data reports use an estimate that is based on the attending physician's clinical estimate of gestational age (labeled "Gestational Age Based on Obstetric Estimate"). The CDC's National Center for Health Statistics<sup>3</sup> is transitioning to the obstetric estimate, and WISH and other DHS birth data reports now contain an estimate of gestational age that is calculated in a comparable manner. The CDC's evaluation indicates that,

<sup>&</sup>lt;sup>3</sup> http://www.cdc.gov/nchs/data/nvsr/nvsr64/nvsr64\_05.pdf

compared to the estimate based on the time since the mother's last menstrual period, the obstetric estimate results in a smaller proportion of premature births.

#### Prenatal Care Utilization

The Kotelchuck Adequacy of Prenatal Care Utilization Index takes into account the month that prenatal care began and the number of prenatal visits as reported by the mother, and adjusts for gestational age. It includes categories for adequate plus, adequate, intermediate, and inadequate levels of prenatal care utilization. Adequate plus and adequate categories were combined for the analyses. The number of prenatal visits is compared to the expected number for gestational age, based on the <a href="American College of Obstetricians and Gynecologists">American College of Obstetricians and Gynecologists</a> prenatal care standards for uncomplicated pregnancies. Prenatal care utilization is considered inadequate if the first visit was after month four of the pregnancy or if the actual number of visits was less than half of the recommended number of visits. Prenatal care was considered intermediate if care began in the first four months of pregnancy and the ratio of the actual number of visits to the expected number of visits was greater than or equal to 0.5 and less than 0.8. Prenatal care was adequate if care began in the first four months of pregnancy and the ratio of the actual number of visits to the expected number of visits was greater than or equal to 0.8.

# Mortality

**Infant Deaths:** An infant death is the death of an infant who was born alive but died before its first birthday. The birth-death cohort is not completed until one full year after the end of the birth cohort calendar year. This publication reports mortality data on infants who died during 2017, rather than the mortality experience of the 2017 birth cohort, to allow timelier reporting on infant mortality.

**Neonatal Deaths**: A neonatal death is the death of an infant who was born alive but died within the first 27 days of life (aged <28 days).

**Postneonatal Deaths:** A postneonatal death is the death of an infant who was born alive but died sometime between 28 days to one year of life.

Causes of Death: Causes of death in this report have been coded according to the International Classification of Diseases (ICD), Tenth Edition. Before the 1999 data year, causes of death were coded according to the ICD, Ninth Edition. This change does not affect comparability between years for four of the five leading causes of infant death shown in this report (Conditions Originating in the Perinatal Period; Congenital Malformations, Deformations and Chromosomal Anomalies; Sudden Infant Death Syndrome or SIDS; and Accidents).

**Fetal Deaths:** A fetal death or stillbirth is "a fetus which, after complete expulsion or extraction from the woman, does not breathe or show other evidence of life such as beating of the heart, pulsation of the umbilical cord, or definite movement of the voluntary muscles." By Wisconsin statute, a stillbirth of at least 20 weeks' gestation or 350 grams must be reported. (Note: A fetal death report is not used for induced abortions.) Some stillbirths (fetal deaths) to Wisconsin residents are not included in this publication because some were less than 20 weeks' gestation, some occurred outside Wisconsin, and an unknown number that occurred within the state were not reported.

**Perinatal Deaths**: Includes all reported fetal deaths of 20 or more weeks of gestation and all neonatal deaths (deaths of live-born infants occurring before 28 days of age).

**Maternal Deaths:** Includes all deaths where the cause of death was medically related to the pregnancy. There are pregnancy-associated maternal deaths that are not included in this report.

# **Perinatal Regions**

These tables present birth information for each of the seven perinatal regions in Wisconsin (see Map 2). These regions were informally designated in the early 1970s by the Wisconsin Association for Perinatal Care, in cooperation with organizations representing public and private providers of perinatal services. Perinatal regions represent geographical boundaries within which arrangements exist among providers of perinatal services to facilitate optimal treatment for pregnant women and newborns. Within each region, selected hospitals with the capability to provide neonatal and maternal intensive care have been self-designated as perinatal centers. Women with high-risk pregnancies and deliveries are often referred to or transported to these perinatal centers, as are high-risk babies born outside perinatal centers. Health care providers within the regions also collaborate in other ways to address perinatal health care issues as they arise.

# **Neighborhood Poverty**

The census block groups of the mothers' residences (for births and fetal deaths) or infants' residences (for infant deaths) were obtained by geocoding the residential addresses. Estimates of neighborhood poverty (i.e., the percentage of households with incomes below the federal poverty level for each census block group) were obtained from the U.S. Census Bureau's 2012-2016 American Community Survey five-year estimates (public use detailed table C17002) and applied to the birth and death certificate data. Area level poverty categories were low (0.0-9.9%), moderate-low (10.0-19.9%), moderate-high (20.0-39.9%), and high (40.0-100.0%).

## **Urban-Rural Classification**

The 2017 report is the first year that the <u>NCHS Urban-Rural Classification Scheme for Counties</u> was included, which is based on proximity and economic ties to nearby metropolitan statistical areas (MSAs). The six urban-rural classifications are as follows:

Large central metro: NCHS-defined "central" counties of MSAs of 1 million or more population. Includes:

Milwaukee County

Large fringe metro: NCHS-defined "fringe" counties of MSAs of 1 million or more population. Includes:

Waukesha County Washington County St. Croix County
Kenosha County Ozaukee County Pierce County

Medium metro: Counties within MSAs of 250,000-999,999 population. Includes:

Dane County Douglas County Iowa County
Brown County Oconto County Kewaunee County

Columbia County Green County

Small metro: Counties within MSAS of 50,000 to 249,999 population. Includes:

Racine County
Outagamie County
Uinnebago County
Marathon County
La Crosse County
Sheboygan County
Chippewa County
Calumet County

Rock County Fond du Lac County

*Micropolitan*: Counties in micropolitan statistical areas. Includes:

Walworth CountyPortage CountyMarinette CountyDodge CountySauk CountyLincoln CountyJefferson CountyGrant CountyFlorence CountyManitowoc CountyDunn CountyMenominee County

Wood County Shawano County

Noncore: Counties not within micropolitan statistical areas. Includes:

Waupaca County Vilas County Washburn County Barron County Adams County **Burnett County** Monroe County Taylor County Marguette County Jackson County **Bayfield County** Polk County Oneida County Langlade County Rusk County Clark County Green Lake County Price County Vernon County Richland County **Buffalo County** Trempealeau County Lafayette County Forest County Door County Sawyer County Pepin County Crawford County Iron County Juneau County

Waushara County Ashland County

### **Birth Certificate Checklists**

Beginning in 2011, data on maternal risk factors, pregnancy, delivery, and some dimensions of birth outcomes are collected in the following birth certificate checklists:

- Risk factors in this pregnancy
- Infections present and/or treated during this pregnancy
- Obstetric procedures
- Onset of labor
- Characteristics of labor and delivery
- Method of delivery
- Maternal morbidity
- Abnormal conditions of the newborn
- Congenital anomalies of the newborn

These checklists include data that were not previously collected on the birth certificate and do not include all of the data that was collected prior to 2011.

Definitions for these groups were adapted from the 2003 revision of the U.S. Standard Certificate of Live Birth. See specification for items on the birth certificate at <a href="http://www.cdc.gov/nchs/nvss/vital\_certificate\_revisions.htm">http://www.cdc.gov/nchs/nvss/vital\_certificate\_revisions.htm</a>.

Risk Factors in this Pregnancy

Diabetes Prepregnancy	Glucose intolerance requiring treatment diagnosed prior to this pregnancy.
Diabetes Gestational	Glucose intolerance requiring treatment diagnosed during this pregnancy.
Hypertension Prepregnancy	Elevation of blood pressure above normal for age, gender, and physiological condition diagnosed prior to the onset of this pregnancy.
Hypertension Gestational	Elevation of blood pressure above normal for age, gender, and physiological condition diagnosed during this pregnancy. May include proteinuria (protein in the urine) without seizures or coma and pathologic edema (generalized swelling, including swelling of the hands, legs and face). (PIH, Preeclampsia)
Eclampsia	Gestational hypertension with proteinuria with generalized seizures or coma. May include pathologic edema.
Previous Preterm Birth	History of pregnancy(ies) resulting in a live birth of less than 37 completed weeks of gestation.
Other Previous Poor Pregnancy Outcome	History of pregnancies continuing into the 20 <sup>th</sup> week of gestation (post menstrual age) and resulting in perinatal death (including fetal and neonatal deaths), small-for-gestational age, or intrauterine growth-restricted birth.
Pregnancy Resulted from Infertility Treatment	Any assisted reproduction technique used to initiate the pregnancy.
Infertility Treatment - Fertility-enhancing drugs, artificial insemination, or intrauterine insemination	Any fertility-enhancing drugs (e.g., Clomid, Pergonal), artificial insemination, or intrauterine insemination used to initiate the pregnancy.
Infertility Treatment - Assisted Reproductive Technology	Any assisted reproduction technology (ART)/technical procedures (e.g., IVF, GIFT, ZIFT) used to initiate the pregnancy.
Mother had a Previous Cesarean Delivery	Previous operative birth in which the infant is extracted through an incision in the maternal abdominal and uterine walls.

Infections Present and/or Treated During this Pregnancy

Gonorrhea	A positive test for Neisseria gonorrhoeae present at the time of pregnancy diagnosis or confirmed diagnosis during the pregnancy with or without documentation of treatment.
	Documentation of treatment is adequate if a definitive diagnosis is not present in the available record.

Infections Present and/or Treated During this Pregnancy

Syphilis	A positive test for Treponema pallidum present at the time of pregnancy diagnosis or confirmed diagnosis during the pregnancy with or without documentation of treatment.  Documentation of treatment is adequate if a definitive diagnosis is not present in the available record.
Chlamydia	A positive test for Chlamydia trachomatis present at the time of pregnancy diagnosis or confirmed diagnosis during the pregnancy, with or without documentation of treatment.  Documentation of treatment is adequate if a definitive diagnosis is not present in the available record.
Hepatitis B	A positive test for hepatitis B virus present at the time of pregnancy diagnosis or confirmed diagnosis during the pregnancy with or without documentation of treatment. Documentation of treatment is adequate if a definitive diagnosis is not present in the available record.
Hepatitis C	A positive test for hepatitis C virus present at the time of pregnancy diagnosis or confirmed diagnosis during the pregnancy with or without documentation of treatment. Documentation of treatment is adequate if a definitive diagnosis is not present in the available record.

# **Obstetric Procedures**

• # * * * * * * * * * * * * * * * * * *	
Cervical Cerclage	Circumferential banding or suture of the cervix to prevent or treat passive dilation.
Tocolysis	Administration of any agent with the intent to inhibit preterm uterine contractions to extend the length of the pregnancy.
External Cephalic Version	Attempted conversion of a fetus from a non-vertex presentation to a vertex presentation by external manipulation.
External Cephalic Version Successful	Successful attempted conversion of a fetus from a non-vertex presentation to a vertex presentation by external manipulation.
External Cephalic Version Failed	Failed attempted conversion of a fetus from a non-vertex presentation to a vertex presentation by external manipulation.

# Onset of Labor

Premature Rupture of the Membranes	Spontaneous tearing of the amniotic sac (natural breaking of the "bag of waters"), 12 hours or more before labor begins.
Precipitous Labor	Labor that progresses rapidly and lasts for less than three hours.
Prolonged Labor	Labor that progresses slowly and lasts for 20 hours or more.

**Characteristics of Labor and Delivery** 

Induction of Labor	Initiation of uterine contractions by medical and/or surgical means for the purpose of delivery before the spontaneous
	onset of labor.
Augmentation of Labor	Stimulation of uterine contractions by drug or manipulative technique with the intent to reduce the time to delivery.
Non-vertex Presentation	Includes any non-vertex fetal presentation (e.g., breech, shoulder, brow, face presentations, and transverse lie) in the active phase of labor or at delivery other than vertex.
Steroids for Fetal Lung Maturation Received by the Mother Prior to Delivery	Steroids (glucosteroids) for fetal lung maturation received by the mother prior to delivery. Includes betamethasone, dexamethasone, or hydrocortisone specifically given to accelerate fetal lung maturation in anticipation of preterm delivery.
Antibiotics Received by the Mother During Labor	Antibiotics received by the mother during labor. Includes antibacterial medications given systemically (intravenous or intramuscular) to the mother in the interval between the onset of labor and the actual delivery (Ampicillin, Penicillin, Clindamycin, Erythromycin, Gentamicin, Cefataxime, Ceftriaxone, Vancomycin).
Clinical Chorioamnionitis Diagnosed During Labor or Maternal Temperature ≥38° C (100.4° F)	Clinical chorioamnionitis diagnosed during labor or maternal temperature ≥38° C (100.4° F). A clinical diagnosis of chorioamnionitis during labor made by the delivery attendant. Usually includes more than one of the following: fever, uterine tenderness and/or irritability, leukocytosis, and fetal tachycardia. Any recorded maternal temperature at or above 38° C (100.4° F).
Moderate/Heavy Meconium Staining of the Amniotic Fluid	Moderate/heavy meconium staining of the amniotic fluid. Staining of the amniotic fluid caused by passage of fetal bowel contents during labor and/or delivery which is more than enough to cause a greenish color change of an otherwise thin fluid.
Fetal Intolerance of Labor	Fetal intolerance of labor such that one or more of the following actions was taken: in-utero resuscitation measures, further fetal assessment, or operative delivery. In-utero resuscitative measures such as any of the following: maternal position change, oxygen administration to the mother, intravenous fluid administered to the mother, amnioinfusion, support of maternal blood pressure, and administration of uterine relaxing agents. Further fetal assessment includes any of the following: scalp pH, scalp stimulation, acoustic stimulation. Operative delivery: operative intervention to shorten time to delivery of the fetus such as forceps, vacuum, or cesarean delivery.
Epidural or Spinal Anesthesia During Labor	Epidural or spinal anesthesia during labor. Administration to the mother of a regional anesthetic for control of the pain of labor (i.e., delivery of the agent into a limited space with the distribution of the analgesic effect limited to the lower body).

**Maternal Morbidity** 

	·
Maternal Transfusion	Includes infusion of whole blood or packed red blood cells within the period specified.
Third or Fourth Degree Perineal Laceration	Third degree laceration extends completely through the perineal skin, vaginal mucosa, perineal body and anal sphincter. Fourth degree laceration is all of the above with extension through the rectal mucosa.
Ruptured Uterus	Tearing of the uterine wall.
Unplanned Hysterectomy	Surgical removal of the uterus that was not planned prior to admission for delivery. Includes an anticipated or possible but not definitively planned procedure.
Admission to Intensive Care Unit	Any admission, planned or unplanned, of the mother to a facility or unit designated as providing intensive care.
Unplanned Operating Room Procedure Following Delivery	Any transfer of the mother back to a surgical area for an operative procedure that was not planned prior to the admission for delivery.

# Abnormal Conditions of the Newborn

Assisted Ventilation Required Immediately Following Delivery	Infant given manual breaths with bag and mask or bag and endotracheal tube within the first several minutes from birth for any duration. Excludes free flow oxygen only and laryngoscopy for aspiration of meconium.
Assisted Ventilation Required for More Than Six Hours	Infant given mechanical ventilation (breathing assistance) by any method for more than six hours. Includes conventional, high frequency and/or continuous positive pressure (CPAP).
NICU Admission	Admission into a facility or unit staffed and equipped to provide continuous mechanical ventilatory support for the newborn.
Newborn Given Surfactant Replacement Therapy	Infant given endotracheal instillation of a surface active suspension for the treatment of surfactant deficiency either due to preterm birth or pulmonary injury resulting in decreased lung compliance (respiratory distress). Includes both artificial and extracted natural surfactant.
Antibiotics Received by the Newborn for Suspected Neonatal Sepsis	Any antibacterial drug, received by the newborn for suspected neonatal sepsis, given systemically (intravenous or intramuscular) (e.g., penicillin, ampicillin, gentamicin, cefotaxime, etc.).

# Abnormal Conditions of the Newborn

Seizure or Serious Neurologic Dysfunction	Seizure defined as any involuntary repetitive convulsive movement or behavior. Serious neurologic dysfunction defined as severe alteration of alertness such as obtundation, stupor, or coma (i.e., hypoxic-ischemic encephalopathy). Excludes lethargy or hypotonia in the absence of other neurologic findings. Exclude symptoms associated with central nervous system (CNS) congenital anomalies.
Significant Birth Injury	Skeletal fracture(s), peripheral nerve injury, or soft tissue (or solid organ) hemorrhage which requires intervention. Present immediately following delivery or manifesting soon after delivery. Includes any bony fracture or weakness or loss of sensation, but excludes fractured clavicles and transient facial nerve palsy. Soft tissue hemorrhage requiring evaluation and/or treatment includes sub-galeal (progressive extravasation within the scalp) hemorrhage, giant cephalohematoma, extensive truncal, facial and/or extremity ecchymosis accompanied by evidence of anemia and/or hypovolemia and or hypotension. Solid organ hemorrhage includes subcapsular hematoma of the liver, fractures of the spleen, or adrenal hematoma. All require confirmation by diagnostic imaging or exploratory laparotomy.

Congenital Anomalies of the Newborn

Congenital Anomalies of the Newborn		
Anencephaly	Partial or complete absence of the brain and skull. Also called anencephalus, acrania, or absent brain. Babies with craniorachischisis (anencephaly with contiguous spine defect) should also be included in this category.	
Meningomyelocele/Spina Bifida	Spina bifida refers to herniation of the meninges and/or spinal cord tissue through a bony defect of spine closure.  Meningomyelocele refers to herniation of meninges and spinal cord tissue. Babies with meningocele (herniation of meninges without spinal cord tissue) should also be included in this category. Both open and closed (covered with skin) lesions should be included. Do not include spina bifida occulta (a midline bony spinal defect without protrusion of the spinal cord or meninges).	
Cyanotic Congenital Heart Disease	Congenital heart defects which cause cyanosis. Includes but is not limited to TGA—transposition of the great arteries (vessels), TOF—teratology of Fallot, pulmonary or pulmonic valvular atresia, tricuspid atresia, truncus arteriosus, TAPVR - total/partial anomalous pulmonary venous return with or without obstruction.	
Congenital Diaphragmatic Hernia	Defect in the formation of the diaphragm allowing herniation of abdominal organs into the thoracic cavity.	

Congenital Anomalies of the Newborn

Congenital Anomalies	
Omphalocele	A defect in the anterior abdominal wall, accompanied by herniation of some abdominal organs through a widened umbilical ring into the umbilical stalk. The defect is covered by a membrane (different from gastroschisis, see below), although this sac may rupture. Also called exomphalos. Do not include umbilical hernia (completely covered by skin) in this category.
Gastroschisis	An abnormality of the anterior abdominal wall, lateral to the umbilicus, resulting in herniation of the abdominal contents directly into the amniotic cavity. Differentiated from omphalocele by the location of the defect and absence of a protective membrane.
Limb Reduction Defect	Complete or partial absence of a portion of an extremity secondary to failure to develop. Excluding congenital amputation and dwarfing syndromes.
Cleft Lip With or Without Cleft Palate	Cleft lip with or without cleft palate is defined as incomplete closure of the lip. May be unilateral, bilateral, or median; all should be included in this category.
Cleft Palate Alone	Cleft palate is defined as incomplete fusion of the palatal shelves. This may be limited to the soft palate or may also extend into the hard palate. Cleft palate in the presence of cleft lip should be included in the "Cleft Lip with or without cleft palate" category, rather than here.
Down Syndrome	Trisomy 21.
Down Syndrome - Karyotype Confirmed	Karyotype confirmed for Trisomy 21.
Down Syndrome - Karyotype Pending	Karyotype pending for Trisomy 21.
Suspected Chromosomal Disorder	Includes any constellation of congenital malformations resulting from or compatible with known syndromes caused by detectable defects in chromosome structure.
Suspected Chromosomal Disorder - Karyotype Confirmed	Karyotype confirmed for suspected chromosomal disorder.
Suspected Chromosomal Disorder - Karyotype Pending	Karyotype pending for suspected chromosomal disorder.
Hypospadias	Incomplete closure of the male urethra resulting in the urethral meatus opening on the ventral surface of the penis.

# Limitations of the Data

Vital records information has some limitations related to completeness, accuracy, and quality control procedures. The reporting of live births and infant deaths is considered to be essentially complete for births occurring in Wisconsin. A cooperative exchange program between all states has been in operation for several decades to allocate birth and death

certificates to the state of residence. Although this exchange program is not totally complete, the incompleteness is not of sufficient magnitude to seriously affect statistics on natality and infant mortality.

The statistics in the report reflect 2017 birth record data as of August, 2018. Any corrections made to individual 2017 birth records during quality control procedures after August 2018 are not included in this report.

# Changes Beginning with 2011 Births

On January 1, 2011, the <u>Wisconsin State Vital Records Office</u> began collecting birth certificate information based on the 2003 U.S. Standard Certificate of Live Birth using a new electronic system. The new electronic system performs edit checks as hospital and Vital Records staff enter birth certificate information. Staff in the Office of Health Informatics perform an annual review of birth certificates and death certificates for inconsistencies and outlying values. Previous to January 1, 2011, Wisconsin had been collecting birth certificate information based on the 1989 U.S. Standard Certificate of Live Birth.

There were many changes made to birth data collection when Wisconsin began using the 2003 U.S. Standard Certificate of Live Birth. Extreme caution should be used when analyzing data based on new data fields that have not been previously collected. Wisconsin has no trend data to validate these data. Wisconsin Vital Records analysts have compared this new data to national data, but because many states are new to collecting this information, the data quality is in question. For example, according to the Wisconsin Department of Health Services, Division of Medicaid Services, almost 50 percent of births in 2014 were paid for by Wisconsin Medicaid/BadgerCare, but Medicaid/BadgerCare was indicated as the principal payer source on only 38 percent of 2014 birth certificates. Also, in some cases Wisconsin birth hospitals are reporting private insurance and some Medicaid/BadgerCare under the "Other" category and then naming the actual insurance carrier.

**New Data Fields**: New fields include but are not limited to: date of first prenatal care visit, date of last prenatal care visit, mother married at any time in her life, mother's height, mother's prepregnancy weight, mother's weight at delivery, mother received WIC food for herself during the pregnancy, principal source of payment for this delivery, APGAR score at 10 minutes, infant breastfed at discharge, mother lives with someone who smokes.

**Data Fields with Changed Values:** Fields that continue to be collected, but with different values, include but are not limited to: mother's education, mother's race and ethnicity, maternal smoking, infant transferred following delivery. Specific changes for some of these fields are listed below:

- 1. Prior to 2011, the number of years of education completed was reported. Beginning in 2011, education was collected using the following categories:
  - 8<sup>th</sup> Grade or Less
  - 9<sup>th</sup> 12<sup>th</sup> Grade, No Diploma
  - High School Graduate or GED Completed
  - Some College Credit, No Degree
  - Associate Degree (e.g., AA, AS)
  - Bachelor's Degree (e.g., BA, AB, BS)
  - Master's Degree (e.g., MA, MS, MEng, MEd, MSW, MBA)
  - Doctorate or Professional Degree (e.g., PhD, EdD, MD, DDS, DVM, JD)

- 2. For race changes, see Race/Ethnicity Classifications.
- 3. Prior to 2011, mothers were asked if they smoked during pregnancy and the average number of cigarettes smoked per day. Beginning in 2011, the average number of cigarettes per day was collected using the following categories:
  - Three Months before Pregnancy
  - First Three Months of Pregnancy
  - Second Three Months of Pregnancy
  - Third Trimester of Pregnancy
- 4. Beginning in 2011, the infant transferred field includes only infants that were transferred externally within 24 hours of the delivery. Neonatal Intensive Care Unit (NICU) admission is collected under the Abnormal Conditions of the Newborn checklist. Conversely, before 2011, this field was used to indicate whether an infant was transferred and to which facility the infant was transferred. This included transfers to high-risk nurseries internally or to external facilities.
- 5. Beginning in 2011, month prenatal care began is calculated using the date of last normal menses and the date of the first prenatal care visit. If either of these fields is unknown, the date of birth and the clinician's estimate of gestational age are used to calculate the month prenatal care began. This change has caused a drop in the reported number of mothers with first-trimester prenatal care. In previous years, the month prenatal care began was self-reported.
- 6. "Method of delivery" categories changed in 2011. The categories are vaginal/spontaneous, vaginal/forceps, vaginal/vacuum, cesarean, and unknown. Additional fields collected under the Risk Factors checklist include asking if the mother had any previous cesarean deliveries, and, if so, how many. The field for previous cesarean deliveries had to be combined with the field for final route and method of delivery to calculate vaginal birth after cesarean (VBAC), primary C-section, and repeat C-section. However, there are birth records for which the previous cesarean deliveries item was unknown and cesarean was indicated as the final route and method of delivery. Those records were categorized as C-section. Previous to 2011, the categories for method of delivery were: spontaneous, VBAC, forceps, primary C-section, repeat C-section, other, and unknown.

Extreme caution should be used when comparing data from these fields over time as they are not directly comparable to their pre-2011 counterparts. Wisconsin Vital Records staff recommends that no attempt be made to compare data over time based on these fields.

**Data Fields Eliminated**: Fields that were removed include but are not limited to: APGAR score at one minute, alcohol use during pregnancy, and the birth checklist fields listed below.

- Anemia
- Cardiac Disease
- Acute or Chronic Lung Disease
- Genital Herpes
- Other STD
- Hydramnios/Oligohydramnios

- Hemoglobinopathy
- Incompetent Cervix
- Previous Infant 4000+ Grams
- Renal Disease
- Rh Sensitization
- Uterine Bleeding

- Other Medical Risk Factor (Specify)
- Amniocentesis
- Electronic Fetal Monitoring
- Ultrasound
- Postpartum Sterilization
- Other Obstetric Procedures (Specify)
- Abruptio Placenta
- Placenta Previa
- Seizures During Labor
- Dysfunctional Labor
- Cord Prolapse
- Anesthetic Complications
- Other Events of Labor and/or Delivery (Specify)
- Newborn Anemia
- Hyaline Membrane Disease (RDS)
- Meconium Aspiration Syndrome
- Seizures
- Other Abnormal Conditions of the Newborn (Specify)
- Hydrocephalus

- Microcephalus
- Other Central Nervous System Anomalies (Specify)
- Heart Malformations
- Other Circulatory/Respiratory Anomalies (Specify)
- Rectal Atresia/Stenosis
- Tracheo-esophageal Fistula/Esophageal Atresia
- Other Gastrointestinal Anomalies (Specify)
- Malformed Genitalia
- Renal Agenesis
- Other Urogenital Anomalies (Specify)
- Polydactyly/Syndactyly/Adactyly
- Club Foot
- Other Musculoskeletal/Integumental Anomalies (Specify)
- Other Chromosomal Anomalies (Specify)
- Other Congenital Anomalies (Specify)

# Survey Instrument (Birth Certificate Worksheet) Data Elements

Below is a list of data elements collected on the birth certificate worksheet. Wisconsin does not publish the actual survey instrument in order to safeguard the data. Note: On this list, "n" means it was a new field beginning with 2011, and "r" means the field was revised beginning with 2011.

Mother's Medical Record Number<sup>n</sup> Newborn's Medical Record Number<sup>n</sup>

Child's Name

First

Middle

Last

Suffix

Mother's Current Legal Name

First

Middle

Last

Suffix

Mother's Name Prior to First Marriage

First

Middle

Last

Suffix

Mother's Birthplace

Country

U.S. State/U.S. territory

Mother's Date of Birth

Mother's Hispanic Origin<sup>r</sup>

```
Not Spanish/Hispanic/Latina
       Mexican, Mexican American, Chicana
       Puerto Rican
       Cuban
       Other
       Other Specify
       Unknown
Mother's Racer
       White
       Black or African American
       American Indian or Alaska Native
               American Indian or Alaska Native Specify Tribe 1
               American Indian or Alaska Native Specify Tribe 2
       Asian Indian
       Chinese
       Filipino
       Japanese
       Korean
       Vietnamese
       Laotian
       Hmong
       Other Asian
               Other Asian Specify 1
               Other Asian Specify 2
       Native Hawaiian
       Guamanian or Chamorro
       Samoan
       Other Pacific Islander
               Other Pacific Islander Specify 1
               Other Pacific Islander Specify 2
       Other
               Other Specify 1
               Other Specify 2
       Unknown
Mother's Height
       Feetn
       Inches<sup>n</sup>
Mother's Prepregnancy Weight<sup>n</sup>
Mother's Education<sup>r</sup>
Cigarette Smoking Before and During Pregnancy<sup>r</sup>
       Three Months Prior to Pregnancy
       First Three Months of Pregnancy
       Second Three Months of Pregnancy
       Third Trimester of Pregnancy
Does the Mother Live with Someone who Smokes<sup>n</sup>
Did Mother get WIC Food for Herself<sup>n</sup>
Mother's Marital Status
       Has the Mother Ever Been Legally Married<sup>n</sup>
       Was the Mother Married at Any Time During this Pregnancy
       Paternity Acknowledgement Completed in Hospital<sup>n</sup>
```

```
Mother's SSN
SSN Requested for Child
Mother's Country of Residence<sup>r</sup>
Mother's State of Residence
Mother's County of Residence
Mother's City/Village/Township of Residence
        City/Village/Township Indicator
Mother's Residence Address
        Street Name and Number<sup>n</sup>
        Type of Apartment<sup>n</sup>
        Apartment Number<sup>n</sup>
        Zip Code<sup>n</sup>
Mother's Mailing Address
        In Care of Indicator<sup>n</sup>
        In Care of Text<sup>n</sup>
        Address
        Type of Apartment
        Apartment Number
        City
        State
        Zip Code
Mother's Phone Number<sup>n</sup>
Mother's Type of Phone<sup>n</sup>
Husband's Current Legal Name
        First
        Middle
        Last
        Suffix
Husband's Birth Name
        First<sup>n</sup>
        Middle<sup>n</sup>
        Last<sup>n</sup>
        Suffix<sup>n</sup>
Husband's Birthplace
        Country
        US State/US territory
Husband's Date of Birth
Husband's Hispanic Origin<sup>r</sup>
        Not Spanish/Hispanic/Latino
        Mexican, Mexican American, Chicano
        Puerto Rican
        Cuban
        Other
        Other Specify
        Unknown
Husband's Race<sup>r</sup>
        White
        Black or African American
        American Indian or Alaska Native
                American Indian or Alaska Native Specify Tribe 1
```

```
American Indian or Alaska Native Specify Tribe 2
       Asian Indian
       Chinese
       Filipino
       Japanese
       Korean
       Vietnamese
       Laotian
       Hmong
       Other Asian
               Other Asian Specify 1
               Other Asian Specify 2
       Native Hawaiian
       Guamanian or Chamorro
       Samoan
       Other Pacific Islander
               Other Pacific Islander Specify 1
               Other Pacific Islander Specify 2
       Other
               Other Specify 1
               Other Specify 2
       Unknown
Husband's Education<sup>r</sup>
Husband's SSN
Type of Birth Facility
Facility Name
State of Birth
County of Birth
City/Village/Township of Birth
City/Village/Township Indicator of Birth
Address of Birth
Zip Code of Birth
Facility NPI<sup>n</sup>
Did Mother Receive Prenatal Care
Date of First Prenatal Care Visit<sup>n</sup>
Date of Last Prenatal Care Visit<sup>n</sup>
Total Number of Prenatal Visits for this Pregnancy
Date of Last Normal Menses
Number of Previous Live Births Now Living
Number of Previous Live Births Now Dead
Date of Last Live Birth
Number of Other Pregnancy Outcomes - spontaneous or induced losses or ectopic pregnancies<sup>n</sup>
Date of Last Other Pregnancy Outcome<sup>n</sup>
Risk Factors In this Pregnancy
       None
       Diabetes
               prepregnancy
               gestational
       Hypertension
               prepregnancy
```

```
gestational
        Eclampsia
        Previous Preterm Birth<sup>r</sup>
        Other Previous Poor Pregnancy Outcome<sup>n</sup>
        Pregnancy resulted from infertility treatment<sup>n</sup>
                 Fertility Enhancing Drugs, AI or Intrauterine Insemination<sup>n</sup>
                 Assisted Reproductive Technology<sup>n</sup>
        Mother had a previous cesarean delivery<sup>n</sup>
                 Number of Previous C-sections<sup>n</sup>
        Unknown
Infections Present and/or Treated During This Pregnancy
        Gonorrhear
        Syphilis<sup>r</sup>
        Chlamydia<sup>r</sup>
        Hep B<sup>n</sup>
        Hep C<sup>n</sup>
        Unknown
Obstetric Procedures
        None
        Cervical Cerclage<sup>n</sup>
        Tocolysis
        External Cephalic Version<sup>n</sup>
                Successful<sup>n</sup>
                Failedn
        Unknown
Onset of Labor
        None
        PROM
        Precipitous Labor
        Prolonged Labor
        Unknown
Child's Date of Birth
Time of Birth
Principal Source of Payment for this Delivery<sup>n</sup>
Mother Transferred for Maternal Medical or Fetal Indications for Delivery
Mother Transferred From Facility Name<sup>n</sup>
Attendant's Name
        First
        Last
Attendant's Title
Attendant's License Number
Attendant's NPI<sup>n</sup>
Mother's Weight at Delivery<sup>n</sup>
Characteristics of Labor and Delivery
        None
        Induction of labor
        Augmentation of labor
        Non-Vertex Presentation<sup>r</sup>
```

Steroids (glucocorticoids) for fetal lung maturation received by the mother prior to delivery<sup>n</sup>

Antibiotics received by the mother during labor<sup>n</sup>

Clinical Chorioamnionitis diagnosed during labor or maternal temperature  $\geq 38^{\circ}C(100.4^{\circ}F)$ 

Moderate/Heavy Meconium Staining of the Amniotic Fluid

Fetal intolerance of labor such that one or more of the following actions was taken: in-utero resuscitative measures, further fetal assessment, or operative delivery<sup>r</sup>

Epidural or Spinal Anesthesia During Labor<sup>n</sup>

Unknown

# Method of Delivery

Was Delivery with Forceps Attempted but Unsuccessful<sup>n</sup>

Was Delivery with Vacuum Extraction Attempted but Unsuccessful<sup>n</sup>

Fetal Presentation at Birth

Final Route and Method of Delivery

If cesarean, was a trial of labor attempted<sup>n</sup>

#### Maternal Morbidity

None

Maternal Transfusion<sup>n</sup>

Third or Fourth Degree Perineal Laceration<sup>n</sup>

Ruptured Uterus<sup>n</sup>

Unplanned Hysterectomy<sup>n</sup>

Admission to Intensive Care Unit<sup>n</sup>

Unplanned Operating Room Procedure Following Delivery<sup>n</sup>

Unknown

Birthweight

Length at Birth

Infant's Sex

Obstetric Estimate of Gestation

APGAR at 5 Minutes

APGAR at 10 Minutes (if APGAR at 5 minutes is less than 6)<sup>n</sup>

Plurality

If Not Single Birth, Order

Number of Infants Born Alive (multiple births only)<sup>n</sup>

Abnormal Conditions of the Newborn

None

Assisted Ventilation Required Immediately Following Delivery<sup>n</sup>

Assisted Ventilation Required for More Than 6 Hours<sup>n</sup>

NICU Admission<sup>n</sup>

Newborn Given Surfactant Replacement Therapy<sup>n</sup>

Antibiotics Received by the Newborn for Suspected Neonatal Sepsis<sup>n</sup>

Seizure or Serious Neurologic Dysfunction

Significant Birth Injury (Skeletal Fracture(s), Peripheral Nerve Injury, and/or Soft

Tissue/Solid Organ Hemorrhage Which Requires Intervention)

Unknown

# Congenital Anomalies of the Newborn

None

Anencephaly

Meningomyelocele/Spina bifida

Cyanotic Congenital Heart Disease<sup>n</sup>

Congenital Diaphragmatic Hernia Omphalocele<sup>r</sup> Gastroschisis<sup>r</sup> Limb Reduction Defect<sup>n</sup> Cleft Lip with or without Cleft Palate<sup>r</sup> Cleft Palate alone<sup>n</sup> Down Syndrome Karyotype Confirmed<sup>n</sup> Karyotype Pending<sup>n</sup> Suspected Chromosomal Disorder<sup>r</sup> Karyotype Confirmed<sup>n</sup>

Karyotype Pending<sup>n</sup>

Hypospadias<sup>n</sup> Unknown

Is the Infant Being Breastfed at Discharge<sup>n</sup> Newborn Screening Blood Card Number<sup>n</sup> Vaccine Administered to Infant 1<sup>n</sup> Date Vaccine Administered to Infant 1<sup>n</sup> Vaccine Administered to Infant 2<sup>n</sup> Date Vaccine Administered to Infant 2<sup>n</sup> Was Infant Transferred Within 24 Hours of Delivery If Infant Transferred Name of Facility Transferred To<sup>r</sup> Infant Died Infant's Date of Death

<sup>&</sup>lt;sup>n</sup> = New field from 2011

r = Revised field from 2011

# **Regions**

Map 1. Wisconsin Department of Health Services Regions, 2018



Source: Wisconsin Department of Health Services, Division of Public Health, Office of Health Informatics.

Notes: Vernon County was reassigned from the Western to the Southern Region effective February 2004.

This change is reflected in the data presented for 2003 and subsequent years. Comparisons with regional data in previous publications may be affected.

Map 2. Wisconsin Perinatal Regions, 2018



Source: Wisconsin Department of Health Services, Division of Public Health, Office of Health Informatics.