

**FORWARDHEALTH
PRIOR AUTHORIZATION DRUG ATTACHMENT FOR XYREM AND XYWAV**

INSTRUCTIONS: Type or print clearly. Before completing this form, read the Prior Authorization Drug Attachment for Xyrem and Xywav Instructions, F-01430A. Providers may refer to the Forms page of the ForwardHealth Portal at <https://www.forwardhealth.wi.gov/WIPortal/Subsystem/Publications/ForwardHealthCommunications.aspx?panel=Forms> for the completion instructions.

Pharmacy providers are required to have a completed Prior Authorization Drug Attachment for Xyrem and Xywav form signed and dated by the prescriber before submitting a prior authorization (PA) request on the Portal, by fax, or by mail. Providers may call Provider Services at 800-947-9627 with questions.

SECTION I – MEMBER INFORMATION

1. Name – Member (Last, First, Middle Initial)

2. Member ID Number

3. Date of Birth – Member

SECTION II – PRESCRIPTION INFORMATION

4. Drug Name

5. Drug Strength

6. Date Prescription Written

7. Directions for Use

8. Refills

9. Name – Prescriber

10. National Provider Identifier – Prescriber

11. Address – Prescriber (Street, City, State, Zip+4 Code)

12. Phone Number – Prescriber

SECTION III A – CLINICAL INFORMATION

Note: A copy of the member's current medical records that support a clinical correlation between the member's test results and the member's medical condition of narcolepsy with cataplexy or narcolepsy without cataplexy must be submitted with the PA request, including the following:

- Test results and provider interpretation for the polysomnogram) and Multiple Sleep Latency Test (MSLT)
- For members with excessive daytime sleepiness (EDS), a copy of the Epworth Sleepiness Scale (ESS) questionnaire, Maintenance of Wakefulness Test (MWT), or MSLT
- For renewal PA requests, medical record documentation demonstrating clinical improvement, including a decrease in cataplexy or a decrease in the member's daytime sleepiness, supported by an ESS questionnaire, MWT, or MSLT

13. Diagnosis Code and Description



14. Does the member have narcolepsy with cataplexy? Yes No

If yes, indicate in the space below the cataplexy symptoms experienced by the member and how frequently they occur.

15. Does the member have narcolepsy without cataplexy? Yes No

16. Is the member 7 years of age or older? Yes No

17. Does the member have a succinic semialdehyde dehydrogenase deficiency? Yes No

18. As required by the Xywav and Xyrem Risk Evaluation and Mitigation Strategy (REMS) Program:

- Has the prescriber counseled the member on the contraindication between Xyrem or Xywav and alcohol? Yes No
- Has the member agreed to be abstinent from alcohol while being treated with Xyrem or Xywav? Yes No

19. Indicate which symptom(s) of narcolepsy Xyrem or Xywav is being used to treat.

- Cataplexy
- Excessive Daytime Sleepiness
- Other _____

20. Does the member have a history of substance abuse, addiction, or diversion? Yes No

21. Is the member taking any sedative hypnotics? Yes No

22. Is the member taking central nervous system (CNS) depressants (for example, anxiolytics, barbiturates, or opioids)? Yes No

If yes, indicate the CNS depressants and daily doses.

1. _____
2. _____
3. _____

Are any of the above listed CNS depressants contributing to the member's daytime sleepiness? Yes No

If no, indicate how the prescriber evaluated the CNS depressants and determined they are not contributing to the member's daytime sleepiness.

23. Has the member had an overnight polysomnogram sleep study followed by an MSLT? Yes No

24. Does the member have EDS that interferes with normal activities on a daily basis? Yes No

25. Has the member completed an ESS questionnaire, MWT, or MSLT? Yes No

-
26. Has the prescriber ruled out or treated the member for each of the following potential causes of EDS? Yes No
- Other sleep disorders including sleep apnea
 - Chronic pain or illness that disrupts normal sleep patterns
 - Mood disorders such as depression
 - Caffeine or nicotine use causing poor quality of nighttime sleep

-
27. Has the member experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction with a stimulant? Yes No

If yes, list the stimulant and dose, specific details about the unsatisfactory therapeutic response or clinically significant adverse drug reaction, and the approximate dates the stimulant was taken in the space provided.

-
28. Does the member have a medical condition(s) preventing the use of a stimulant? Yes No

If yes, list the medical condition(s) that prevents the use of a stimulant in the space provided.

-
29. Is there a clinically significant drug interaction(s) between another medication(s) the member is taking and stimulants? Yes No

If yes, list the medication(s) and drug interaction(s) in the space provided.

-
30. Has the member experienced an unsatisfactory therapeutic response after the medication has been titrated to a maximum recommended daily dose or experienced a clinically significant adverse drug reaction with armodafinil or modafinil? Yes No

If yes, list the drug and dose, specific details about the unsatisfactory therapeutic response or clinically significant adverse drug reaction, and the approximate dates armodafinil or modafinil or were taken in the space provided.

-
31. Does the member have a medical condition(s) preventing the use of armodafinil or modafinil? Yes No

If yes, list the medical condition(s) that prevents the use of armodafinil or modafinil in the space provided.

32. Is there a clinically significant drug interaction(s) between another medication(s) the member is taking and armodafinil or modafinil? Yes No

If yes, list the medication(s) and drug interaction(s) in the space provided.

33. Has the member experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction with tricyclic antidepressant (TCA), selective serotonin reuptake inhibitor (SSRI), or serotonin norepinephrine reuptake inhibitor (SNRI)? Yes No

If yes, list the TCA, SSRI, or SNRI and dose, specific details about the unsatisfactory therapeutic response or clinically significant adverse drug reaction, and the approximate dates the TCA, SSRI, or SNRI was taken in the space provided.

SECTION III B – ADDITIONAL CLINICAL INFORMATION FOR XYWAV PA REQUESTS

34. PA requests for Xywav must include detailed clinical justification for prescribing Xywav instead of Xyrem. This clinical information must document why the member cannot use Xyrem, including why it is medically necessary that the member receive Xywav instead of Xyrem.

SECTION IV – AUTHORIZED SIGNATURE

35. **SIGNATURE** – Prescriber

36. Date Signed

SECTION V – ADDITIONAL INFORMATION

37. Include any additional information in the space below. Additional diagnostic and clinical information explaining the need for the drug requested may also be included here.
