

INFORMED CONSENT FOR MEDICATION

Dosage and / or Side Effect information last revised on 05/20/2020

Completion of this form is voluntary. If informed consent is not given, the medication cannot be administered without a court order unless in an emergency.

This consent is maintained in the client's record and is accessible to authorized users.

Name – Patient / Client (Last, First MI)		ID Number	Living Unit	Date of Birth
Name – Individual Preparing This Form		Name – Staff Contact		Name / Telephone Number – Institution

MEDICATION CATEGORY	MEDICATION	RECOMMENDED DAILY TOTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE
Alpha Adrenergic Agonist Antihypertensive	Catapres, Duraclon, Kapvay (clonidine)	Oral: 0.025mg-2.4mg	

The anticipated dosage range is to be individualized, may be above or below the recommended range but no medication will be administered without your informed and written consent.

Recommended daily total dosage range of manufacturer, as stated in *Physician's Desk Reference* (PDR) or another standard reference.

This medication will be administered Orally Injection Other – Specify:

1. Reason for Use of Psychotropic Medication and Benefits Expected (note if this is 'Off-Label' Use)

Include DSM-5 diagnosis or the diagnostic impression ("working hypothesis.")

2. Alternative mode(s) of treatment other than OR in addition to medications include

Note: Some of these would be applicable only in an inpatient environment.

- | | |
|---|---|
| <input type="checkbox"/> Environment and/or staff changes | <input type="checkbox"/> Rehabilitation treatments/therapy (OT, PT, AT) |
| <input type="checkbox"/> Positive redirection and staff interaction | <input type="checkbox"/> Treatment programs and approaches (habilitation) |
| <input type="checkbox"/> Individual and/or group therapy | <input type="checkbox"/> Use of behavior intervention techniques |

Other Alternatives:

3. Probable consequences of NOT receiving the proposed medication are

Impairment of Work Activities Family Relationships Social Functioning

Possible increase in symptoms leading to potential

- | | |
|--|--|
| <input type="checkbox"/> Use of seclusion or restraint | <input type="checkbox"/> Limits on recreation and leisure activities |
| <input type="checkbox"/> Limits on access to possessions | <input type="checkbox"/> Intervention of law enforcement authorities |
| <input type="checkbox"/> Limits on personal freedoms | <input type="checkbox"/> Risk of harm to self or others |
| <input type="checkbox"/> Limit participation in treatment and activities | |

Other Consequences:

Note: These consequences may vary depending upon whether or not the individual is in an inpatient setting. It is also possible that in unusual situations, little or no adverse consequences may occur if the medications are not administered.

See Page 2

Client Initial _____ Date _____

4. Possible side effects, warnings, and cautions associated with this medication are listed below. This is not an all-inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text, such as the PDR. As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects in order to enhance care and treatment.

Continued – Possible side effects, warnings, and cautions associated with this medication.

Most Common Side Effects: hypotension, skin rash, dry mouth, drowsiness, dizziness, headache, constipation, and fatigue.

Check with your doctor immediately if any of the following side effects occur: Signs and symptoms of overdose: Difficulty in breathing; dizziness (extreme) or faintness; feeling cold; pinpoint pupils of eyes; slow heartbeat; unusual tiredness or weakness (extreme).

Less Common Side Effects: decreased sexual ability; dizziness, lightheadedness, or fainting, especially when getting up from a lying or sitting position; dry, itching, or burning eyes; loss of appetite; nausea or vomiting; nervousness; mental depression; swelling of feet and lower legs.

Rare Side Effects: paleness or cold feeling in fingertips and toes; vivid dreams or nightmares. These side effects may go away during treatment as your body adjusts to the medicine. However, check with your doctor if any of the listed side effects continue or are bothersome.

Caution:

- Do not suddenly stop taking this medication. This medication should be tapered (stopped slowly). Please consult your physician before stopping this medication.
- When first starting this medication, it can make you dizzy while standing up. Take extra caution by standing slowly from a seated or lying position.
- This medication can impair physical or mental abilities; use caution about performing tasks which require mental alertness, such as driving, operating machinery, or anything else that could be dangerous if you are not alert.

Warning

Withdrawal: Do not discontinue therapy without consulting your physician. Sudden cessation of clonidine treatment has, in some cases, resulted in symptoms such as nervousness, agitation, headache, and tremor accompanied or followed by a rapid rise in blood pressure and elevated catecholamine concentrations in the plasma. The likelihood of such reactions appears to be greater after administration of higher doses or taking this medication with beta-blocker treatment and special caution is therefore advised in these situations. Rare instances of hypertensive encephalopathy (symptoms of headache, vomiting, confusion, or balance issues), cerebrovascular accidents (stroke), and death have been reported after clonidine withdrawal.

Because children commonly have gastrointestinal illnesses that lead to vomiting, they may be particularly susceptible to hypertensive episodes resulting from abrupt inability to take medication.

See standard reference text for an all-inclusive list of side effects.

By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather, it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager, or psychologist.
4. I have the right to request a review at any time of my record, pursuant to § 51.30(4)(d) or § 51.30(5)(b).
5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager, or agency/facility client rights specialist may be contacted for assistance.
6. My consent permits the dose to be changed within the **anticipated dosage range** without signing another consent.
7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s), and the probable consequences that may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate, and complete.
8. This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The need for and continued use of this medication will be reviewed at least quarterly by the Interdisciplinary Team. The goal, on behalf of the client, will be to arrive at and maintain the client at the minimum effective dose.

SIGNATURES

DATE SIGNED

Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client <input type="checkbox"/> Self <input type="checkbox"/> Parent <input type="checkbox"/> Guardian (POA-HC)	
Staff Present at Oral Discussion	Title	

Client / Parent of Minor / Guardian (POA-HC) Comments

As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.

Verbal Consent

Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received <input type="checkbox"/> Yes <input type="checkbox"/> No
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received