DEPARTMENT OF HEALTH SERVICES

Division of Mental Health and Substance Abuse Services F-24277 (07/2015)

STATE OF WISCONSIN 42 CFR483.420(a)(2) DHS 134.31(3)(o) DHS 94.03 & 94.09 §§ 51.61(1)(g) & (h)

Client Initial _____ Date ____

INFORMED CONSENT FOR MEDICATION

Dosage and / or Side Effect information last revised on 08/08/2016

This consent is maintained in the client's record and is accessible to at Name – Patient / Client (Last, First MI)			ID Number	Living Unit	Date of Birth	
rame rationer short (Last, rinser	v <i>)</i>		12 Number	Living Offic	Bate of Birtin	
Name – Individual Preparing This Form Name		Name – Staff Co	ntact	Name / Telephone N	Name / Telephone Number – Institution	
MEDICATION CATEGORY	MEDI	CATION		ECOMMENDED DTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE	
Sedative, Hypnotic	Ambien; Ambien CR Edular; Intermezzo; Zolpimist (zolpidem)		5mg to 10mg (5mg in elderly or women) CR: 6.25mg to 12.5mg (6.5mg in elderly) Inermezzo: 1.75 mg for women, 3.5 mg for men Zolpimist: 10 mg daily (elderly 5 mg daily)		men	
The anticipated dosage range is to be without your informed and written con Recommended daily total dosage raths medication will be administered	onsent. inge of manufacti	•		ference (PDR) or another st		
Reason for Use of Psychotrop Include DSM-5 diagnosis or the control of the c			ted (note if this is	'Off-Label' Use)		
2. Alternative mode(s) of treatme Note: Some of these would be ap Environment and/or staff change Positive redirection and staff inte Individual and/or group therapy Other Alternatives:	pplicable only in a s		nment. Rehabilitation tr Treatment prog	reatments/therapy (OT, PT, a rams and approaches (habil rintervention techniques		
3. Probable consequences of NO						
Impairment of Work Activities	; □ Fa	amily Relationships	3	☐ Social Functioning		
Possible increase in symptoms le Use of seclusion or restraint Limits on access to possessions Limits on personal freedoms		al		ation and leisure activities aw enforcement authorities self or others		
☐ Limit participation in treatment ar Other Consequences:	nd activities					

4. Possible side effects, warnings, and cautions associated with this medication are listed below. This is not an all-inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text, such as the PDR. As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects in order to enhance care and treatment.

Continued – Possible side effects, warnings, and cautions associated with this medication.

Most Common Side Effects: headache, sleepiness, dizziness, drowsiness the next day, diarrhea, grogginess or feeling as if you have been drugged. fatigue.

Caution: Serious side effects: getting out of bed while not being fully awake anddoing an activity that you do not know you aredoing; abnormal thoughts and behavior. Symptoms include more outgoing or aggressive behavior thannormal, confusion, agitation, hallucinations, wors-ening of depression, suicidal thoughts or actions.; memory loss, anxiety, severe allergic reactions. Symptoms include swelling of the tongue or throat, trouble breathing.

Warning:

- Need to evaluate for co-morbid diagnoses: Revaluate if insomnia persists after 7 to 10 days of use.
- Severe anaphylactic/anaphylactoid reactions: Angioedema and anaphylaxis have been reported. Do not rechallenge if such reactions occur.
- Abnormal thinking, behavioral changes, complex behaviors: May include "sleep-driving" and hallucinations. Immediately evaluate any new onset behavioral changes.
- Depression: Worsening of depression or, suicidal thinking may occur. Prescribe the least amount feasible to avoid intentional overdose.
- Withdrawal effects: Symptoms may occur with rapid dose reduction or discontinuation.
- CNS depressant effects: Use can impair alertness and motor coordination. If used in combination with other CNS depressants, dose reductions may be needed due to additive effects. Do not use with alcohol.
- Elderly/debilitated patients: Use lower dose due to impaired motor, cognitive performance and increased sensitivity.
- Patients with hepatic impairment, mild to moderate COPD, impaired drug metabolism or hemodynamic responses, mild to
 moderate sleep apnea: Use with caution and monitor closely.

2

See PDR for an all-inclusive list of side effects.	

CI	ient Initial	 Date _	

By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

- 1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather, it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
- 2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
- 3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager, or psychologist.
- 4. I have the right to request a review at any time of my record, pursuant to § 51.30(4)(d) or § 51.30(5)(b).
- 5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager, or agency/facility client rights specialist may be contacted for assistance.
- 6. My consent permits the dose to be changed within the anticipated dosage range without signing another consent.
- 7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s), and the probable consequences that may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate, and complete.

 the information to be specific, accurate, and complete. This medication consent is for a period effective immediately and not to exceed need for and continued use of this medication will be reviewed at least quarter client, will be to arrive at and maintain the client at the minimum effective dose 	ly by the Interdisciplinary Tea				
SIGNATURES		DATE	SIGNED		
Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client				
Staff Present at Oral Discussion	Title				
Client / Parent of Minor / Guardian (POA-HC) Comments					
As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.					
Verbal Consent					
Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received ☐ Yes ☐ No			
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received			

3