INFORMED CONSENT FOR MEDICATION

Dosage and / or Side Effect information last revised on 05/20/2020

Completion of this form is voluntary. If informed consent is not given, the medication cannot be administered without a court order unless in an emergency. cinteined in the client's record and is accessible to authorized

I his consent is maintained in the	ellents record ar	nd is accessible to aut	inorizea u	sers.			
Name – Patient / Client (Last, First MI)			ID Num			Date of Birth	
, Name – Individual Preparing This Form		Name – Staff Co	Name – Staff Contact		Name / Telephone Number – Institution		
MEDICATION CATEGORY		MEDICATION	MEDICATION		ECOMMENDED DTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE	
Antihistamine (sedative, antianx	ety) Benadryl (diphenhy	dryl 25mg –		25mg – 300ı	ng		
The anticipated dosage range is without your informed and writte Recommended daily total dosag This medication will be administe	n consent. e range of manufa	•	hysician's		C C		
1. Reason for Use of Psychot Include DSM-5 diagnosis or t				if this is 'Off	Label' Use)		
Positive redirection and staff interaction							
3. Probable consequences of	-				_		
Impairment of Uvrk Activ	ties] Family Relationships	S		Social Functioning		
Possible increase in symptom	s leading to pote	ential					
 Use of seclusion or restraint Limits on access to possessions Limits on personal freedoms Limit participation in treatment and activities Other Consequences: 			 Limits on recreation and leisure activities Intervention of law enforcement authorities Risk of harm to self or others 				
Note: These consequence unusual situations, little o					an inpatient setting. It is als ot administered.	o possible that in	

See Page 2

Client Initial

4. Possible side effects, warnings, and cautions associated with this medication are listed below. This is not an all-inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text, such as the PDR. As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects in order to enhance care and treatment.

Continued – Possible side effects, warnings, and cautions associated with this medication.

Most Common Side Effects: dizziness, drowsiness, fatigue; dry mouth, nose, or throat; gastrointestinal upset, stomach pain, or nausea; headache; thickening of mucus.

Less Common Side Effects: change in vision; body aches or pain; clumsiness or unsteadiness; congestion/nasal stuffiness; constipation; difficult or painful urination/urinary retention; hoarseness; increased sweating; joint pain; muscle pains or stiffness; skin tenderness/itching; nightmares; ringing or buzzing in ears; skin rash/ redness; swollen joints; heartburn; tender swollen glands in neck; tremor; unusual excitement; nervousness/ restlessness; insomnia; tremor; irritability; nausea/vomiting.

Rare Side Effects: Although rare, check with you physician as soon as possible if any of the following side effects occur: abdominal or stomach pain; burning; chills; clay-colored stools or dark urine; cough; difficulty swallowing or breathing, shortness of breath; tightness in chest or wheezing; dizziness; fast or irregular heartbeat; fever; headache; itching; prickly sensations; puffiness or swelling of the eyelids or around the eyes, face, lips or tongue; seizures; unusual tiredness or weakness.

Check with your doctor as soon as possible if any of the following rare side effects occur: Sore throat; unusual bleeding or bruising; unusual tiredness or weakness.

Caution

- Antihistamines will add to the effects of alcohol and other CNS depressants (medicines that slow down the nervous system, possibly causing drowsiness). Some examples of CNS depressants are sedatives, tranquilizers, or sleeping medicine; prescription pain medicine or narcotics; barbiturates; medicine for seizures; muscle relaxants; or anesthetics, including some dental anesthetics. Check with your doctor before taking any of the above while you are using this medicine.
- This medicine may cause some people to become drowsy or less alert than they are normally. Even if taken at bedtime, it may cause some people to feel drowsy or less alert on arising. Some antihistamines are more likely to cause drowsiness than others.
- Make sure you know how you react to the antihistamine you are taking before you drive, use machines, or do anything else that could be dangerous if you are not alert.

See standard reference text for an all-inclusive list of side effects.

F-24277

By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

- 1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather, it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
- 2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
- 3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager, or psychologist.
- 4. I have the right to request a review at any time of my record, pursuant to § 51.30(4)(d) or § 51.30(5)(b).
- 5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager, or agency/facility client rights specialist may be contacted for assistance.
- My consent permits the dose to be changed within the **anticipated dosage range** without signing another consent. 6.
- 7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s), and the probable consequences that may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate, and complete.
- 8. This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The need for and continued use of this medication will be reviewed at least quarterly by the Interdisciplinary Team. The goal, on behalf of the client, will be to arrive at and maintain the client at the minimum effective dose.

SIGNATURES		DATE SIGNED
Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client	
	Parent Guardian (POA-HC)	
Staff Present at Oral Discussion	Title	

Client / Parent of Minor / Guardian (POA-HC) Comments

As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.

Verbal Consent						
Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received				
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received				