

INFORMED CONSENT FOR MEDICATION

Dosage and / or Side Effect information last revised on 12/17/2010

Completion of this form is voluntary. If not completed, the medication cannot be administered without a court order unless in an emergency.
 This consent is maintained in the client's record and is accessible to authorized users.

Name – Patient / Client (Last, First, MI)		ID Number	Living Unit	Birthdate
Name – Individual Preparing This Form		Name – Staff Contact		Name / Telephone Number – Institution

MEDICATION CATEGORY	MEDICATION	RECOMMENDED DAILY TOTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE
Beta-Adrenergic Blocker	Inderal (propranolol)	20mg –320mg	

The anticipated dosage range is to be individualized, may be above or below the recommended range but no medication will be administered without your informed and written consent.

Recommended daily total dosage range of manufacturer, as stated in *Physician's Desk Reference* (PDR) or another standard reference.

This medication will be administered Orally Injection Other – Specify:

1. Reason for Use of Psychotropic Medication and Benefits Expected (note if this is 'Off Label' Use)

Include DSM IV diagnosis or the diagnostic "working hypothesis."

2. Alternative mode(s) of treatment other than or in addition to medications include

Note: Some of these would be applicable only in an inpatient environment.

- | | |
|--|--|
| <input type="checkbox"/> -Environment and / or staff changes | <input type="checkbox"/> -Rehabilitation treatments / therapy (OT, PT, AT) |
| <input type="checkbox"/> -Positive redirection and staff interaction | <input type="checkbox"/> -Treatment programs and approaches (habilitation) |
| <input type="checkbox"/> -Individual and / or group therapy | <input type="checkbox"/> -Use of behavior intervention techniques |

Other Alternatives:

3. Probable consequences of NOT receiving the proposed medication are

Impairment of -Work Activities -Family Relationships -Social Functioning

Possible increase in symptoms leading to potential

- | | |
|---|---|
| <input type="checkbox"/> -Use of seclusion or restraints | <input type="checkbox"/> -Limits on recreation and leisure activities |
| <input type="checkbox"/> -Limits on access to possessions | <input type="checkbox"/> -Intervention of law enforcement authorities |
| <input type="checkbox"/> -Limits on personal freedoms | <input type="checkbox"/> -Risk of harm to self or others |
| <input type="checkbox"/> -Limit participation in treatment and activities | |

Other consequences

Note: These consequences may vary, depending upon whether or not the individual is in an inpatient setting. It is also possible that in unusual situations, little or no adverse consequences may occur if the medications are not administered.

4. Possible side effects, warnings and cautions associated with this medication are listed below. This is not an all inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text such as the PDR or the United States Pharmacopoeia Dispensing Information (USPDI). As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects, in order to enhance care and treatment.

Continued – Possible side effects, warnings and cautions associated with this medication.

The most common side effects include dizziness or lightheadedness, unusually slow pulse, diarrhea, drowsiness, nausea, unusual tiredness or weakness and numbness or tingling of the fingers and / or toes.

Less common side effects include breathing difficulty or wheezing; mental confusion, swelling of ankles, feet and lower legs; anxiety and / or nervousness; constipation, headache, nightmares, trouble sleeping, seeing, hearing, or sensing things not there, feeling of sadness and other symptoms of depression.

Serious skin reactions can occur during treatment with this medicine. Check with your doctor right away if you have any of the following symptoms while taking this medicine: blistering, peeling, or loosening of the skin; chills; cough; diarrhea; fever; itching; joint or muscle pain; red skin lesions; sore throat; sores, ulcers, or white spots in your mouth or lips; or unusual tiredness or weakness.

CAUTION – These medications should be used cautiously with individuals who have diabetes, asthma, or narrow angle glaucoma.

WARNINGS

Angina Pectoris: There have been reports of exacerbation of angina and, in some cases, myocardial infarction, following abrupt discontinuance of propranolol therapy. Therefore, when discontinuance of propranolol is planned, the dosage should be gradually reduced over at least a few weeks, and the patient should be cautioned against interruption or cessation of therapy without the physician's advice. If propranolol therapy is interrupted and exacerbation of angina occurs, it usually is advisable to reinstitute propranolol therapy and take other measures appropriate for the management of unstable angina pectoris. Since coronary artery disease may be unrecognized, it may be prudent to follow the above advice in patients considered at risk of having occult atherosclerotic heart disease who are given propranolol for other indications.

Hypersensitivity and Skin Reactions: Hypersensitivity reactions, including anaphylactic/anaphylactoid reactions, have been associated with the administration of propranolol. Cutaneous reactions, including Stevens-Johnson Syndrome, toxic epidermal necrolysis, exfoliative dermatitis, erythema multiforme, and urticaria, have been reported with use of propranolol.

Cardiac Failure: Sympathetic stimulation may be a vital component supporting circulatory function in patients with congestive heart failure, and its inhibition by beta blockade may precipitate more severe failure. Although beta blockers should be avoided in overt congestive heart failure, some have been shown to be highly beneficial when used with close follow-up in patients with a history of failure who are well compensated and are receiving diuretics as needed. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle. In Patients without a History of Heart Failure, continued use of beta blockers can, in some cases, lead to cardiac failure.

Nonallergic Bronchospasm (e.g., Chronic Bronchitis, Emphysema): In general, patients with bronchospastic lung disease should not receive beta-blockers. Propranolol should be administered with caution in this setting since it may provoke a bronchial asthmatic attack by blocking bronchodilation produced by endogenous and exogenous catecholamine stimulation of beta-receptors.

Major Surgery: The necessity or desirability of withdrawal of beta-blocking therapy prior to major surgery is controversial. It should be noted, however, that the impaired ability of the heart to respond to reflex adrenergic stimuli in propranolol-treated patients may augment the risks of general anesthesia and surgical procedures. Propranolol is a competitive inhibitor of beta-receptor agonists and its effects can be reversed by administration of such agents, e.g., dobutamine or isoproterenol. However, such patients may be subject to protracted severe hypotension.

Diabetes and Hypoglycemia: Beta-adrenergic blockade may prevent the appearance of certain premonitory signs and symptoms (pulse rate and pressure changes) of acute hypoglycemia, especially in labile insulin-dependent diabetics. In these patients, it may be more difficult to adjust the dosage of insulin. Propranolol therapy, particularly when given to infants and children, diabetic or not, has been associated with hypoglycemia especially during fasting as in preparation for surgery. Hypoglycemia has been reported in patients taking propranolol after prolonged physical exertion and in patients with renal insufficiency.

Thyrotoxicosis: Beta-adrenergic blockade may mask certain clinical signs of hyperthyroidism. Therefore, abrupt withdrawal of propranolol may be followed by an exacerbation of symptoms of hyperthyroidism, including thyroid storm. Propranolol may change thyroid-function tests, increasing T4 and reverse T3, and decreasing T3.

Wolff-Parkinson-White Syndrome: Beta-adrenergic blockade in patients with Wolff-Parkinson-White syndrome and tachycardia has been associated with severe bradycardia requiring treatment with a pacemaker. In one case, this result was reported after an initial dose of 5mg propranolol.

See PDR, USPDI or US Hospital Formulary Service for all-inclusive list of side effects.

Client Initial _____ Date _____

Medication : Inderal - (propranolol)

By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager or psychologist.
4. I have the right to request a review at any time of my record, pursuant to ss. 51.30(4)(d) or 51.30(5)(b).
5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager or agency / facility client rights specialist may be contacted for assistance.
6. My consent permits the dose to be changed within the **anticipated dosage range** without signing another consent.
7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s) and the probable consequences, which may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate and complete.
8. This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The need for and continued use of this medication will be reviewed at least quarterly by the Interdisciplinary Team. The goal, on behalf of the client, will be to arrive at and maintain the client at the minimum effective dose.

SIGNATURES

DATE SIGNED

Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client <input type="checkbox"/> Self <input type="checkbox"/> Parent <input type="checkbox"/> Guardian (POA-HC)	
Staff Present at Oral Discussion	Title	

Client / Parent of Minor / Guardian (POA-HC) Comments

As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.

Verbal Consent

Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received

Client Initial _____ Date _____