

INFORMED CONSENT FOR MEDICATION

Dosage and / or Side Effect information last revised on 06/08/2017

Completion of this form is voluntary. If informed consent is not given, the medication cannot be administered without a court order unless in an emergency.

This consent is maintained in the client's record and is accessible to authorized users.

Name – Patient / Client (Last, First MI)	ID Number	Living Unit	Date of Birth
Name – Individual Preparing This Form	Name – Staff Contact	Name / Telephone Number – Institution	

MEDICATION CATEGORY	MEDICATION	RECOMMENDED DAILY TOTAL DOSAGE RANGE	ANTICIPATED DOSAGE RANGE
MAO Inhibitor	Parnate (tranylcypromine)	2.5mg– 30mg	

The anticipated dosage range is to be individualized, may be above or below the recommended range but no medication will be administered without your informed and written consent.

Recommended daily total dosage range of manufacturer, as stated in *Physician's Desk Reference* (PDR) or another standard reference.

This medication will be administered ☐ Orally ☐ Injection ☐ Other – Specify:

1. Reason for Use of Psychotropic Medication and Benefits Expected (note if this is 'Off-Label' Use)

Include DSM-5 diagnosis or the diagnostic impression ("working hypothesis.")

2. Alternative mode(s) of treatment other than OR in addition to medications include

Note: Some of these would be applicable only in an inpatient environment.

- | | |
|---|---|
| <input type="checkbox"/> Environment and/or staff changes | <input type="checkbox"/> Rehabilitation treatments/therapy (OT, PT, AT) |
| <input type="checkbox"/> Positive redirection and staff interaction | <input type="checkbox"/> Treatment programs and approaches (habilitation) |
| <input type="checkbox"/> Individual and/or group therapy | <input type="checkbox"/> Use of behavior intervention techniques |

Other Alternatives:

3. Probable consequences of NOT receiving the proposed medication are

Impairment of ☐ Work Activities ☐ Family Relationships ☐ Social Functioning

Possible increase in symptoms leading to potential

- | | |
|--|--|
| <input type="checkbox"/> Use of seclusion or restraint | <input type="checkbox"/> Limits on recreation and leisure activities |
| <input type="checkbox"/> Limits on access to possessions | <input type="checkbox"/> Intervention of law enforcement authorities |
| <input type="checkbox"/> Limits on personal freedoms | <input type="checkbox"/> Risk of harm to self or others |
| <input type="checkbox"/> Limit participation in treatment and activities | |

Other Consequences:

Note: These consequences may vary depending upon whether or not the individual is in an inpatient setting. It is also possible that in unusual situations, little or no adverse consequences may occur if the medications are not administered.

See Page 2

Client Initial _____ Date _____

-
4. Possible side effects, warnings, and cautions associated with this medication are listed below. This is not an all-inclusive list but is representative of items of potential clinical significance to you. For more information on this medication, you may consult further with your physician or refer to a standard text, such as the PDR. As part of monitoring some of these potential side effects, your physician may order laboratory or other tests. The treatment team will closely monitor individuals who are unable to readily communicate side effects in order to enhance care and treatment.
-

Continued – Possible side effects, warnings, and cautions associated with this medication.

Most Common Side Effects

Check with your doctor as soon as possible if any of the following more common side effects occur: Dizziness or lightheadedness (severe), especially when getting up from a lying or sitting position.

Other more common side effects include: Blurred vision; decreased amount of urine; decreased sexual ability; dizziness or lightheadedness (mild), especially when getting up from a lying or sitting position; drowsiness; headache (mild); increased appetite (especially for sweets) or weight gain; increased sweating; muscle twitching during sleep; nausea; restlessness; shakiness or trembling; tiredness and weakness; trouble in sleeping.

Less Common Side Effects

Check with your doctor as soon as possible if any of the following less common side effects occur: Diarrhea; fast or pounding heartbeat; swelling of feet or lower legs; unusual excitement or nervousness.

Other less common side effects include: Blurred vision; decreased amount of urine; decreased sexual ability; dizziness or lightheadedness (mild), especially when getting up from a lying or sitting position; drowsiness; headache (mild); increased appetite (especially for sweets) or weight gain; increased sweating; muscle twitching during sleep; nausea; restlessness; shakiness or trembling; tiredness and weakness; trouble in sleeping.

Rare Side Effects

Check with your doctor as soon as possible if any of the following rare side effects occur: Dark urine; fever; skin rash; slurred speech; sore throat; staggering walk; yellow eyes or skin.

Other rare side effects include: Chills; constipation; decreased appetite; dryness of mouth.

Caution

Stop taking this medicine and get emergency help immediately if any of the following effects occur:

Symptoms of unusually high blood pressure (hypertensive crisis); chest pain (severe); enlarged pupils; fast or slow heartbeat; headache (severe); increased sensitivity of eyes to light; increased sweating (possibly with fever or cold, clammy skin); nausea and vomiting; stiff or sore neck.

Warning

Hypertensive Crisis

The most important reaction associated with tranylcypromine is the occurrence of hypertensive crises which have sometimes been fatal. These crises are characterized by some or all of the following symptoms: occipital headache which may radiate frontally, palpitation, neck stiffness or soreness, nausea or vomiting, sweating (sometimes with fever and sometimes with cold, clammy skin), and photophobia. Either tachycardia or bradycardia may be present, and associated constricting chest pain and dilated pupils may occur. Intracranial bleeding, sometimes fatal in outcome, has been reported in association with the paradoxical increase in blood pressure.

In all patients taking tranylcypromine, blood pressure should be followed closely to detect evidence of any pressor response. It is emphasized that full reliance should not be placed on blood pressure readings, but that the patient should also be observed frequently. Therapy should be discontinued immediately upon the occurrence of palpitation or frequent headaches during therapy with tranylcypromine. These signs may be prodromal of a hypertensive crisis.

See standard reference text for an all-inclusive list of side effects.

By my signature below, I GIVE consent for the named medication on Page 1 and anticipated dosage range. My signature also indicates that I understand the following:

1. I can refuse to give consent or can withdraw my consent at any time with written notification to the institution director or designee. This will not affect my right to change my decision at a later date. If I withdraw consent after a medication is started, I realize that the medication may not be discontinued immediately. Rather, it will be tapered as rapidly as medically safe and then discontinued so as to prevent an adverse medical consequence, such as seizures, due to rapid medication withdrawal.
2. Questions regarding this medication can be discussed with the Interdisciplinary Team, including the physician. The staff contact person can assist in making any necessary arrangements.
3. Questions regarding any behavior support plan or behavior intervention plan, which correspond with the use of the medication, can be directed to the client's social worker, case manager, or psychologist.
4. I have the right to request a review at any time of my record, pursuant to § 51.30(4)(d) or § 51.30(5)(b).
5. I have a legal right to file a complaint if I feel that client rights have been inappropriately restricted. The client's social worker, case manager, or agency/facility client rights specialist may be contacted for assistance.
6. My consent permits the dose to be changed within the **anticipated dosage range** without signing another consent.
7. I understand the reasons for the use of the medication, its potential risks and benefits, other alternative treatment(s), and the probable consequences that may occur if the proposed medication is not given. I have been given adequate time to study the information and find the information to be specific, accurate, and complete.
8. This medication consent is for a period effective immediately and not to exceed fifteen (15) months from the date of my signature. The need for and continued use of this medication will be reviewed at least quarterly by the Interdisciplinary Team. The goal, on behalf of the client, will be to arrive at and maintain the client at the minimum effective dose.

SIGNATURES**DATE SIGNED**

Client – If Presumed Competent to Consent/Parent of Minor/Guardian (POA-HC)	Relationship to Client <input type="checkbox"/> Self <input type="checkbox"/> Parent <input type="checkbox"/> Guardian (POA-HC)	
Staff Present at Oral Discussion	Title	
Client / Parent of Minor / Guardian (POA-HC) Comments		

As parent/guardian (POA-HC) was not available for signature, he/she was verbally informed of the information in this consent.

Verbal Consent

Obtained by – PRINT – Staff Name	Date Obtained	Written Consent Received <input type="checkbox"/> Yes <input type="checkbox"/> No
Obtained from – PRINT – Parent / Guardian (POA-HC) Name	Date Expires	Date Received