



DIVISION OF PUBLIC HEALTH

1 WEST WILSON STREET  
P O BOX 2659  
MADISON WI 53701-2659

Jim Doyle  
Governor

Kevin R. Hayden  
Secretary

State of Wisconsin

Department of Health and Family Services

608-266-1251  
FAX: 608-267-2832  
TTY: 888-701-1253  
dhfs.wisconsin.gov

Date: March 14, 2008

To: Local Health Departments

From: Jeffrey P. Davis, MD, Chief Medical Officer and  
State Epidemiologist for Communicable Diseases  
*Jeffrey P. Davis MD*

Jonathan L. Temte, MD, PhD, Chair  
Wisconsin Council on Immunization Practices  
*Jonathan L. Temte MD PhD*

Re: Flood water exposure and implications for vaccinating adults

According to the Centers for Disease Control and Prevention (CDC), outbreaks of infectious diseases after a flood in the U.S. are unusual and mass immunization programs are not routinely recommended. Exposure to flood water alone is not a reason to give tetanus-containing vaccine or any other vaccine.

**Tetanus Vaccine:**

The need for tetanus vaccination after exposure to flood water depends on the presence of a wound, the condition of the wound and the likelihood that the injured person is susceptible to tetanus. The recommendations for use of tetanus-containing vaccine that apply during a flood are the same as those that apply at any other time. Tdap is preferred to Td for adults who have never received Tdap because immunity against pertussis wanes. Use a Tdap formulation which has been licensed for adults. Td should be used for adults previously vaccinated with Tdap and for persons  $\geq 65$  years of age.

- Persons with severe wounds should be evaluated by a medical provider. Those who sustain puncture wounds or wounds that may be contaminated with soil or fecal material should receive tetanus-containing vaccine if more than 5 years have elapsed since their last tetanus booster. Persons who have not completed a primary tetanus series (a minimum of 3 doses of tetanus-and diphtheria-containing vaccine) must be vaccinated and given tetanus immune globulin (TIG) as soon as possible.
- Persons who sustain clean, minor wounds should receive tetanus-containing vaccine if they have not completed a primary tetanus series or if more than 10 years have elapsed since their last tetanus booster.
- All persons should receive a tetanus booster every 10 years.

**Hepatitis A vaccine:**

There is no specific reason to give hepatitis A vaccine routinely during floods. No transmission of hepatitis A virus from contaminated water has been identified in the U.S. since the 1980s. Hepatitis A vaccine is not routinely recommended for sewage workers.

**Hepatitis B vaccine:**

Hepatitis B vaccine is recommended for people who will be providing direct patient care or who are otherwise expected to have routine contact with other people's bodily fluids in the course of their work activities. There is no special need for hepatitis B vaccination during a flood.

CC: Lori Getter, Wisconsin Emergency Management