Wisconsin Public Psychiatry Network Teleconference (WPPNT)

- This teleconference is brought to you by the Wisconsin Department of Health Services (DHS) Bureau of Prevention, Treatment, and Recovery and the University of Wisconsin-Madison, Department of Psychiatry.
- The Department of Health Services makes no representations or warranty as to the accuracy, reliability, timeliness, quality, suitability or completeness of or results of the materials in this presentation. Use of information contained in this presentation may require express authority from a third party.

1

The Columbia Suicide Severity Rating Scale (C-SSRS): Improved Risk Assessment and the Positive Impact on Suicide Prevention

Jeffrey Garbelman, Ph.D.

Suicide prevention efforts depend upon appropriate identification & screening

CDC Self-Directed Violence: Uniform Definitions Adopted Columbia Definitions

(link to C-SSRS in CDC document)



Source: Posner K, Oquendo MA, Gould M, Stanley B, Davies M. Columbia Classification Algorithm of Suicide Assessment (C-CASA): Classification of Suicidal Events in the FDA's Pediatric Suicidal Risk Analysis of Antidepressants. Am J Psychiatry. 2007; 164:1035-1043. http://cssrs.columbia.edu/

SELF-DIRECTED VIOLENCE SURVEILLANCE: UNIFORM DEFINITIONS AND RECOMMENDED DATA ELEMENTS 21

How to Fix the Problem…
Columbia - Suicide Severity Rating Scale

Posner, K.; Brent, D.; Lucas, C.; Gould, M.; Stanley, B.; Brown, G.; Zelazny, J.; Fisher, P.; Burke, A.; Oquendo, M.; Mann, J.

- Developed by leading experts (collaboration with Beck's group) for National Adolescent Attempter's Study in response to need for a measure to assess *both* behavior and ideation
- Evidence-based and supported
- Feasible, low-burden – short administration time (average is a few minutes)

includes only the most essential, evidence-based items needed in a thorough assessment

Severity of Ideation Subscale

Simply....

- 1-5 rating for suicidal ideation, of increasing severity (from a wish to die to an active thought of killing oneself with plan and intent)

Two Screen Questions for Ideation

- *Have you wished you were dead or wished you could go to sleep and not wake up?*
- *Have you actually had any thoughts of killing yourself?*

If answer is "No" to both, no more questions on ideation

- Relevant behaviors assessed in one additional question
- All items include definitions for each term and standardized questions for each category are included to guide the interviewer for facilitating improved identification

Suicidal Ideation

1. Wish to die

– *Have you wished you were dead or wished you could go to sleep and not wake up?*

2. Active Thoughts of Killing Oneself

– *Have you actually had any thoughts of killing yourself?*

**** If "NO" to both these questions Suicidal Ideation Section is finished.****
**** If "YES" to 'Active thoughts' ask the following three questions.****

3. Associated Thoughts of Methods

– *Have you been thinking about how you might do this?*

4. Some Intent

– *Have you had these thoughts and had some intention of acting on them?*

5. Plan and Intent

– *Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?*

Auditory hallucinations qualify as ideation

15



Transition Data

- Cumulative Probabilities of transition:
- Ideation to Plan 34 %
- **Plan to Attempt 72 %**

Fowler, 2012;

SSI Total Score by Highest Level of Ideation on the C-SSRS



American Foundation *for* Suicide Prevention

$F(5,185) = 14.35$, $p < .001$

Currier, Brown & Stanley (2009)

58

Clinical Monitoring Guidance: Threshold for Next Steps

SUICIDAL IDEATION	
Ask questions 1 and 2. If both are negative, proceed to "Suicidal Behavior" section. If the answer to question 2 is "yes," ask questions 3, 4 and 5. If the answer to question 1 and/or 2 is "yes", complete "Intensity of Ideation" section below.	Lifetime: Time He/She Felt Most Suicidal
1. Wish to be Dead Subject endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up. ***Have you wished you were dead or wished you could go to sleep and not wake up?*** If yes, describe:	Yes ☐ No ☐
2. Non-Specific Active Suicidal Thoughts General, non-specific thoughts of wanting to end one's life/commit suicide (e.g. *"I've thought about killing myself"*) without thoughts of ways to kill oneself/associated methods, intent, or plan. ***Have you actually had any thoughts of killing yourself?*** If yes, describe:	Yes ☐ No ☐
3. Active Suicidal Ideation with Any Methods (Not Plan) without Intent to Act Subject endorses thoughts of suicide and has thought of at least one method during the assessment period. This is different than a specific plan with time, place or method details worked out (e.g. thought of method to kill self but not a specific plan). Includes person who would say, *"I thought about taking an overdose but I never made a specific plan as to when, where or how I would actually do it ..... and I would never go through with it".* ***Have you been thinking about how you might do this?*** If yes, describe:	Yes ☐ No ☐
4. Active Suicidal Ideation with Some Intent to Act, without Specific Plan Active suicidal thoughts of killing oneself and subject reports having some intent to act on such thoughts, as opposed to *"I have the thoughts but I definitely will not do anything about them".* ***Have you had these thoughts and had some intention of acting on them?*** If yes, describe:	Yes ☐ No ☐
5. Active Suicidal Ideation with Specific Plan and Intent Thoughts of killing oneself with details of plan fully or partially worked out and subject has some intent to carry it out. ***Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?*** If yes, describe:	Yes ☐ No ☐

Indicates Need for Next Step

Data Confirmation… 4 and 5 Predicts Attempts in National Attempter Study

(Posner et al., *AJP* December 2011)

- C-SSRS Lifetime Ideation at baseline, types 4 and 5, predicted suicide attempts in adolescent suicide attempters, followed over a year
  - OR = 3.26, 95% CI: 1.02-10.45, p = 0.047
  - **Beck SSI NOT predictive**
- C-SSRS Lifetime Ideation, types 4 and 5, predicted actual, interrupted or aborted attempts on CSHF
  - OR = 2.76, 95% CI: 1.07-7.12, p = 0.036

Advantages….Operationalized Criteria for Next Steps

- Allows for setting parameters for triggering next steps whatever they may be
  - e.g., 4 or 5 on ideation item to indicate need for immediate referral
  - Decreases unnecessary referrals, interventions, exclusions, etc.

In the past, people didn't know what to manage, so they would hear **any answer and intervene…*

62

Thresholds facilitate identification of those at highest, triage, and care delivery

4/5 → Psych consult

3 → Consult to Care team

Example: Streamlining Care in Hospital Policies

PROCEDURE:

Question	Trigger
Level 4/5 Yes to question 4 or 5	• Nursing Order to call MD for Psych Consult • Nursing Interventions (print on Kardex): • Pt Safety Monitor – 1:1 Observation • Pt Safety Monitor – Within arm's reach at all times • Complete Self Harm Safety Assessment every shift • Affix Suicide Risk Magnet to door • Revise Diet order to Safe tray • Alerts to ATC, Nutrition Services, Environmental Services and Security • Progress note for chart
Level 3 Yes to question 3 (and no to question 4 and 5)	• Consult to Care Team • Nursing Interventions (prints on kardex): • Pt Safety Monitor – 1:1 Observation • Pt Safety Monitor – Within arm's reach at all times • Complete Self Harm Safety Assessment every shift • Affix Suicide Risk Magnet to door • Revise Diet order to Safe Tray • Alerts to ATC, Nutrition Services, Environmental Services, Spruce Facilitator and Security • Progress note for chart

(Reading Hospital Policy)

65

Suicidal Behavior Subscale

Suicidal Behavior section and the C-SSRS screening version

Same Screening Questions

***If 1 and 2 are no, ideation section is done**

This is the C-SSRS Screener

***Minimum of 3 Questions**

COLUMBIA-SUICIDE SEVERITY RATING SCALE
Posner, Brent, Lucas, Gould, Stanley, Brown, Fisher, Zelazny, Burke, Oquendo, & Mann
Screen Version

SUICIDE IDEATION DEFINITIONS AND PROMPTS:	Past month	
Ask questions that are in bolded and underlined. The rest of the information at each question is for staff information only.	Yes	NO
Ask Questions 1 and 2		
1) Wish to be Dead: Person endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up? *Have you wished you were dead or wished you could go to sleep and not wake up?*		
2) Suicidal Thoughts: General non-specific thoughts of wanting to end one's life/commit suicide, *"I've thought about killing myself"* without general thoughts of ways to kill oneself/associated methods, intent, or plan." *Have you actually had any thoughts of killing yourself?*		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.		
3) Suicidal Thoughts with Method (without Specific Plan or Intent to Act): Person endorses thoughts of suicide and has thought of a least one method during the assessment period. This is different than a specific plan with time, place or method details worked out. *"I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it...and I would never go through with it."* *Have you been thinking about how you might kill yourself?*		
4) Suicidal Intent (without Specific Plan): Active suicidal thoughts of killing oneself and patient reports having some intent to act on such thoughts, as opposed to *"I have the thoughts but I definitely will not do anything about them."* *Have you had these thoughts and had some intention of acting on them?*		
5) Suicide Intent with Specific Plan: Thoughts of killing oneself with details of plan fully or partially worked out and person has some intent to carry it out. *Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?*		
6) Suicide Behavior Question *"Have you ever done anything, started to do anything, or prepared to do anything to end your life?"* Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc. **If YES, ask:** *How long ago did you do any of these?* ☐Over a year ago? ☐Between three months and a year ago? ☐Within the last three months?		

6) Suicide Behavior Question
"Have you ever done anything, started to do anything, or prepared to do anything to end your life?"

Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.

If YES, ask: How long ago did you do any of these?

- Over a year ago?
- Between three months and a year ago?
- Within the last three months?

Other Suicidal Behaviors....
Interrupted Attempt

- When person starts to take steps to end their life but someone or something stops them
  - Bottle of pills or gun in hand but someone grabs it
  - On ledge poised to jump
- Question:
  - *Has there been a time when you started to do something to end your life but someone or something stopped you before you actually did anything?*

Aborted Attempt

- When person begins to take steps towards making a suicide attempt, ***but stops themselves*** before they actually have engaged in any self-destructive behavior
- Example:
  - Man plans to drive his car off the road at high speed at a chosen destination. On the way to the destination, he changes his mind and returns home
  - Man walks up to the roof to jump, but changes his mind and turns around
  - She has gun in her hand, but then puts it down
- Question:
  - *Has there been a time when you started to do something to end your life but you stopped yourself before you actually did anything?*

41

Preparatory Acts or Behavior

- Definition:
  - Any other behavior (beyond saying something) with suicidal intent
- Examples
  - Collecting or buying pills
  - Purchasing a gun
  - Writing a will or a suicide note
- Question:
  - *Have you taken any steps towards making a suicide attempt or preparing to kill yourself (such as, collecting pills, getting a gun, giving valuables away, writing a suicide note)?*

42

Research Supported Items/Criteria

- **Preparatory Behavior**
  - Those with recent preparatory behavior (e.g., collecting pills, razors, or loaded weapon) **8x** more likely to commit suicide (Brown & Beck, unpublished)
  - Those who reported at worst point **10x** more likely to commit suicide (Brown & Beck, unpublished)
  - Preparatory Behavior on SSI predicted completed suicide and history of suicide attempts (Joiner et al., 2003)
- **Interrupted Suicide Attempts**
  - **3x** more likely to commit suicide (Steer, Beck & Lester, 1988)
  - No difference in levels of intent
- **Aborted Suicide Attempts**
  - 29-50% of samples report aborted attempts (Marzuk et al., 1997; Barber et al., 1998)
    - Evidencing this as common suicidal behavior
  - Subjects who made aborted attempts **2x** as likely to have made a suicide attempt (Barber et al., 1998)
  - Intent scores similar with suicide attempters

32

eC-SSRS...Depressed Subjects... *ALL* Behaviors Are Prevalent and Predictive

n = 28,699 administrations

Each behavior is EQUALLY PREDICTIVE to an attempt

No Behavior: 28,303
Actual Attempt: 70
Interrupted Attempt: 178
Aborted Attempt: 223
Preparatory Behavior: 71



Multiple behaviors = greater risk

***Only 1.7% had any worrisome answer**
***Only .9% with ~50,000 administrations**

472 Interrupted, Aborted and Preparatory (87%)
vs. 70 Actual Attempts (13%) *Mundt et al., 2011* 10

Suicidal Behavior Section and non – screening versions of C-SSRS

*Lifetime/Recent C-SSRS Version
* Since Last Visit Clinical Version

SUICIDAL BEHAVIOR *(Check all that apply, so long as these are separate events; must ask about all types)*	**Lifetime**	**Recent**
Actual Attempt: A potentially self-injurious act committed with at least some wish to die, *as a result of act*. Behavior was in part thought of as method to kill oneself. Intent does not have to be 100%. If there is ***any*** intent/desire to die associated with the act, then it can be considered an actual suicide attempt. ***There does not have to be any injury or harm***, just the potential for injury or harm. If person pulls trigger while gun is in mouth but gun is broken so no injury results, this is considered an attempt. Inferring Intent: Even if an individual denies intent/wish to die, it may be inferred clinically from the behavior or circumstances. For example, a highly lethal act that is clearly not an accident so no other intent but suicide can be inferred (e.g., gunshot to head, jumping from window of a high floor/story). Also, if someone denies intent to die, but they thought that what they did could be lethal, intent may be inferred. ***Have you made a suicide attempt?*** ***Have you done anything to harm yourself?*** ***Have you done anything dangerous where you could have died?*** ***What did you do?*** ***Did you_____ as a way to end your life?*** ***Did you want to die (even a little) when you____?*** ***Were you trying to end your life when you ____?*** ***Or Did you think it was possible you could have died from____?*** ***Or did you do it purely for other reasons / without ANY intention of killing yourself (like to relieve stress, feel better, get sympathy, or get something else to happen)?*** (Self-Injurious Behavior without suicidal intent) If yes, describe:	Yes No ☐ ☐ Total # of Attempts ____	Yes No ☐ ☐ Total # of Attempts ____
Has subject engaged in Non-Suicidal Self-Injurious Behavior?	Yes No ☐ ☐	Yes No ☐ ☐

Interrupted Attempt: When the person is interrupted (by an outside circumstance) from starting the potentially self-injurious act *(if not for that, actual attempt would have occurred)*. Overdose: Person has pills in hand but is stopped from ingesting. Once they ingest any pills, this becomes an attempt rather than an interrupted attempt. Shooting: Person has gun pointed toward self, gun is taken away by someone else, or is somehow prevented from pulling trigger. Once they pull the trigger, even if the gun fails to fire, it is an attempt. Jumping: Person is poised to jump, is grabbed and taken down from ledge. Hanging: Person has noose around neck but has not yet started to hang - is stopped from doing so. ***Has there been a time when you started to do something to end your life but someone or something stopped you before you actually did anything?*** If yes, describe:	Yes ☐ No ☐ Total # of interrupted ____	Yes ☐ No ☐ Total # of interrupted ____
Aborted or Self-Interrupted Attempt: When person begins to take steps toward making a suicide attempt, but stops themselves before they actually have engaged in any self-destructive behavior. Examples are similar to interrupted attempts, except that the individual stops him/herself, instead of being stopped by something else. ***Has there been a time when you started to do something to try to end your life but you stopped yourself before you actually did anything?*** If yes, describe:	Yes ☐ No ☐ Total # of aborted or self-interrupted ____	Yes ☐ No ☐ Total # of aborted or self-interrupted ____
Preparatory Acts or Behavior: Acts or preparation towards imminently making a suicide attempt. This can include anything beyond a verbalization or thought, such as assembling a specific method (e.g., buying pills, purchasing a gun) or preparing for one's death by suicide (e.g., giving things away, writing a suicide note). ***Have you taken any steps towards making a suicide attempt or preparing to kill yourself (such as collecting pills, getting a gun, giving valuables away or writing a suicide note)?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
Suicidal Behavior: Suicidal behavior was present during the assessment period?	Yes ☐ No ☐	Yes ☐ No ☐

Have you made a suicide attempt?

Have you done anything to harm yourself?

Have you done anything dangerous where you could have died?

What did you do?

Did you_____ as a way to end your life?

Did you want to die (even a little) when you_____?

Were you trying to end your life when you _____?

Or Did you think it was possible you could have died from_____?

Lethality

	Most Recent Attempt Date:	Most Lethal Attempt Date:	Initial/First Attempt Date:
Actual Lethality/Medical Damage: 0. No physical damage or very minor physical damage (e.g., surface scratches). 1. Minor physical damage (e.g., lethargic speech; first-degree burns; mild bleeding; sprains). 2. Moderate physical damage; medical attention needed (e.g., conscious but sleepy, somewhat responsive; second-degree burns; bleeding of major vessel). 3. Moderately severe physical damage; *medical* hospitalization and likely intensive care required (e.g., comatose with reflexes intact; third-degree burns less than 20% of body; extensive blood loss but can recover; major fractures). 4. Severe physical damage; *medical* hospitalization with intensive care required (e.g., comatose without reflexes; third-degree burns over 20% of body; extensive blood loss with unstable vital signs; major damage to a vital area). 5. Death	*Enter Code* ____	*Enter Code* ____	*Enter Code* ____
Potential Lethality: Only Answer if Actual Lethality=0 Likely lethality of actual attempt if no medical damage (the following examples, while having no actual medical damage, had potential for very serious lethality: put gun in mouth and pulled the trigger but gun fails to fire so no medical damage; laying on train tracks with oncoming train but pulled away before run over). 0 = Behavior not likely to result in injury 1 = Behavior likely to result in injury but not likely to cause death 2 = Behavior likely to result in death despite available medical care	*Enter Code* ____	*Enter Code* ____	*Enter Code* ____

Lethality

(Compilation of Beck Medical Lethality Rating Scale)

What actually happened in terms of medical damage?

For example if there was a cut, did it require a Band-Aid or a bandage? Did it bleed a little bit or profusely?

Actual Lethality/Medical Damage:

0. No physical damage or very minor physical damage (e.g. surface scratches).
1. Minor physical damage (e.g. lethargic speech; first-degree burns; mild bleeding; sprains).
2. Moderate physical damage; medical attention needed (e.g. conscious but sleepy, somewhat responsive; second-degree burns; bleeding of major vessel).
3. Moderately severe physical damage; *medical* hospitalization and likely intensive care required (e.g. comatose with reflexes intact; third-degree burns less than 20% of body; extensive blood loss but can recover; major fractures).
4. Severe physical damage; *medical* hospitalization with intensive care required (e.g. comatose without reflexes; third-degree burns over 20% of body; extensive blood loss with unstable vital signs; major damage to a vital area).
5. Death

Why Potential Lethality?

Likely lethality of attempt if no medical damage. Examples of why this is important are cases in which there was no actual medical damage but the potential for very serious lethality

- Laying on tracks with an oncoming train but pulling away before run over
- Put gun in mouth and pulled trigger but it failed to fire

Potential Lethality: Only Answer if Actual Lethality=0

Likely lethality of actual attempt if no medical damage (the following examples, while having no actual medical damage, had potential for very serious lethality: put gun in mouth and pulled the trigger but gun fails to fire so no medical damage; laying on train tracks with oncoming train but pulled away before run over).

0 = Behavior not likely to result in injury
1 = Behavior likely to result in injury but not likely to cause death
2 = Behavior likely to result in death despite available medical care

50

Adding Suicidal Behaviors to Outcome Decisions or 'triage points'- Example

Prediction in Non-Suicidal Adults & Adolescents

- Confirmed By e-CSSRS data: 35,007 (3776 subjects) across depression, epilepsy, insomnia, fibromyalgia
  - Patients with baseline **prior ideation of 4 or 5** or **prior behavior** are 4-5x more likely to report suicidal behavior at follow up than patients with negative baseline report.
  - Patients with **both** are 8x more likely to report suicidal behavior
- Prediction in non-suicidal adolescents – emergency department follow-up study (King et al.)

Centerstone Alert and Monitoring System

****Largest Provider of Behavioral Healthcare in the United States**

Alert and Monitoring System

The Electronic Health Record (EHR) is designed to offer assistance to providers assessing service recipients for high suicide risk. Based on information collected in the applicable Columbia SSRS tool, a service recipient can be identified as being at high risk for suicide. Those who will be considered at high risk for suicide will have a positive endorsement of **either** of the following (research found these to be highly predictive of completed suicides):

a. A positive endorsement, relative to the past 30 days, in the **"Suicidal Thoughts" section of item # 4** (Have you had these thoughts and had some intention of acting on them?) **or item # 5** (Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?).
b. A positive endorsement, relative to the past 90 days, in the **"Suicide Behavior" section of item # 6** (Have you ever done anything, started to do anything, or prepared to do anything to end your life?).

66

Streamlining Identification, Triage & Care Delivery in the Military

Fort Carson

COLUMBIA-SUICIDE SEVERITY RATING SCALE
Screening Version with Triage Patients—3 step form

II. EACH Response Protocol to C-SSRS Screening

Suicide Ideation Level

LEVEL	SEVERITY	MANAGEMENT PROTOCOL
0	Low risk	ROUTINE Behavioral Health Referral at physician discretion
1 & 2	Mild	ROUTINE Behavioral Health Referral at discharge
3	Moderate	Review by Care Team- Consider safety precautions and telephone consult with Behavioral Health
4 & 5	Serious	EMERGENT ACTION NECESSARY: Behavioral Health Consultation and Patient Safety Monitor/ Procedures

Suicide Behavior History

	MANAGEMENT PROTOCOL
1 week ago and less	ACUTE: Behavioral Health Consultation and Patient Safety precautions
Between 1 week and 3 months ago	CONCERN: Care Team Review, safety precautions and telephone consultation with Behavioral Health
Over 3 months ago	DISCRETIONARY: Consider Behavioral Health Referral at discharge

III. REFERENCE ONLY: SUICIDE IDEATION DEFINITIONS AND PROMPTS
Note: Wording may be adjusted for children and young adolescents

1	**Ideation I** Wish to be Dead:	***Have you wished you were dead or wished you could go to sleep and not wake up?*** Person endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up?
2	**Ideation II** Suicidal Thoughts:	***Have you had any actual thoughts of killing yourself?*** General non-specific thoughts of wanting to end one's life/commit suicide. *"I've thought about killing myself" without general thoughts of ways to kill oneself/associated methods, intent, or plan."*
3	**Suicidal Thoughts with Method** *(without Specific Plan or Intent to Act):*	***Have you been thinking about how you might kill yourself?*** Person endorses thoughts of suicide and has thought of a least one method during the assessment period. This is different than a specific plan with time, place or method details worked out. *"I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it…and I would never go through with it."*
4	**Suicidal Intent I** *(without Specific Plan):*	***Have you had these thoughts and had some intention of acting on them?*** Active suicidal thoughts of killing oneself and patient reports having some intent to act on such thoughts, as opposed to *"I have the thoughts but I definitely will not do anything about them."*
5	**Suicide Intent II** *(with Specific Plan)*	***Have you started to work out or worked out the details of how to kill yourself?*** Thoughts of killing oneself with details of plan fully or partially worked out and person has some intent to carry it out.

68

Intensity of Ideation Subscale

Intensity of Ideation Subscale

- Is not a part of the C-SSRS screen version
- Available in the Lifetime/Recent and Since Last Visit Clinical Version
- Can be used as a part of informing clinical judgment
- Or it can be incorporated into formal triage points

SUICIDAL IDEATION *Ask questions 1 and 2. If both are negative, proceed to "Suicidal Behavior" section. If the answer to question 2 is "yes", ask questions 3, 4 and 5. If the answer to question 1 and/or 2 is "yes", complete "Intensity of Ideation" section below.*	Lifetime: Time He/She Felt Most Suicidal	Past 1 month
1. Wish to be Dead Subject endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up. ***Have you wished you were dead or wished you could go to sleep and not wake up?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
2. Non-Specific Active Suicidal Thoughts General non-specific thoughts of wanting to end one's life/commit suicide (e.g., *"I've thought about killing myself"*) without thoughts of ways to kill oneself/associated methods, intent, or plan during the assessment period. ***Have you actually had any thoughts of killing yourself?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
3. Active Suicidal Ideation with Any Methods (Not Plan) without Intent to Act Subject endorses thoughts of suicide and has thought of at least one method during the assessment period. This is different than a specific plan with time, place or method details worked out (e.g., thought of method to kill self but not a specific plan). Includes person who would say, *"I thought about taking an overdose but I never made a specific plan as to when, where or how I would actually do it...and I would never go through with it."* ***Have you been thinking about how you might do this?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
4. Active Suicidal Ideation with Some Intent to Act, without Specific Plan Active suicidal thoughts of killing oneself and subject reports having some intent to act on such thoughts, as opposed to *"I have the thoughts but I definitely will not do anything about them."* ***Have you had these thoughts and had some intention of acting on them?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
5. Active Suicidal Ideation with Specific Plan and Intent Thoughts of killing oneself with details of plan fully or partially worked out and subject has some intent to carry it out. ***Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?*** If yes, describe:	Yes ☐ No ☐	Yes ☐ No ☐
INTENSITY OF IDEATION *The following features should be rated with respect to the most severe type of ideation (i.e., 1-5 from above, with 1 being the least severe and 5 being the most severe). Ask about time he/she was feeling the most suicidal.* Lifetime - ***Most Severe Ideation:*** ___ (Type # (1-5)) ___ (Description of Ideation) Recent - ***Most Severe Ideation:*** ___ (Type # (1-5)) ___ (Description of Ideation)	Most Severe	Most Severe
Frequency ***How many times have you had these thoughts?*** (1) Less than once a week (2) Once a week (3) 2-5 times in week (4) Daily or almost daily (5) Many times each day	____	____
Duration ***When you have the thoughts how long do they last?*** (1) Fleeting - few seconds or minutes (2) Less than 1 hour/some of the time (3) 1-4 hours/a lot of time (4) 4-8 hours/most of day (5) More than 8 hours/persistent or continuous	____	____
Controllability ***Could/can you stop thinking about killing yourself or wanting to die if you want to?*** (1) Easily able to control thoughts (2) Can control thoughts with little difficulty (3) Can control thoughts with some difficulty (4) Can control thoughts with a lot of difficulty (5) Unable to control thoughts (0) Does not attempt to control thoughts	____	____
Deterrents ***Are there things - anyone or anything (e.g., family, religion, pain of death) - that stopped you from wanting to die or acting on thoughts of committing suicide?*** (1) Deterrents definitely stopped you from attempting suicide (2) Deterrents probably stopped you (3) Uncertain that deterrents stopped you (4) Deterrents most likely did not stop you (5) Deterrents definitely did not stop you (0) Does not apply	____	____
Reasons for Ideation ***What sort of reasons did you have for thinking about wanting to die or killing yourself? Was it to end the pain or stop the way you were feeling (in other words you couldn't go on living with this pain or how you were feeling) or was it to get attention, revenge or a reaction from others? Or both?*** (1) Completely to get attention, revenge or a reaction from others (2) Mostly to get attention, revenge or a reaction from others (3) Equally to get attention, revenge or a reaction from others and to end/stop the pain (4) Mostly to end or stop the pain (you couldn't go on living with the pain or how you were feeling) (5) Completely to end or stop the pain (you couldn't go on living with the pain or how you were feeling) (0) Does not apply	____	____

Intensity of Ideation

- Once types of ideation are determined, few follow-up questions about most severe thought
  - Frequency
  - Duration
  - Controllability
  - Deterrents
  - Reasons for ideation (stop the pain or make someone angry—stop the pain is worse)
- Gives you a 2-25 score that will help inform clinical judgment about risk

All these items significantly predictive of suicide (on SSI)/minimum amount of info needed for tracking and severity

19

Clinical Monitoring Guidance

- For Intensity of Ideation, risk is greater when:
  - Thoughts are more frequent
  - Thoughts are of longer duration
  - Thoughts are less controllable
  - Fewer deterrents to acting on thoughts
  - Stopping the pain is the reason

… systematically assessing using the C-SSRS decreases burden

Jeffrey Garbelman, Ph.D.

jeffrey.garbelman@va.gov
jeffrey.garbelman@wi.gov
jeffrey.garbelman@att.net

For questions and other inquiries,
email Dr. Kelly Posner at:
posnerk@nyspi.columbia.edu

Website address for more information
on the C-SSRS:
http://www.cssrs.columbia.edu/

44