



Wisconsin Public Psychiatry Network Teleconference (WPPNT)

- This teleconference is brought to you by the Wisconsin Department of Health Services (DHS) Bureau of Prevention, Treatment and Recovery and the University of Wisconsin-Madison, Department of Psychiatry.
- The Department of Health Services makes no representations or warranty as to the accuracy, reliability, timeliness, quality, suitability or completeness of or results of the materials in this presentation. Use of information contained in this presentation may require express authority from a third party.



PROMOTING RECOVERY
PROPS, RENEÉ SUTTAN & ERICA O'REILLY



WHY EARLY INTERVENTION?


Duration of Untreated Psychosis: median of 74 Weeks

Medications: lower dose, longer time, less side effects

Family Involvement: Increase supports and education

Proactive with self medicating with alcohol, illicit drugs, tobacco

(RAISE)



PRODROME AND PSYCHOSIS

Prodromal Symptom/Clinical High Risk:
Occurs before first episode of psychosis
Subtle changes that are affecting functioning, such as:
changes in mood, sleep or appetite
changes in concentration
changes in thought

Psychosis:
The experience of loss of contact with reality, and is not part of the person's cultural group belief system of experiences. Psychosis typically involves hallucinations and/or delusions (NAMI).
10-20% of people who are at clinical high risk for schizophrenia or psychosis actually develop the illness (Kane)
Key Difference: In active psychosis, insight is limited (Compton & Brossard, 2009)

RECOVERY

Recovery is a process of change through which involved individuals improve their health and wellness. Recovery does not mean a cure. It is a process, not a destination.
Recovery is individual to each person and means working toward management of symptoms as well as areas of life that are the most important to an individual's overall well-being (SAMHSA, 2014)

Hope: The foundation of recovery
Resilience: Ability to adapt to challenges and change
Health: Exercise, substance, tobacco
Home: Safety and shelter, basic needs are met
Community: Support by peers, family, co-workers. Foundation of trust and stigma reduction
Purpose: Job, work, school, structure, "get up and go" (SAMHSA, 2014)

SHARED DECISION MAKING


Recovery based
Assumption that there are multiple experts in the room, consumers being one of the experts
Enhances the quality of care and informed decisions. Consumers are taking a part in their treatment, being accountable for themselves
Relationship and trust with providers is key (Adams & Drake, 2006)

"Collaborative process in which the consumer and the team member share knowledge and information and actively participate and treatment decisions, resulting in agreement on a preferred treatment approach" (OnTrack NY, Recovery Coach Manual)

CONSUMER FEEDBACK

"You guys have helped me learn a lot and get into the adult world"

"PROPs has greatly surpassed my expectations. PROPs has provided me with care and support. They are friendly and particularly helpful"



REFERRALS

Total referrals since February 2015: 47

Current Roster: 12

Discharges: 2 moved, 1 inappropriate dx, 2 inappropriate fit, 1 suicide death

Why were these referrals not appropriate?

Other dx, personality disorder dx or substance use dx: 8


Trauma related psychosis: 2

Too Young: 3

Didn't engage in outreach process: 11

Psychotic symptoms < 3 years: 4

At least 3 consumers who did not engage were most likely appropriate, the other 9 consumers declined formal assessment



DIAGNOSTIC CHALLENGES


Substance use and abuse complicating factor

Previous providers do not want to use dx of schizophrenia or psychosis (stigma)

Previous dx of schizophrenia

Lack of research resources and funds to use structured assessments

Trauma-based symptoms look remarkably similar to psychosis



MEDICATION TREATMENT

Just ONE of the important components of coordinated specialty care


Low dose monotherapy with antipsychotic medication

- Risperidone, aripiprazole, ziprasidone, lurasidone
- Perphenazine, loxapine

Common side effects:

- Sedation
- Restlessness
- Weight gain

Routine monitoring



PARENTS AND FAMILY


Primary supporters

Understanding of change in functioning

Consumers trust their parents

Need Support

- Parent Group
- Family Therapy



WORK/SCHOOL/HOSPITALIZATIONS

Consumers working:

- full time: 2
- part time: 1

Consumers enrolled in school (n=12):

- high school: 5
- college: 3

Consumer hospitalizations (n=18):

- hospitalizations before admission: 29
- hospitalizations post admission: 9



RESOURCES

Adams, J.R. & Drake, R.E. (2006). Shared Decision-Making and Evidence-Based Practice. *Community Mental Health Journal*, 42 (1).

Cohn, J. (2015) "Stop the Madness". <http://highline.huffingtonpost.com/articles/en/stop-the-madness/>

Compton, M. T. & Broussard, B. (2009). *The First Episode of Psychosis: A Guide for Patients and Their Families*. Oxford University Press.

Heinssen, R.K., Goldstein, A. B., & Azrin, S.T. (2014). Evidenced-Based Treatments for First Episode Psychosis: Components of Coordinated Specialty Care. RAISE.

Navigate. [Navigateconsultants.org](http://navigateconsultants.org)

OnTrack New York. Center for Practice Innovations. Unpublished, available online <http://practiceinnovations.org/OnTrackUSA/tabid/253/Default.aspx>

Prevent and Recovery in Early Psychosis. Prepwellness.org

Promoting Recovery from Onset of Psychosis. PropsJourneymhc.org

Recognition and Prevention Program, RAP.org

