

Wisconsin sexually transmitted disease surveillance report, 2011

2011 Manitowoc County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 185 | 100.0% | 230 | 17 | 100.0% | 21 | 0 | - | 0 | 202 | 100.0% | 251 |
| Sex | | | | | | | | | | | | |
| Male | 54 | 29.2% | 134 | 6 | 35.3% | 15 | - | - | - | 60 | 29.7% | 149 |
| Female | 131 | 70.8% | 325 | 11 | 64.7% | 27 | - | - | - | 142 | 70.3% | 352 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 108 | 58.4% | 139 | 10 | 58.8% | 13 | - | - | - | 118 | 58.4% | 152 |
| African American | 10 | 5.4% | 1,437 | 4 | 23.5% | 575 | - | - | - | 14 | 6.9% | 2,011 |
| American Indian | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 6 | 3.2% | 301 | 0 | 0.0% | 0 | - | - | - | 6 | 3.0% | 301 |
| Other | 2 | 1.1% | - | 0 | 0.0% | - | - | - | - | 2 | 1.0% | - |
| Unknown | 59 | 31.9% | - | 3 | 17.6% | - | - | - | - | 62 | 30.7% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 5 | 2.7% | 216 | 1 | 5.9% | 43 | - | - | - | 6 | 3.0% | 259 |
| Not Hispanic | 116 | 62.7% | 148 | 11 | 64.7% | 14 | - | - | - | 127 | 62.9% | 162 |
| Unknown | 64 | 34.6% | - | 5 | 29.4% | - | - | - | - | 69 | 34.2% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 3 | 1.6% | 58 | 0 | 0.0% | 0 | - | - | - | 3 | 1.5% | 58 |
| 15-19 | 41 | 22.2% | 754 | 3 | 17.6% | 55 | - | - | - | 44 | 21.8% | 809 |
| 20-24 | 92 | 49.7% | 2,001 | 8 | 47.1% | 174 | - | - | - | 100 | 49.5% | 2,175 |
| 25-29 | 19 | 10.3% | 423 | 3 | 17.6% | 67 | - | - | - | 22 | 10.9% | 490 |
| 30-34 | 20 | 10.8% | 504 | 2 | 11.8% | 50 | - | - | - | 22 | 10.9% | 555 |
| 35-39 | 4 | 2.2% | 87 | 0 | 0.0% | 0 | - | - | - | 4 | 2.0% | 87 |
| 40-44 | 1 | 0.5% | 17 | 0 | 0.0% | 0 | - | - | - | 1 | 0.5% | 17 |
| 45-49 | 2 | 1.1% | 29 | 1 | 5.9% | 14 | - | - | - | 3 | 1.5% | 43 |
| 50+ | 3 | 1.6% | 10 | 0 | 0.0% | 0 | - | - | - | 3 | 1.5% | 10 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 185 | 100.0% | 230 | 17 | 100.0% | 21 | - | - | - | 202 | 100.0% | 251 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



WI DEPARTMENT OF HEALTH SERVICES
 Division of Public Health
 Bureau of Communicable Diseases and Emergency Response
 STD Control Section
 P-00415B (2011)