

## Colorectal cancer screening, adults 50+

Health Priority A: Access to Primary and Preventive Health Services

Objective A2b: Increase provider screening for chronic diseases and other health risks.

2010 Target: No target established

Indicator: Colorectal cancer screening, adults 50+

### Percent of Wisconsin Adults Aged 50+ Who Have Ever Had a Blood Stool Test

Year (N)	Total		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-
2001	50%	3%	47%	4%	52%	4%
(N)	1,306		525		781	
2002	49%	2%	47%	4%	51%	3%
(N)	1,834		703		1,131	
2004	47%	3%	44%	4%	50%	4%
(N)	2,009		786		1,223	
2006	44%	2%	40%	4%	47%	3%
(N)	2,428		982		1,446	
2008	39%	2%	36%	3%	42%	3%
(N)	4,091		1,684		2,407	
2010	35%	2%	33%	3%	37%	3%
(N)	2,981		1,152		1,829	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

**Continued on next page**

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services

## Colorectal cancer screening, adults 50+, continued

### Percent of Wisconsin Adults Aged 50+ Who Have Ever Had a Blood Stool Test, by Race/Ethnicity

Years (N)	Total		African American*		White*	
	Percent	+/-	Percent	+/-	Percent	+/-
2001-2002 (N)	50%	2%	40%	7%	50%	2%
	3,140		170		2,777	
2002, 2004 (N)	48%	2%	36%	8%	48%	2%
	3,844		251		3,456	
2004, 2006 (N)	45%	2%	31%	7%	46%	2%
	4,467		297		3,988	
2006, 2008 (N)	41%	2%	32%	8%	42%	2%
	6,543		397		5,822	
2008, 2010 (N)	38%	2%	32%	8%	38%	2%
	7,010		513		6,164	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Other race categories are not shown, as there are fewer than 100 cases for ages 50+.

N        Number in sample. See data documentation.  
 +/-     Confidence interval. See data documentation.  
 \*        Non-Hispanic

**Continued on next page**

## Colorectal cancer screening, adults 50+, continued

### Percent of Wisconsin Adults Aged 50+ Who Have Ever Had a Sigmoidoscopy or Colonoscopy

Year (N)	Total		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-
2001	57%	3%	57%	4%	58%	3%
(N)	1,302		521		781	
2002	56%	2%	58%	4%	55%	3%
(N)	1,826		698		1,128	
2004	59%	3%	58%	4%	61%	3%
(N)	2,002		486		1,216	
2006	64%	2%	64%	4%	64%	3%
(N)	2,428		979		1,449	
2008	67%	2%	67%	4%	68%	3%
(N)	4,040		1,673		2,367	
2010	69%	2%	69%	3%	69%	3%
(N)	2,983		1,148		1,835	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Continued on next page

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services

## Colorectal cancer screening, adults 50+, continued

### Percent of Wisconsin Adults Aged 50+ Who Have Ever Had a Sigmoidoscopy or Colonoscopy, by Race/Ethnicity

Years (N)	Total		African American*		White*	
	Percent	+/-	Percent	+/-	Percent	+/-
2001-2002 (N)	57%	2%	49%	8%	57%	2%
	3,128		165		2,771	
2002, 2004 (N)	58%		56%	10%	58%	2%
	3,829		239		3,421	
2004, 2006 (N)	62%	2%	54%	8%	62%	2%
	4,459		294		3,986	
2006, 2008 (N)	66%	2%	66%	8%	67%	2%
	6,468		387		5,769	
2008, 2010 (N)	68%	2%	68%	8%	69%	2%
	7,007		514		6,160	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Other race categories are not shown, as there are fewer than 100 cases for ages 50+.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

\* Non-Hispanic.

**Continued on next page**

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services

## Colorectal cancer screening, adults 50+, continued

### Percent of Wisconsin Adults Aged 50+ Who Have Had a Sigmoidoscopy or Colonoscopy in the Past 5 Years

Year (N)	Total		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-
2001	46%	3%	46%	5%	46%	4%
(N)	1,294		518		776	
2002	47%	3%	49%	4%	45%	3%
(N)	1,822		698		1,124	
2004	50%	3%	50%	4%	51%	3%
(N)	1,980		780		1,200	
2006	54%	2%	54%	4%	55%	3%
(N)	2,419		978		1,441	
2008	57%	2%	57%	4%	57%	3%
(N)	4,069		1,677		2,392	
2010	56%	2%	58%	4%	55%	3%
(N)	2,953		1,137		1,816	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation.

Continued on next page

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services

**Percent of Wisconsin Adults Aged 50+ Who Have Had a Sigmoidoscopy or Colonoscopy in the Past 5 years, by Race/Ethnicity**

Years (N)	Total		African American*		White*	
	Percent	+/-	Percent	+/-	Percent	+/-
2001-2002	46%	2%	45%	10%	47%	3%
(N)	3,116		165		2,761	
2002,2004	49%	2%	52%	9%	49%	2%
(N)	3,802		224		3,402	
2004,2006	52%	2%	49%	8%	53%	2%
(N)	4,404		285		3,945	
2006, 2008	58%	2%	61%	8%	59%	2%
(N)	6,511		398		5,795	
2008, 2010	57%	2%	62%	9%	57%	2%
(N)	6,961		509		6,125	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: Other race categories are not shown, as there are fewer than 100 cases for ages 50+.

N        Number in sample. See data documentation.  
 +/-     Confidence interval. See data documentation.  
 \*        Non-Hispanic.