

Cigarette smoking, high school students

Health Priority C: Alcohol and Other Substance Use and Addiction

Objective C2c: Reduce the percentage of youth who report using tobacco in the past 30 days to 22.4 percent.

2010 Target: 22.4%

Indicator: Cigarette smoking, high school students

Percent of Wisconsin High School Students Who Smoked Cigarettes in the Past 30 Days

Year (N)	Total		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-
1999	38%	3%	38%	4%	38%	4%
(N)	1,336		682		648	
2001	33%	2%	29%	3%	37%	3%
(N)	2,120		1,031		1,084	
2003	24%	2%	23%	3%	24%	3%
(N)	2,121		1,019		1,097	
2005	23%	2%	24%	3%	22%	2%
(N)	2,389		1,163		1,219	
2007	20%	2%	21%	3%	20%	3%
(N)	2,013		1,012		1,000	
2009	17%	1%	18%	2%	16%	2%
(N)	2,368		1,152		1,213	

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation

Continued on next page

Cigarette smoking, continued

Percent of Wisconsin High School Students Who Smoked Cigarettes in the Past 30 Days, by Race/Ethnicity

Years (N)	Total		Afr. Am.*		Am. Ind.*		Asian*		Hispanic		White*	
	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-
1999-2003	31%	1%	17%	5%	53%	12%	26%	7%	37%	10%	32%	2%
(N)	5,577		397		104		213		154		4,379	
2001-2005	26%	1%	15%	4%	44%	9%	25%	6%	28%	7%	27%	1%
(N)	6,630		468		136		272		203		5,219	
2003-2007	22%	1%	14%	4%	40%	9%	20%	5%	23%	6%	23%	1%
(N)	6,317		316		111		246		183		5,111	
2005-2009	20%	1%	13%	4%	32%	9%	19%	5%	19%	6%	20%	1%
(N)	6,673		357		103		272		171		5,278	

Source: Wisconsin Youth Risk Behavior Survey, Department of Public Instruction.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation

*Non-Hispanic