

Immunizations, adults 65+

Health Priority E: Existing, Emerging, and Re-emerging Communicable Diseases

Objective E2c: By 2010, at least 90 percent of Wisconsin residents 65 years of age and older and individuals with chronic health conditions will be fully immunized in accordance with current Advisory Committee on Immunization Practices (ACIP) recommendations.

2010 Target: Greater than or equal to 90%

Indicator: Immunizations, adults 65+

Percent of Wisconsin Adults Aged 65 and Older Who Are Immunized in Accordance with ACIP Recommendations

Year (N)	Flu Shot in Past 12 Months		Pneumonia Shot (Ever)		Both	
	Percent	+/-	Percent	+/-	Percent	+/-
2001	70%	4%	65%	4%	55%	4%
(N)	590		590		590	
2002	74%	3%	71%	3%	61%	3%
(N)	847		829		829	
2003	72%	3%	67%	3%	58%	3%
(N)	805		783		783	
2004	74%	3%	70%	4%	62%	4%
(N)	921		895		895	
2005	72%	3%	66%	4%	56%	4%
(N)	1,121		1,083		1,082	
2006	72%	3%	72%	3%	59%	4%
(N)	1,063		1,032		1,054	
2007	74%	3%	69%	3%	61%	3%
(N)	1,928		1,880		1,880	
2008	73%	3%	70%	3%	60%	3%
(N)	1,863		1,826		1,832	
2009	72%	3%	70%	4%	60%	4%
(N)	1,280		1,260		1,259	
2010	68%	3%	73%	3%	59%	3%
(N)	1,356		1,333		1,328	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: ACIP also recommends tetanus and diphtheria boosters for adults aged 65+. Data on these immunizations are not reported in available sources.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation

Continued on next page

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services

Immunizations, adults 65+, continued

Percent of Wisconsin Adults Aged 65 and Older Who Are Immunized in Accordance with ACIP Recommendations, by Sex

Year (N)	Flu Shot in Past 12 months				Pneumonia Shot (Ever)				Both			
	Males		Females		Males		Females		Males		Females	
	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-	Percent	+/-
2001	75%	6%	67%	5%	63%	7%	67%	5%	56%	7%	56%	5%
(N)	204		384		196		379		196		379	
2002	76%	5%	73%	4%	68%	6%	72%	4%	60%	6%	61%	4%
(N)	281		566		267		562		267		562	
2003	71%	5%	73%	4%	62%	6%	70%	4%	54%	6%	60%	4%
(N)	285		520		272		511		272		511	
2004	76%	5%	73%	4%	67%	6%	73%	5%	61%	6%	62%	5%
(N)	310		611		298		597		298		597	
2005	74%	5%	71%	5%	64%	6%	67%	4%	57%	6%	55%	5%
(N)	398		723		376		707		376		706	
2006	72%	5%	72%	4%	71%	5%	73%	4%	60%	6%	59%	4%
(N)	387		681		362		670		369		685	
2007	72%	5%	76%	3%	66%	5%	72%	4%	59%	6%	63%	4%
(N)	686		1,242		652		1,228		652		1,228	
2008	72%	5%	74%	4%	67%	5%	73%	4%	57%	5%	62%	4%
(N)	705		1,158		679		1,147		682		1,150	
2009	68%	6%	75%	4%	64%	6%	74%	4%	56%	6%	63%	5%
(N)	536		744		524		736		523		736	
2010	65%	5%	71%	4%	69%	5%	77%	3%	53%	5%	62%	4%
(N)	475		881		461		872		458		870	

Source: Wisconsin Behavioral Risk Factor Survey, Office of Health Informatics, Division of Public Health, Department of Health Services.

Note: ACIP also recommends tetanus and diphtheria boosters for adults aged 65+. Data on these immunizations are not reported in available sources.

N Number in sample. See data documentation.

+/- Confidence interval. See data documentation

Prepared for Tracking the State Health Plan, 2010—State-Level Data

<http://dhs.wisconsin.gov/statehealthplan/track2010> (July 2011)

Office of Health Informatics

Division of Public Health

Wisconsin Department of Health Services