

Syphilis Incidence

Health Priority F: High-Risk Sexual Behavior

Objective F3a: By 2010, the incidence of primary and secondary syphilis in Wisconsin will be 0.2 cases per 100,000 population.

2010 Target: 0.2 cases per 100,000 population

Indicator: Syphilis incidence

Reported Primary and Secondary Syphilis Incidence in Wisconsin, by Sex

Year	Cases per 100,000 Population	Reported Cases		
		Total	Males	Females
2000	0.9	47	22	25
2001	0.5	25	12	13
2002	0.5	30	24	6
2003	0.3	15	14	1
2004	0.6	32	29	3
2005	0.7	41	39	2
2006	1.2	66	63	3
2007	1.2	68	62	6
2008	1.1	65	51	14
2009	0.8	45	35	10

Source: Bureau of Communicable Diseases, Division of Public Health, Department of Health Services.

Reported Primary and Secondary Syphilis Incidence in Wisconsin, by Race

Year	Cases per 100,000 Population	Reported Cases						
		Total	African American*	American Indian *	Asian *	Hispanic	White *	Other, Unknown
2000	0.9	47	43	2	0	1	0	1
2001	0.5	25	18	0	1	1	5	0
2002	0.5	30	9	1	0	3	15	2
2003	0.3	15	3	1	0	0	11	0
2004	0.6	32	3	0	0	3	26	0
2005	0.7	41	10	1	0	3	27	0
2006	1.2	65	17	0	1	8	38	1
2007	1.2	68	33	0	1	8	25	1
2008	1.1	65	35	0	1	2	26	1
2009	0.8	45	26	0	1	1	14	3

Source: Bureau of Communicable Diseases, Division of Public Health, Department of Health Services.

* Non-Hispanic

Continued on next page

Syphilis Incidence, continued

Reported Primary and Secondary Syphilis Incidence in Wisconsin, by Age

Year	Cases per 100,000 Population	Reported Cases						
		Total	Ages 0-9	Ages 10-14	Ages 15-19	Ages 20-24	Ages 25-29	Ages 30+
2000	0.9	47	0	0	3	5	5	34
2001	0.5	25	0	0	1	3	4	17
2002	0.5	30	0	0	0	2	8	20
2003	0.3	15	0	0	1	1	2	11
2004	0.6	32	0	0	1	2	4	25
2005	0.7	41	0	0	1	4	11	25
2006	1.2	65	0	0	4	8	8	45
2007	1.2	68	0	0	3	10	10	45
2008	1.1	65	0	0	4	11	15	35
2009	0.8	45	0	1	2	9	8	25

Source: Bureau of Communicable Diseases, Division of Public Health, Department of Health Services.