

Wisconsin sexually transmitted disease surveillance report, 2012

Bayfield County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 15 | 100.0% | 101 | 0 | - | 0 | 0 | - | 0 | 15 | 100.0% | 101 |
| Sex | | | | | | | | | | | | |
| Male | 4 | 26.7% | 54 | - | - | - | - | - | - | 4 | 26.7% | 54 |
| Female | 11 | 73.3% | 150 | - | - | - | - | - | - | 11 | 73.3% | 150 |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 5 | 33.3% | 38 | - | - | - | - | - | - | 5 | 33.3% | 38 |
| African American | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| American Indian | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 1 | 6.7% | 1,724 | - | - | - | - | - | - | 1 | 6.7% | 1,724 |
| Other | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Unknown | 9 | 60.0% | - | - | - | - | - | - | - | 9 | 60.0% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Not Hispanic | 4 | 26.7% | 27 | - | - | - | - | - | - | 4 | 26.7% | 27 |
| Unknown | 11 | 73.3% | - | - | - | - | - | - | - | 11 | 73.3% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 15-19 | 7 | 46.7% | 716 | - | - | - | - | - | - | 7 | 46.7% | 716 |
| 20-24 | 7 | 46.7% | 902 | - | - | - | - | - | - | 7 | 46.7% | 902 |
| 25-29 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 30-34 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 35-39 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 40-44 | 1 | 6.7% | 108 | - | - | - | - | - | - | 1 | 6.7% | 108 |
| 45-49 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 50+ | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Northern | 15 | 100.0% | 101 | - | - | - | - | - | - | 15 | 100.0% | 101 |
| Southeastern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
 Division of Public Health
 Bureau of Communicable Diseases and
 Emergency Preparedness
 STD Control Section
 P-00415B (2012)