

Wisconsin sexually transmitted disease surveillance report, 2011

2011 Rock County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 767 | 100.0% | 479 | 79 | 100.0% | 49 | 4 | 100.0% | 2 | 850 | 100.0% | 531 |
| Sex | | | | | | | | | | | | |
| Male | 212 | 27.6% | 267 | 28 | 35.4% | 35 | - | - | - | 243 | 28.6% | 306 |
| Female | 551 | 71.8% | 682 | 51 | 64.6% | 63 | - | - | - | 603 | 70.9% | 746 |
| Unknown | 4 | 0.5% | - | 0 | 0.0% | - | - | - | - | 4 | 0.5% | - |
| Race | | | | | | | | | | | | |
| White | 386 | 50.3% | 258 | 27 | 34.2% | 18 | - | - | - | 417 | 49.1% | 279 |
| African American | 159 | 20.7% | 1,944 | 32 | 40.5% | 391 | - | - | - | 191 | 22.5% | 2,335 |
| American Indian | 4 | 0.5% | 622 | 0 | 0.0% | 0 | - | - | - | 4 | 0.5% | 622 |
| Asian/Pacific Islander | 6 | 0.8% | 307 | 0 | 0.0% | 0 | - | - | - | 6 | 0.7% | 307 |
| Other | 21 | 2.7% | - | 1 | 1.3% | - | - | - | - | 22 | 2.6% | - |
| Unknown | 191 | 24.9% | - | 19 | 24.1% | - | - | - | - | 210 | 24.7% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 70 | 9.1% | 665 | 2 | 2.5% | 19 | - | - | - | 74 | 8.7% | 702 |
| Not Hispanic | 482 | 62.8% | 322 | 53 | 67.1% | 35 | - | - | - | 537 | 63.2% | 359 |
| Unknown | 215 | 28.0% | - | 24 | 30.4% | - | - | - | - | 239 | 28.1% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 1 | 0.1% | 9 | 0 | 0.0% | 0 | - | - | - | 1 | 0.1% | 9 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 12 | 1.6% | 111 | 0 | 0.0% | 0 | - | - | - | 12 | 1.4% | 111 |
| 15-19 | 260 | 33.9% | 2,305 | 23 | 29.1% | 204 | - | - | - | 283 | 33.3% | 2,509 |
| 20-24 | 275 | 35.9% | 2,762 | 31 | 39.2% | 311 | - | - | - | 306 | 36.0% | 3,073 |
| 25-29 | 125 | 16.3% | 1,167 | 12 | 15.2% | 112 | - | - | - | 138 | 16.2% | 1,289 |
| 30-34 | 53 | 6.9% | 541 | 6 | 7.6% | 61 | - | - | - | 60 | 7.1% | 613 |
| 35-39 | 23 | 3.0% | 220 | 6 | 7.6% | 57 | - | - | - | 29 | 3.4% | 277 |
| 40-44 | 10 | 1.3% | 89 | 1 | 1.3% | 9 | - | - | - | 12 | 1.4% | 107 |
| 45-49 | 4 | 0.5% | 32 | 0 | 0.0% | 0 | - | - | - | 4 | 0.5% | 32 |
| 50+ | 2 | 0.3% | 4 | 0 | 0.0% | 0 | - | - | - | 3 | 0.4% | 6 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southern | 767 | 100.0% | 479 | 79 | 100.0% | 49 | - | - | - | 850 | 100.0% | 531 |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



WI DEPARTMENT OF HEALTH SERVICES
 Division of Public Health
 Bureau of Communicable Diseases and Emergency Response
 STD Control Section
 P-00415B (2011)