

Wisconsin sexually transmitted disease surveillance report, 2013

Douglas County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 192 | 100.0% | 439 | 17 | 100.0% | 39 | 1 | 100.0% | 2 | 210 | 100.0% | 480 |
| Sex | | | | | | | | | | | | |
| Male | 50 | 26.0% | 232 | 3 | 17.6% | 14 | - | - | - | 53 | 25.2% | 246 |
| Female | 142 | 74.0% | 640 | 14 | 82.4% | 63 | - | - | - | 157 | 74.8% | 707 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 122 | 63.5% | 291 | 11 | 64.7% | 26 | - | - | - | 133 | 63.3% | 317 |
| African American | 6 | 3.1% | 1,460 | 1 | 5.9% | 243 | - | - | - | 8 | 3.8% | 1,946 |
| American Indian | 4 | 2.1% | 418 | 1 | 5.9% | 104 | - | - | - | 5 | 2.4% | 522 |
| Asian/Pacific Islander | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| Other | 1 | 0.5% | - | 0 | 0.0% | - | - | - | - | 1 | 0.5% | - |
| Unknown | 59 | 30.7% | - | 4 | 23.5% | - | - | - | - | 63 | 30.0% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 1 | 0.5% | 225 | 0 | 0.0% | 0 | - | - | - | 1 | 0.5% | 225 |
| Not Hispanic | 127 | 66.1% | 293 | 15 | 88.2% | 35 | - | - | - | 143 | 68.1% | 330 |
| Unknown | 64 | 33.3% | - | 2 | 11.8% | - | - | - | - | 66 | 31.4% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 1 | 0.5% | 37 | 0 | 0.0% | 0 | - | - | - | 1 | 0.5% | 37 |
| 15-19 | 54 | 28.1% | 1,880 | 13 | 76.5% | 453 | - | - | - | 67 | 31.9% | 2,333 |
| 20-24 | 79 | 41.1% | 2,585 | 2 | 11.8% | 65 | - | - | - | 82 | 39.0% | 2,683 |
| 25-29 | 34 | 17.7% | 1,051 | 1 | 5.9% | 31 | - | - | - | 35 | 16.7% | 1,082 |
| 30-34 | 16 | 8.3% | 622 | 0 | 0.0% | 0 | - | - | - | 16 | 7.6% | 622 |
| 35-39 | 6 | 3.1% | 222 | 0 | 0.0% | 0 | - | - | - | 6 | 2.9% | 222 |
| 40-44 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | - | - | - | 0 | 0.0% | 0 |
| 45-49 | 1 | 0.5% | 28 | 1 | 5.9% | 28 | - | - | - | 2 | 1.0% | 56 |
| 50+ | 1 | 0.5% | 7 | 0 | 0.0% | 0 | - | - | - | 1 | 0.5% | 7 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |
| Western | 192 | 100.0% | 439 | 17 | 100.0% | 39 | - | - | - | 210 | 100.0% | 480 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases and
Emergency Preparedness
STD Control Section
P-00415B (7/2014)