

Wisconsin sexually transmitted disease surveillance report, 2013

Sheboygan County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 312 | 100.0% | 272 | 32 | 100.0% | 28 | 7 | 100.0% | 6 | 351 | 100.0% | 306 |
| Sex | | | | | | | | | | | | |
| Male | 82 | 26.3% | 142 | 10 | 31.3% | 17 | 6 | 85.7% | 10 | 98 | 27.9% | 169 |
| Female | 230 | 73.7% | 406 | 22 | 68.8% | 39 | 1 | 14.3% | 2 | 253 | 72.1% | 446 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 229 | 73.4% | 213 | 22 | 68.8% | 20 | 5 | 71.4% | 5 | 256 | 72.9% | 238 |
| African American | 29 | 9.3% | 1,517 | 6 | 18.8% | 314 | 2 | 28.6% | 105 | 37 | 10.5% | 1,935 |
| American Indian | 2 | 0.6% | 362 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 2 | 0.6% | 362 |
| Asian/Pacific Islander | 13 | 4.2% | 273 | 1 | 3.1% | 21 | 0 | 0.0% | 0 | 14 | 4.0% | 294 |
| Other | 8 | 2.6% | - | 1 | 3.1% | - | 0 | 0.0% | - | 9 | 2.6% | - |
| Unknown | 31 | 9.9% | - | 2 | 6.3% | - | 0 | 0.0% | - | 33 | 9.4% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 35 | 11.2% | 611 | 2 | 6.3% | 35 | 2 | 28.6% | 35 | 39 | 11.1% | 681 |
| Not Hispanic | 225 | 72.1% | 207 | 25 | 78.1% | 23 | 5 | 71.4% | 5 | 255 | 72.6% | 234 |
| Unknown | 52 | 16.7% | - | 5 | 15.6% | - | 0 | 0.0% | - | 57 | 16.2% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 10-14 | 1 | 0.3% | 13 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 0.3% | 13 |
| 15-19 | 76 | 24.4% | 998 | 4 | 12.5% | 53 | 0 | 0.0% | 0 | 80 | 22.8% | 1,051 |
| 20-24 | 133 | 42.6% | 1,952 | 9 | 28.1% | 132 | 1 | 14.3% | 15 | 143 | 40.7% | 2,099 |
| 25-29 | 60 | 19.2% | 751 | 8 | 25.0% | 100 | 1 | 14.3% | 13 | 69 | 19.7% | 864 |
| 30-34 | 21 | 6.7% | 296 | 5 | 15.6% | 71 | 2 | 28.6% | 28 | 28 | 8.0% | 395 |
| 35-39 | 12 | 3.8% | 165 | 1 | 3.1% | 14 | 1 | 14.3% | 14 | 14 | 4.0% | 192 |
| 40-44 | 4 | 1.3% | 49 | 2 | 6.3% | 24 | 1 | 14.3% | 12 | 7 | 2.0% | 85 |
| 45-49 | 2 | 0.6% | 21 | 2 | 6.3% | 21 | 1 | 14.3% | 11 | 5 | 1.4% | 53 |
| 50+ | 3 | 1.0% | 8 | 1 | 3.1% | 3 | 0 | 0.0% | 0 | 4 | 1.1% | 10 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 312 | 100.0% | 272 | 32 | 100.0% | 28 | 7 | 100.0% | 6 | 351 | 100.0% | 306 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases and
Emergency Preparedness
STD Control Section
P-00415B (07/2014)