



Reported Cases of HIV Infection Wood County, Wisconsin, 1982-2015

	New Diagnoses by Year of Diagnosis(a)									Prevalence(c)		
	1982-2015(b)		2010-2014			2015			Cases	%	Rate(e)	
	Cases	%	Cases	Avg(d)	%	Rate(e)	Cases	%				Rate(e)
Total cases(f)	48	100.0%	5	1.0	100.0%	2.2	0	0.0%	0.0	41	100.0%	55.7
Disease Status(g)												
(Missing)	3	6.3%	0	0.0	0.0%	0.0	-	-	-	2	4.9%	2.7
HIV	35	72.9%	5	1.0	100.0%	2.2	-	-	-	13	31.7%	17.7
AIDS	10	20.8%	0	0.0	0.0%	0.0	-	-	-	26	63.4%	35.3
Sex at Birth												
Female	12	25.0%	1	0.2	20.0%	2.7	-	-	-	13	31.7%	34.9
Male	36	75.0%	4	0.8	80.0%	3.6	-	-	-	28	68.3%	77.0
Race/Ethnicity												
White	38	79.2%	-	-	-	-	-	-	-	33	80.5%	48.2
African American	-	-	-	-	-	-	-	-	-	-	-	-
Hispanic	5	10.4%	0	0.0	0.0%	0.0	-	-	-	-	-	-
American Indian	0	0.0%	0	0.0	0.0%	0.0	-	-	-	0	0.0%	0.0
Asian/Pacific Islander	0	0.0%	0	0.0	0.0%	0.0	-	-	-	0	0.0%	0.0
Multi-racial	-	-	0	0.0	0.0%	0.0	-	-	-	-	-	-
Unknown	-	-	0	0.0	0.0%	-	-	-	-	-	-	-
Age(h)												
(Missing)	2	4.2%	0	0.0	0.0%	0.0	-	-	-	2	4.9%	-
<5	1	2.1%	1	0.2	20.0%	23.7	-	-	-	0	0.0%	0.0
5-14	1	2.1%	0	0.0	0.0%	0.0	-	-	-	1	2.4%	11.2
15-19	1	2.1%	0	0.0	0.0%	0.0	-	-	-	0	0.0%	0.0
20-24	5	10.4%	0	0.0	0.0%	0.0	-	-	-	0	0.0%	0.0
25-29	6	12.5%	0	0.0	0.0%	0.0	-	-	-	0	0.0%	0.0
30-34	4	8.3%	1	0.2	20.0%	24.0	-	-	-	1	2.4%	24.0
35-39	12	25.0%	1	0.2	20.0%	25.7	-	-	-	1	2.4%	25.7
40-44	6	12.5%	1	0.2	20.0%	22.8	-	-	-	5	12.2%	118.5
45-49	6	12.5%	1	0.2	20.0%	16.5	-	-	-	6	14.6%	122.8
50-54	3	6.3%	0	0.0	0.0%	0.0	-	-	-	12	29.3%	194.9
55-59	1	2.1%	0	0.0	0.0%	0.0	-	-	-	6	14.6%	101.4
60+	0	0.0%	0	0.0	0.0%	0.0	-	-	-	7	17.1%	37.4
Risk exposure												
MSM	24	50.0%	4	0.8	80.0%	-	-	-	-	18	43.9%	-
IDU	5	10.4%	0	0.0	0.0%	-	-	-	-	3	7.3%	-
MSM & IDU	4	8.3%	0	0.0	0.0%	-	-	-	-	2	4.9%	-
Heterosexual	5	10.4%	0	0.0	0.0%	-	-	-	-	12	29.3%	-
Other/Unknown	10	20.8%	1	0.2	20.0%	-	-	-	-	6	14.6%	-

Year of HIV Diagnosis	Cases	Rate(e)
Before 2006	38	-
2006	1	1.4
2007	1	1.4
2008	0	0.0
2009	3	4.1
2010	1	1.3
2011	0	0.0
2012	0	0.0
2013	1	1.4
2014	3	4.1
2015	0	0.0

a. New diagnoses include only individuals whose initial HIV diagnosis was made in Wisconsin.
b. The first cases of HIV infection in Wisconsin were diagnosed in 1982. Thus, these represent cumulative cases from 1982 through the specified date.
c. Prevalent cases include all cases presumed to be alive and living in Wisconsin, regardless of the state of initial HIV report.
d. The average annual cases in the specified period.
e. Cases per 100,000 population. Rates not available for risk exposure groups.
f. Breakdown not shown if total is < 5. See Technical Notes for suppression rule for race/ethnicity breakdown.
g. Disease status when first diagnosed with HIV infection, except for prevalent cases, where it is the current disease status.
h. Age when first diagnosed with HIV infection, except for prevalent cases, where it is the current age.