

**Reported Cases of HIV Infection
WISCONSIN, 2014-2019**

	New Diagnoses by Year of Diagnosis ^a									2019 Prevalence ^b		
	2014-2018				2019							
	Cases	Avg ^c	%	Rate ^d	Cases	%	Rate ^d	Cases	%	Rate ^d		
Total cases^e	1,142	228.4	100.0%	4.0	214	100.0%	3.7	6,750	100.0%	116.8		
Disease status^f												
(missing)	4	0.8	0.4%	0.0	0	0.0%	0.0	3	0.0%	0.1		
HIV	897	179.4	78.5%	3.1	169	79.0%	2.9	3,535	52.4%	61.1		
HIV Stage 3 (AIDS)	241	48.2	21.1%	0.8	45	21.0%	0.8	3,212	47.6%	55.6		
Gender^g												
Female	155	31.0	13.6%	1.1	43	20.1%	1.5	1,353	20.0%	46.6		
Male	962	192.4	84.2%	6.7	167	78.0%	5.8	5,302	78.5%	184.4		
Race/ethnicity^h												
White	403	80.6	35.3%	1.7	68	31.8%	1.4	2,885	42.7%	60.9		
Black/African American	495	99.0	43.3%	25.0	101	47.2%	25.1	2,592	38.4%	644.0		
Hispanic	168	33.6	14.7%	8.7	33	15.4%	8.2	946	14.0%	235.3		
American Indian	8	1.6	0.7%	2.8	-	-	-	32	0.5%	55.8		
Asian/Pacific Islander	26	5.2	2.3%	3.0	-	-	-	102	1.5%	56.1		
Multi-racial or unknown	42	8.4	3.7%	-	-	-	-	193	2.9%	-		
Ageⁱ												
0-14	2	0.4	0.2%	0.0	1	0.5%	0.1	30	0.4%	2.9		
15-19	65	13.0	5.7%	3.4	11	5.1%	2.9	33	0.5%	8.8		
20-29	469	93.8	41.1%	12.3	83	38.8%	10.8	661	9.8%	86.3		
30-39	245	49.0	21.5%	6.8	46	21.5%	6.3	1,206	17.9%	165.3		
40-49	167	33.4	14.6%	4.8	38	17.8%	5.6	1,378	20.4%	202.3		
50-59	145	29.0	12.7%	3.5	24	11.2%	3.0	2,006	29.7%	248.7		
60+	49	9.8	4.3%	0.8	11	5.1%	0.8	1,436	21.3%	104.6		
Risk exposure												
Gay or bisexual male (MSM)	739	147.8	64.7%	-	118	55.1%	-	3,667	54.3%	-		
Injection drug use (IDU)	39	7.8	3.4%	-	11	5.1%	-	481	7.1%	-		
MSM and IDU	33	6.6	2.9%	-	9	4.2%	-	339	5.0%	-		
Heterosexual	91	18.2	8.0%	-	22	10.3%	-	938	13.9%	-		
Perinatal exposure	2	0.4	0.2%	-	1	0.5%	-	70	1.0%	-		
Other or unknown	238	47.6	20.8%	-	53	24.8%	-	1,255	18.6%	-		

WISCONSIN		
Year of HIV diagnosis	Cases	Rate ^d
2010	252	4.4
2011	244	4.3
2012	223	3.9
2013	248	4.3
2014	221	3.8
2015	227	3.9
2016	224	3.9
2017	257	4.4
2018	213	3.7
2019	214	3.7

a. New diagnoses include only individual whose initial HIV diagnosis was made in Wisconsin.

b. Prevalent cases include all cases presumed to be alive and living in Wisconsin, regardless of the state of initial HIV report.

c. The average annual cases in the specified time period.

d. Cases per 100,000 population. Rates not available for risk exposure groups.

e. Breakdowns are not shown if the total is <5.

f. Disease status when first diagnosed with HIV infection, except for prevalent cases, where it is the current disease status.


g. Cases are presented by current gender rather than sex at birth. During 2014-2018, a total of 25 transgender persons were newly diagnosed with HIV in Wisconsin.

h. Race/ethnicity breakdowns are not shown if the cell in the subgroup has either a value of <5 or an estimated population size (denominator) <100. In such cases, the multi-racial or unknown

category along with any cells that equal zero are suppressed to prevent interpolation of the suppressed cell(s). Rates for multi-racial persons are not shown because population denominators for this group are not available in the demographic data source.


i. Age when first diagnosed with HIV infection, except for prevalence cases, where it is the current age.

Geographic distribution of new HIV diagnoses, Wisconsin, 2019 (n=214)



*Excludes 7 people diagnosed in the Department of Corrections
 Source: Wisconsin Department of Health Services

Geographic distribution of people living with HIV, Wisconsin, 2019 (n=6,749)



Source: Wisconsin Department of Health Services