

Sexually Transmitted Infections

Wisconsin 2020




Sexually Transmitted Infections Surveillance Report Data Tables, Wisconsin, 2020

**Table 1. Number of Cases and Rates of Chlamydia, Gonorrhea, Syphilis
In Persons 15-19 Years of Age, Wisconsin, 2016-2020**

| Disease | 2016 | | 2017 | | 2018 | | 2019 | | 2020 | |
|-----------|-------|--------|-------|--------|-------|--------|-------|--------|--------|--------|
| | Cases | Rates* | Cases | Rates* | Cases | Rates* | Cases | Rates* | Cases* | Rates* |
| Chlamydia | 7,369 | 1,852 | 7,303 | 1,836 | 7,482 | 1,881 | 8,197 | 2,061 | 7,015 | 1,852 |
| Gonorrhea | 1,632 | 410 | 695 | 340 | 1,679 | 422 | 1,761 | 443 | 1,767 | 467 |
| Syphilis | 19 | 5 | 28 | 13 | 38 | 10 | 34 | 9 | 32 | 8 |

*Rates per 100,000 population



Because of the Covid-19 pandemic, STI testing services were running at a minimum level in both STD and non-STD clinics and local health departments. Chlamydia cases decreased from 8,197 to 7,015 (14.4% decrease), and no significant change in Gonorrhea and Syphilis cases in 2020 observed.

Wisconsin Sexually Transmitted Disease Surveillance Report, 2020 Persons 15-19 years of age

| | Chlamydia | | | Gonorrhea | | | Syphilis | | | Total | | |
|----------------------------|-----------|---------|--------|-----------|---------|--------|----------|---------|--------|-------|---------|--------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Total | 7,015 | 100% | 1,852 | 1,767 | 100% | 467 | 32 | 100% | 8 | 8,814 | 100% | 2,327 |
| Gender | | | | | | | | | | | | |
| Male | 1,647 | 23.5% | 852 | 677 | 38.3% | 350 | 16 | 50.0% | 8 | 2,340 | 26.5% | 1,210 |
| Female | 5,354 | 76.3% | 2,888 | 1,079 | 61.1% | 582 | 15 | 46.9% | 8 | 6,448 | 73.2% | 3,478 |
| Transgender Female | 1 | 0.0% | - | 2 | 0.1% | - | 1 | 3.1% | - | 4 | 0.0% | - |
| Transgender Male | 1 | 0.0% | - | 1 | 0.1% | - | 0 | 0.0% | - | 2 | 0.0% | - |
| Other/Unknown | 12 | 0.2% | - | 8 | 0.5% | - | 0 | 0.0% | - | 20 | 0.2% | - |
| Race/Ethnicity | | | | | | | | | | | | |
| White | 2,285 | 32.6% | 808 | 280 | 15.8% | 99 | 7 | 22% | 2^ | 2,572 | 29.2% | 910 |
| Black | 2,513 | 35.8% | 7,113 | 1,007 | 57.0% | 2,850 | 19 | 59% | 54 | 3,539 | 40.2% | 10,018 |
| Native American | 64 | 0.9% | 1,455 | 6 | 0.3% | 136^ | 0 | 0.0% | 0 | 70 | 0.8% | 1,591 |
| Asian | 59 | 0.8% | 412 | 12 | 0.7% | 84 | 0 | 0.0% | 0 | 71 | 0.8% | 495 |
| Hispanic | 701 | 10.0% | 1,672 | 145 | 8.2% | 346 | 4 | 12.5% | ** | 850 | 9.6% | 2,027 |
| Other/Unknown | 1,393 | 19.9% | - | 317 | 17.9% | - | 2 | 6.3% | - | 1,712 | 19.4% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 1,036 | 14.8% | 1,311 | 148 | 8.4% | 187 | 1 | 3.1% | ** | 1,185 | 13.4% | 1,500 |
| Northern | 299 | 4.3% | 1,011 | 28 | 1.6% | 95 | 0 | 0.0% | 0 | 327 | 3.7% | 1,106 |
| Southeastern | 3,874 | 55.2% | 2,693 | 1,219 | 69.0% | 847 | 24 | 75.0% | 17 | 5,117 | 58.1% | 3,716 |
| Southern | 1,200 | 17.1% | 1,679 | 312 | 17.7% | 437 | 6 | 18.8% | 8^ | 1,518 | 17.2% | 1,955 |
| Western | 606 | 8.6% | 1,106 | 60 | 3.4% | 109 | 1 | 3.1% | ** | 667 | 7.6% | 1,217 |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

Wisconsin Sexually Transmitted Disease Surveillance Report, 2020
Reported Cases by Region and County
Persons 15-19 years of age

| | Chlamydia | | | Gonorrhea | | | Syphilis | | | Total | | |
|---------------------|--------------|--------------|--------------|--------------|-------------|------------|-----------|-------------|-----------|--------------|--------------|--------------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Total Case | 7,015 | 100% | 1,852 | 1,767 | 100% | 467 | 32 | 100% | 8 | 8,814 | 100% | 2,327 |
| Northeastern | | | | | | | | | | | | |
| Brown | 274 | 3.9% | 1,606 | 51 | 2.9% | 299 | 0 | 0.0% | 0 | 325 | 3.7% | 1,904 |
| Calumet | 18 | 0.3% | 484 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 18 | 0.2% | 484 |
| Door | 13 | 0.2% | 1,019 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 14 | 0.2% | 1,097 |
| Fond Du Lac | 75 | 1.1% | 1,142 | 13 | 0.7% | 198 | 0 | 0.0% | 0 | 88 | 1.0% | 1,340 |
| Green Lake | 12 | 0.2% | 983 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 13 | 0.1% | 1,065 |
| Kewaunee | 9 | 0.1% | 677^ | 1 | 0.1% | ** | 0 | 0.0% | 0 | 10 | 0.1% | 752^ |
| Manitowoc | 55 | 0.8% | 1,160 | 6 | 0.3% | 127^ | 0 | 0.0% | 0 | 61 | 0.7% | 1,287 |
| Marinette | 35 | 0.5% | 1,491 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 36 | 0.4% | 1,533 |
| Marquette | 5 | 0.1% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 5 | 0.1% | ** |
| Menominee | 9 | 0.1% | 2,961^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 9 | 0.1% | 2,961^ |
| Oconto | 32 | 0.5% | 1,437^ | 2 | 0.1% | ** | 0 | 0.0% | 0 | 34 | 0.4% | 1,527 |
| Outagamie | 126 | 1.8% | 1,040 | 23 | 1.3% | 190 | 1 | 3.1% | ** | 150 | 1.7% | 1,239 |
| Shawano | 26 | 0.4% | 1,042 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 28 | 0.3% | 1,122 |
| Sheboygan | 96 | 1.4% | 1,315 | 10 | 0.6% | 137^ | 0 | 0.0% | 0 | 106 | 1.2% | 1,452 |
| Waupaca | 33 | 0.5% | 1,125 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 35 | 0.4% | 1,193 |
| Waushara | 18 | 0.3% | 1,365 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 19 | 0.2% | 1,440 |
| Winnebago | 200 | 2.9% | 1,774 | 34 | 1.9% | 301 | 0 | 0.0% | 0 | 234 | 2.7% | 2,075 |
| Region total | 1,036 | 14.8% | 1,311 | 148 | 8.4% | 187 | 1 | 3.1% | ** | 1,185 | 13.4% | 1,500 |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

| | Chlamydia | | | Gonorrhea | | | Syphilis | | | Total | | |
|---------------------|-----------|---------|--------|-----------|---------|--------|----------|---------|--------|-------|---------|--------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Northern | | | | | | | | | | | | |
| Ashland | 14 | 0.2% | 1,389 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 14 | 0.2% | 1,389 |
| Bayfield | 6 | 0.1% | 881^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 6 | 0.1% | 881^ |
| Florence | 1 | 0.0% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 0.0% | ** |
| Forest | 12 | 0.2% | 2,147 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 13 | 0.1% | 2,326 |
| Iron | 5 | 0.1% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 5 | 0.1% | ** |
| Langlade | 11 | 0.2% | 1,007^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 11 | 0.1% | 1,007^ |
| Lincoln | 15 | 0.2% | 907 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 17 | 0.2% | 1,028 |
| Marathon | 99 | 1.4% | 1,150 | 11 | 0.6% | 128^ | 0 | 0.0% | 0 | 110 | 1.2% | 1,278 |
| Oneida | 16 | 0.2% | 961 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 16 | 0.2% | 961 |
| Portage | 51 | 0.7% | 893 | 10 | 0.6% | 175^ | 0 | 0.0% | 0 | 61 | 0.7% | 1,068 |
| Price | 10 | 0.1% | 1,479^ | 1 | 0.1% | ** | 0 | 0.0% | 0 | 11 | 0.1% | 1,627^ |
| Sawyer | 2 | 0.0% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 2 | 0.0% | ** |
| Taylor | 8 | 0.1% | 611^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 8 | 0.1% | 611^ |
| Vilas | 7 | 0.1% | 689^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 7 | 0.1% | 689^ |
| Wood | 42 | 0.6% | 979 | 3 | 0.2% | ** | 0 | 0.0% | 0 | 45 | 0.5% | 1,049 |
| Region total | 299 | 4.3% | 1,011 | 28 | 1.6% | 95 | 0 | 0.0% | 0 | 327 | 3.7% | 1,106 |
| | | | | | | | | | | | | |
| Southeastern | | | | | | | | | | | | |
| Kenosha | 206 | 2.9% | 1,716 | 57 | 3.2% | 475 | 0 | 0.0% | 0 | 263 | 3.0% | 2,191 |
| Milwaukee | 2,992 | 42.7% | 4,616 | 992 | 56.1% | 1,531 | 19 | 59.4% | 29 | 4,003 | 45.4% | 6,176 |
| Ozaukee | 28 | 0.4% | 467 | 9 | 0.5% | 150^ | 1 | 3.1% | ** | 38 | 0.4% | 634 |
| Racine | 308 | 4.4% | 2,493 | 122 | 6.9% | 988 | 1 | 3.1% | ** | 431 | 4.9% | 3,489 |
| Walworth | 65 | 0.9% | 966 | 5 | 0.3% | ** | 0 | 0.0% | 0 | 70 | 0.8% | 1,040 |
| Washington | 51 | 0.7% | 584 | 3 | 0.2% | ** | 0 | 0.0% | 0 | 54 | 0.6% | 618 |
| Waukesha | 165 | 2.4% | 610 | 25 | 1.4% | 92 | 3 | 9.4% | ** | 193 | 2.2% | 713 |
| Region Total | 3,874 | 54.4% | 2,938 | 1,219 | 69.0% | 924 | 24 | 75.0% | 18 | 5,117 | 58.1% | 3,880 |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

| | Chlamydia | | | Gonorrhea | | | Syphilis | | | Total | | |
|---------------------|-----------|---------|--------|-----------|---------|--------|----------|---------|-------|-------|---------|-------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rate* | Cases | Percent | Rate* |
| Southern | | | | | | | | | | | | |
| Adams | 6 | 0.1% | 716^ | 1 | 0.1% | ** | 0 | 0.0% | 0 | 7 | 0.1% | 835^ |
| Columbia | 45 | 0.6% | 1,279 | 7 | 0.4% | 199^ | 0 | 0.0% | 0 | 52 | 0.6% | 1,478 |
| Crawford | 6 | 0.1% | 646^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 6 | 0.1% | 646^ |
| Dane | 684 | 9.8% | 2,131 | 208 | 11.8% | 648 | 5 | 15.6% | ** | 897 | 10.2% | 2,794 |
| Dodge | 35 | 0.5% | 663 | 5 | 0.3% | ** | 0 | 0.0% | 0 | 40 | 0.5% | 758 |
| Grant | 29 | 0.4% | 633 | 2 | 0.1% | ** | 1 | 3.1% | ** | 32 | 0.4% | 699 |
| Green | 20 | 0.3% | 870 | 3 | 0.2% | ** | 0 | 0.0% | 0 | 23 | 0.3% | 1,000 |
| Iowa | 6 | 0.1% | 416^ | 1 | 0.1% | ** | 0 | 0.0% | 0 | 7 | 0.1% | 485^ |
| Jefferson | 59 | 0.8% | 953 | 6 | 0.3% | 97^ | 0 | 0.0% | 0 | 65 | 0.7% | 1,050 |
| Juneau | 32 | 0.5% | 2,213 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 32 | 0.4% | 2,213 |
| Lafayette | 12 | 0.2% | 1,028 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 14 | 0.2% | 1,200 |
| Richland | 8 | 0.1% | 690^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 8 | 0.1% | 690^ |
| Rock | 241 | 3.4% | 2,240 | 79 | 4.5% | 734 | 0 | 0.0% | 0 | 320 | 3.6% | 2,975 |
| Sauk | 55 | 0.8% | 1,443 | 3 | 0.2% | ** | 0 | 0.0% | 0 | 58 | 0.7% | 1,522 |
| Vernon | 21 | 0.3% | 987 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 22 | 0.2% | 1,034 |
| Region total | 1,200 | 17.9% | 1,438 | 312 | 17.7% | 374 | 6 | 18.8% | 7^ | 1518 | 17.2% | 1,819 |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

| Western | Chlamydia | | | Gonorrhea | | | Syphilis | | | Total | | |
|--------------|------------|-------------|-------------|-----------|-------------|------------|----------|-------------|-----------|------------|-------------|-------------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Barron | 25 | 0.4% | 966 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 25 | 0.3% | 966 |
| Buffalo | 4 | 0.1% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 4 | 0.0% | ** |
| Burnett | 10 | 0.1% | 1319^ | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 10 | 0.1% | 1319^ |
| Chippewa | 42 | 0.6% | 1080 | 4 | 0.2% | ** | 0 | 0.0% | 0 | 46 | 0.5% | 1183 |
| Clark | 15 | 0.2% | 591 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 16 | 0.2% | 630 |
| Douglas | 27 | 0.4% | 1066 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 29 | 0.3% | 1144 |
| Dunn | 30 | 0.4% | 772 | 5 | 0.3% | ** | 0 | 0.0% | 0 | 35 | 0.4% | 901 |
| Eau Claire | 81 | 1.2% | 993 | 15 | 0.8% | 184 | 0 | 0.0% | 0 | 96 | 1.1% | 1177 |
| Jackson | 16 | 0.2% | 1330 | 2 | 0.1% | ** | 1 | 3.1% | ** | 19 | 0.2% | 1579 |
| La Crosse | 193 | 2.6% | 2150 | 18 | 1.0% | 201 | 0 | 0.0% | 0 | 211 | 2.4% | 2351 |
| Monroe | 30 | 0.4% | 992 | 2 | 0.1% | ** | 0 | 0.0% | 0 | 32 | 0.4% | 1058 |
| Pepin | 4 | 0.1% | ** | 1 | 0.1% | ** | 0 | 0.0% | 0 | 5 | 0.1% | ** |
| Pierce | 27 | 0.4% | 750 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 28 | 0.3% | 778 |
| Polk | 33 | 0.5% | 1211 | 4 | 0.2% | ** | 0 | 0.0% | 0 | 37 | 0.4% | 1357 |
| Rusk | 4 | 0.1% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 4 | 0.0% | ** |
| Saint Croix | 37 | 0.5% | 589 | 4 | 0.2% | ** | 0 | 0.0% | 0 | 41 | 0.5% | 653 |
| Trempealeau | 14 | 0.2% | 763 | 1 | 0.1% | ** | 0 | 0.0% | 0 | 15 | 0.2% | 817 |
| Washburn | 14 | 0.2% | 1757 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 14 | 0.2% | 1757 |
| Total | 606 | 8.6% | 1106 | 60 | 3.4% | 109 | 1 | 3.1% | ** | 667 | 7.6% | 1217 |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

Wisconsin Sexually Transmitted Disease Surveillance Report: 2020

Syphilis Cases By Disease Stage

Persons 15-19 years of age

| | Primary | | | Secondary | | | Early Latent | | | Early Total | | |
|----------------------------|---------|---------|--------|-----------|---------|--------|--------------|---------|--------|-------------|---------|--------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Total | 6 | 100% | 2^ | 4 | 100% | *** | 10 | 100% | 3^ | 20 | 100% | 5 |
| Gender | | | | | | | | | | | | |
| Male | 4 | 66.7% | ** | 1 | 25.0% | ** | 6 | 60.0% | 3^ | 11 | 55.0% | 6^ |
| Female | 2 | 33.3% | ** | 3 | 75.0% | ** | 4 | 40.0% | ** | 9 | 45.0% | 5^ |
| Transgender Female | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Transgender Male | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race/Ethnicity | | | | | | | | | | | | |
| White | 1 | 16.7% | ** | 1 | 25.0% | ** | 0 | 0.0% | 0 | 2 | 10.0% | ** |
| Black | 5 | 83.3% | ** | 0 | 0.0% | 0 | 7 | 70.0% | 20^ | 12 | 60.0% | 34 |
| Native American | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | - |
| Asian | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | - |
| Hispanic | 0 | 0.0% | 0 | 3 | 75.0% | ** | 1 | 10.0% | ** | 4 | 20.0% | ** |
| Other/Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 2 | 20.0% | - | 2 | 10.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | ** |
| Northern | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| Southeastern | 4 | 67.7% | ** | 4 | 100% | ** | 10 | 100.0% | 7^ | 18 | 90.0% | 13 |
| Southern | 2 | 33.3% | ** | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 2 | 10.0% | ** |
| Western | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | ** |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

Wisconsin Sexually Transmitted Disease Surveillance Report: 2020
Syphilis Cases By Disease Stage
Persons 15-19 years of age

| | Early Total | | | Late Latent | | | Congenital | | | All Stages Total | | |
|----------------------------|-------------|---------|--------|-------------|---------|--------|------------|---------|--------|------------------|---------|--------|
| | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* | Cases | Percent | Rates* |
| Total | 20 | 100% | 5 | 12 | 100% | 3 | 0 | 100% | 0 | 32 | 100% | 8 |
| Gender | | | | | | | | | | | | |
| Male | 11 | 55.0% | 6^ | 5 | 41.7% | ** | 0 | 0.0% | 0 | 16 | 50.0% | 8 |
| Female | 9 | 45.0% | 5^ | 6 | 50.0% | 3^ | 0 | 0.0% | 0 | 15 | 46.9% | 8 |
| Transgender Female | 0 | 0.0% | - | 1 | 8.3% | ** | 0 | 0.0% | - | 1 | 3.1% | - |
| Transgender Male | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race/Ethnicity | | | | | | | | | | | | |
| White | 2 | 10.0% | ** | 5 | 41.7% | ** | 0 | 0.0% | 0 | 7 | 21.9% | 2^ |
| Black | 12 | 60.0% | 34 | 7 | 58.3% | 20^ | 0 | 0.0% | 0 | 19 | 59.4% | 54 |
| Native American | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | - |
| Asian | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | - |
| Hispanic | 4 | 20.0% | ** | 0 | 0.0% | - | 0 | 0.0% | 0 | 4 | 12.5% | ** |
| Other/Unknown | 2 | 10.0% | ** | 0 | 0.0% | - | 0 | 0.0% | 0 | 2 | 6.3% | ** |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 1 | 8.3% | ** | 0 | 0.0% | 0 | 1 | 3.1% | * |
| Northern | 0 | 0.0% | - | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| Southeastern | 18 | 90.0% | 13 | 6 | 50.0% | 4^ | 0 | 0.0% | 0 | 24 | 75.0% | 17 |
| Southern | 2 | 10.0% | ** | 4 | 33.3% | ** | 0 | 0.0% | 0 | 6 | 18.8% | 8^ |
| Western | 0 | 0.0% | - | 1 | 8.3% | ** | 0 | 0.0% | 0 | 1 | 3.1% | ** |

*Cases per 100,000 population.

**Rates based on case counts five or less than five are not shown.

^Rate is unreliable due to a count <12.

Technical Notes

1. These reports are compiled by the Wisconsin STI Unit and are based on STI case surveillance data collected by the Wisconsin Division of Public Health (DPH). In Wisconsin, state statutes require both laboratories and health care providers to report cases of chlamydia, gonorrhea, syphilis, chancroid, and pelvic inflammatory disease to local health departments. STI case surveillance data is automatically sent to the Wisconsin STI Unit via WEDSS (Wisconsin Electronic Disease Surveillance System).
2. These reports include cases diagnosed within the specified period. Additional cases diagnosed within the specified period may continue to be reported. Thus, this report should be considered provisional and it supersedes all previous reports.
3. Reporting of STI cases can be incomplete. Reporting rates can vary by disease, geographic region, and demographic groups. At any time, reported STI cases represent only part of the total number of diagnosed cases, therefore STI surveillance data likely underestimates total STI morbidity.
4. Rate is calculated based on the population data available on the WISH (Wisconsin Interactive Statistics on Health) system website. In these reports, rates presented are cases per 100,000 population (2019). Rates are not calculated for categories with small numbers of cases. See each report for the rate calculation threshold.
5. In these reports, cases are allocated to the county of residence at the time of diagnosis. Cases diagnosed among inmates of correctional facilities and cases residing outside Wisconsin are not included in this report.
6. All races are non-Hispanic ethnicity. Under the category race, Asian includes Asian Pacific Islander. Under the category race, American Indian Alaska Native is named as Native American.
7. Rates based on case counts five and less are not shown and rates based on counts less than 12 are marked as statistically unreliable.

For more information, contact: Wisconsin STI Unit
1 West Wilson Street
PO Box 2659
Madison, WI 53701-2659
Telephone: 608-266-8479