

Wisconsin sexually transmitted disease surveillance report, 2016

Winnebago County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 787 | 100.0% | 485 | 105 | 100.0% | 65 | 14 | 100.0% | 9 | 906 | 100.0% | 559 |
| Sex | | | | | | | | | | | | |
| Male | 239 | 30.4% | 293 | 53 | 50.5% | 65 | 11 | 78.6% | 14 | 303 | 33.4% | 372 |
| Female | 548 | 69.6% | 680 | 52 | 49.5% | 64 | 3 | 21.4% | 4 | 603 | 66.6% | 748 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 609 | 77.4% | 393 | 63 | 60.0% | 41 | 12 | 85.7% | 8 | 684 | 75.5% | 441 |
| African American | 86 | 10.9% | 3,040 | 31 | 29.5% | 1,096 | 1 | 7.1% | 35 | 118 | 13.0% | 4,171 |
| American Indian | 5 | 0.6% | 504 | 1 | 1.0% | 101 | 0 | 0.0% | 0 | 6 | 0.7% | 605 |
| Asian/Pacific Islander | 16 | 2.0% | 492 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 16 | 1.8% | 492 |
| Other | 22 | 2.8% | - | 4 | 3.8% | - | 1 | 7.1% | - | 27 | 3.0% | - |
| Unknown | 49 | 6.2% | - | 6 | 5.7% | - | 0 | 0.0% | - | 55 | 6.1% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 33 | 4.2% | 671 | 8 | 7.6% | 163 | 1 | 7.1% | 20 | 42 | 4.6% | 854 |
| Not Hispanic | 567 | 72.0% | 361 | 70 | 66.7% | 45 | 13 | 92.9% | 8 | 650 | 71.7% | 414 |
| Unknown | 187 | 23.8% | - | 27 | 25.7% | - | 0 | 0.0% | - | 214 | 23.6% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 10-14 | 3 | 0.4% | 30 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 3 | 0.3% | 30 |
| 15-19 | 215 | 27.3% | 1,892 | 29 | 27.6% | 255 | 0 | 0.0% | 0 | 244 | 26.9% | 2,147 |
| 20-24 | 330 | 41.9% | 2,554 | 31 | 29.5% | 240 | 3 | 21.4% | 23 | 364 | 40.2% | 2,817 |
| 25-29 | 127 | 16.1% | 1,035 | 22 | 21.0% | 179 | 3 | 21.4% | 24 | 152 | 16.8% | 1,239 |
| 30-34 | 57 | 7.2% | 553 | 12 | 11.4% | 116 | 1 | 7.1% | 10 | 70 | 7.7% | 679 |
| 35-39 | 35 | 4.4% | 325 | 8 | 7.6% | 74 | 3 | 21.4% | 28 | 46 | 5.1% | 427 |
| 40-44 | 9 | 1.1% | 76 | 3 | 2.9% | 25 | 2 | 14.3% | 17 | 14 | 1.5% | 118 |
| 45-49 | 4 | 0.5% | 31 | 0 | 0.0% | 0 | 1 | 7.1% | 8 | 5 | 0.6% | 39 |
| 50+ | 7 | 0.9% | 14 | 0 | 0.0% | 0 | 1 | 7.1% | 2 | 8 | 0.9% | 16 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 787 | 100.0% | 485 | 105 | 100.0% | 65 | 14 | 100.0% | 9 | 906 | 100.0% | 559 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
 Division of Public Health
 Bureau of Communicable Diseases
 STD Control Section
 P-00415B (07/2017)