

Wisconsin sexually transmitted disease surveillance report, 2017

Oneida County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 73 | 100.0% | 203 | 2 | 100.0% | 6 | 0 | - | 0 | 75 | 100.0% | 208 |
| Sex | | | | | | | | | | | | |
| Male | 19 | 26.0% | 106 | - | - | - | - | - | - | 20 | 26.7% | 111 |
| Female | 54 | 74.0% | 299 | - | - | - | - | - | - | 55 | 73.3% | 305 |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 68 | 93.2% | 193 | - | - | - | - | - | - | 70 | 93.3% | 199 |
| African American | 1 | 1.4% | 370 | - | - | - | - | - | - | 1 | 1.3% | 370 |
| American Indian | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| Other | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Unknown | 4 | 5.5% | - | - | - | - | - | - | - | 4 | 5.3% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 1 | 1.4% | 258 | - | - | - | - | - | - | 1 | 1.3% | 258 |
| Not Hispanic | 66 | 90.4% | 185 | - | - | - | - | - | - | 68 | 90.7% | 191 |
| Unknown | 6 | 8.2% | - | - | - | - | - | - | - | 6 | 8.0% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 10-14 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 15-19 | 17 | 23.3% | 795 | - | - | - | - | - | - | 17 | 22.7% | 795 |
| 20-24 | 37 | 50.7% | 1,857 | - | - | - | - | - | - | 37 | 49.3% | 1,857 |
| 25-29 | 10 | 13.7% | 494 | - | - | - | - | - | - | 11 | 14.7% | 543 |
| 30-34 | 3 | 4.1% | 188 | - | - | - | - | - | - | 3 | 4.0% | 188 |
| 35-39 | 3 | 4.1% | 149 | - | - | - | - | - | - | 3 | 4.0% | 149 |
| 40-44 | 2 | 2.7% | 78 | - | - | - | - | - | - | 2 | 2.7% | 78 |
| 45-49 | 0 | 0.0% | 0 | - | - | - | - | - | - | 0 | 0.0% | 0 |
| 50+ | 1 | 1.4% | 6 | - | - | - | - | - | - | 2 | 2.7% | 13 |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Northern | 73 | 100.0% | 203 | - | - | - | - | - | - | 75 | 100.0% | 208 |
| Southeastern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | - | - | - | - | - | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population

