

Wisconsin sexually transmitted disease surveillance report, 2017

Saint Croix County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 209 | 100.0% | 253 | 16 | 100.0% | 19 | 8 | 100.0% | 10 | 233 | 100.0% | 282 |
| Sex | | | | | | | | | | | | |
| Male | 63 | 30.1% | 152 | 6 | 37.5% | 14 | 6 | 75.0% | 14 | 75 | 32.2% | 181 |
| Female | 146 | 69.9% | 357 | 10 | 62.5% | 24 | 2 | 25.0% | 5 | 158 | 67.8% | 386 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 179 | 85.6% | 223 | 14 | 87.5% | 17 | 8 | 100.0% | 10 | 201 | 86.3% | 250 |
| African American | 6 | 2.9% | 881 | 1 | 6.3% | 147 | 0 | 0.0% | 0 | 7 | 3.0% | 1,028 |
| American Indian | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| Asian/Pacific Islander | 1 | 0.5% | 85 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 0.4% | 85 |
| Other | 6 | 2.9% | - | 0 | 0.0% | - | 0 | 0.0% | - | 6 | 2.6% | - |
| Unknown | 17 | 8.1% | - | 1 | 6.3% | - | 0 | 0.0% | - | 18 | 7.7% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 13 | 6.2% | 1,144 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 13 | 5.6% | 1,144 |
| Not Hispanic | 178 | 85.2% | 219 | 15 | 93.8% | 18 | 8 | 100.0% | 10 | 201 | 86.3% | 247 |
| Unknown | 18 | 8.6% | - | 1 | 6.3% | - | 0 | 0.0% | - | 19 | 8.2% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 10-14 | 1 | 0.5% | 18 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 0.4% | 18 |
| 15-19 | 59 | 28.2% | 1,081 | 3 | 18.8% | 55 | 1 | 12.5% | 18 | 63 | 27.0% | 1,154 |
| 20-24 | 91 | 43.5% | 1,835 | 5 | 31.3% | 101 | 1 | 12.5% | 20 | 97 | 41.6% | 1,956 |
| 25-29 | 35 | 16.7% | 549 | 3 | 18.8% | 47 | 1 | 12.5% | 16 | 39 | 16.7% | 612 |
| 30-34 | 14 | 6.7% | 230 | 1 | 6.3% | 16 | 2 | 25.0% | 33 | 17 | 7.3% | 279 |
| 35-39 | 5 | 2.4% | 83 | 2 | 12.5% | 33 | 3 | 37.5% | 50 | 10 | 4.3% | 166 |
| 40-44 | 2 | 1.0% | 32 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 2 | 0.9% | 32 |
| 45-49 | 0 | 0.0% | 0 | 1 | 6.3% | 14 | 0 | 0.0% | 0 | 1 | 0.4% | 14 |
| 50+ | 2 | 1.0% | 9 | 1 | 6.3% | 4 | 0 | 0.0% | 0 | 3 | 1.3% | 13 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 209 | 100.0% | 253 | 16 | 100.0% | 19 | 8 | 100.0% | 10 | 233 | 100.0% | 282 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population

