

Wisconsin sexually transmitted disease surveillance report, 2017

Winnebago County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 661 | 100.0% | 408 | 104 | 100.0% | 64 | 12 | 100.0% | 7 | 777 | 100.0% | 479 |
| Sex | | | | | | | | | | | | |
| Male | 200 | 30.3% | 245 | 46 | 44.2% | 56 | 12 | 100.0% | 15 | 258 | 33.2% | 317 |
| Female | 459 | 69.4% | 569 | 57 | 54.8% | 71 | 0 | 0.0% | 0 | 516 | 66.4% | 640 |
| Unknown | 2 | 0.3% | - | 1 | 1.0% | - | 0 | 0.0% | - | 3 | 0.4% | - |
| Race | | | | | | | | | | | | |
| White | 530 | 80.2% | 342 | 63 | 60.6% | 41 | 10 | 83.3% | 6 | 603 | 77.6% | 389 |
| African American | 76 | 11.5% | 2,686 | 34 | 32.7% | 1,202 | 2 | 16.7% | 71 | 112 | 14.4% | 3,959 |
| American Indian | 4 | 0.6% | 403 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 4 | 0.5% | 403 |
| Asian/Pacific Islander | 11 | 1.7% | 338 | 1 | 1.0% | 31 | 0 | 0.0% | 0 | 12 | 1.5% | 369 |
| Other | 20 | 3.0% | - | 3 | 2.9% | - | 0 | 0.0% | - | 23 | 3.0% | - |
| Unknown | 20 | 3.0% | - | 3 | 2.9% | - | 0 | 0.0% | - | 23 | 3.0% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 42 | 6.4% | 854 | 2 | 1.9% | 41 | 0 | 0.0% | 0 | 44 | 5.7% | 895 |
| Not Hispanic | 551 | 83.4% | 351 | 96 | 92.3% | 61 | 12 | 100.0% | 8 | 659 | 84.8% | 419 |
| Unknown | 68 | 10.3% | - | 6 | 5.8% | - | 0 | 0.0% | - | 74 | 9.5% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 10-14 | 5 | 0.8% | 51 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 5 | 0.6% | 51 |
| 15-19 | 143 | 21.6% | 1,258 | 16 | 15.4% | 141 | 0 | 0.0% | 0 | 159 | 20.5% | 1,399 |
| 20-24 | 311 | 47.0% | 2,407 | 37 | 35.6% | 286 | 0 | 0.0% | 0 | 348 | 44.8% | 2,693 |
| 25-29 | 104 | 15.7% | 848 | 21 | 20.2% | 171 | 3 | 25.0% | 24 | 128 | 16.5% | 1,044 |
| 30-34 | 54 | 8.2% | 524 | 8 | 7.7% | 78 | 1 | 8.3% | 10 | 63 | 8.1% | 611 |
| 35-39 | 23 | 3.5% | 214 | 12 | 11.5% | 111 | 1 | 8.3% | 9 | 36 | 4.6% | 334 |
| 40-44 | 8 | 1.2% | 67 | 6 | 5.8% | 51 | 1 | 8.3% | 8 | 15 | 1.9% | 127 |
| 45-49 | 6 | 0.9% | 46 | 1 | 1.0% | 8 | 2 | 16.7% | 15 | 9 | 1.2% | 69 |
| 50+ | 7 | 1.1% | 14 | 3 | 2.9% | 6 | 4 | 33.3% | 8 | 14 | 1.8% | 28 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 661 | 100.0% | 408 | 104 | 100.0% | 64 | 12 | 100.0% | 7 | 777 | 100.0% | 479 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases
STD Control Section
P-00415B (08/2018)

Wisconsin STD Program data