

Wisconsin sexually transmitted disease surveillance report, 2019

Brown County

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total | | | | | | | | | | | | |
| Total | 1,146 | 100.0% | 468 | 245 | 100.0% | 100 | 19 | 100.0% | 8 | 1,410 | 100.0% | 575 |
| Sex | | | | | | | | | | | | |
| Male | 344 | 30.0% | 282 | 124 | 50.6% | 101 | 15 | 78.9% | 12 | 483 | 34.3% | 395 |
| Female | 802 | 70.0% | 653 | 121 | 49.4% | 98 | 4 | 21.1% | 3 | 927 | 65.7% | 755 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Race | | | | | | | | | | | | |
| White | 713 | 62.2% | 314 | 112 | 45.7% | 49 | 14 | 73.7% | 6 | 839 | 59.5% | 370 |
| African American | 204 | 17.8% | 3,625 | 98 | 40.0% | 1,742 | 1 | 5.3% | 18 | 303 | 21.5% | 5,385 |
| American Indian | 87 | 7.6% | 1,319 | 21 | 8.6% | 318 | 0 | 0.0% | 0 | 108 | 7.7% | 1,638 |
| Asian/Pacific Islander | 31 | 2.7% | 524 | 2 | 0.8% | 34 | 0 | 0.0% | 0 | 33 | 2.3% | 558 |
| Other | 84 | 7.3% | - | 6 | 2.4% | - | 1 | 5.3% | - | 91 | 6.5% | - |
| Unknown | 27 | 2.4% | - | 6 | 2.4% | - | 0 | 0.0% | - | 33 | 2.3% | - |
| Hispanic ethnicity | | | | | | | | | | | | |
| Hispanic | 175 | 15.3% | 1,162 | 19 | 7.8% | 126 | 1 | 5.3% | 7 | 195 | 13.8% | 1,295 |
| Not Hispanic | 944 | 82.4% | 411 | 220 | 89.8% | 96 | 2 | 10.5% | 1 | 1,166 | 82.7% | 507 |
| Unknown | 27 | 2.4% | - | 6 | 2.4% | - | 0 | 0.0% | - | 33 | 2.3% | - |
| Age | | | | | | | | | | | | |
| 0-4 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 1 | 5.3% | 6 | 1 | 0.1% | 6 |
| 5-9 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 | 0 | 0.0% | 0 |
| 10-14 | 13 | 1.1% | 80 | 4 | 1.6% | 25 | 0 | 0.0% | 0 | 17 | 1.2% | 105 |
| 15-19 | 324 | 28.3% | 1,857 | 36 | 14.7% | 206 | 1 | 5.3% | 6 | 361 | 25.6% | 2,070 |
| 20-24 | 437 | 38.1% | 2,641 | 78 | 31.8% | 471 | 3 | 15.8% | 18 | 518 | 36.7% | 3,131 |
| 25-29 | 182 | 15.9% | 1,096 | 56 | 22.9% | 337 | 7 | 36.8% | 42 | 245 | 17.4% | 1,476 |
| 30-34 | 90 | 7.9% | 538 | 29 | 11.8% | 173 | 1 | 5.3% | 6 | 120 | 8.5% | 718 |
| 35-39 | 60 | 5.2% | 356 | 18 | 7.3% | 107 | 3 | 15.8% | 18 | 81 | 5.7% | 481 |
| 40-44 | 25 | 2.2% | 133 | 10 | 4.1% | 53 | 1 | 5.3% | 5 | 36 | 2.6% | 192 |
| 45-49 | 7 | 0.6% | 35 | 7 | 2.9% | 35 | 0 | 0.0% | 0 | 14 | 1.0% | 70 |
| 50+ | 8 | 0.7% | 11 | 7 | 2.9% | 10 | 2 | 10.5% | 3 | 17 | 1.2% | 23 |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | 1,146 | 100.0% | 468 | 245 | 100.0% | 100 | 19 | 100.0% | 8 | 1,410 | 100.0% | 575 |
| Northern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southeastern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Southern | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Western | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |
| Unknown | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - | 0 | 0.0% | - |

* Demographic breakdown not available for diseases with less than 5 cases reported.

** Cases per 100,000 population



DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases
STD Control Section
P-00415B (08/2020)