

Wisconsin Sexually Transmitted Infections Surveillance Report
FLORENCE COUNTY, 2021

| | Chlamydia* | | | Gonorrhea* | | | Syphilis* | | | Total* | | |
|----------------------------|------------|---------|--------|------------|---------|--------|-----------|---------|--------|--------|---------|--------|
| | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** | Cases | Percent | Rate** |
| Total cases | 2 | 100.0% | 45 | 1 | 100.0% | 23 | 0 | - | - | 3 | 100.0% | 68 |
| Gender | | | | | | | | | | | | |
| Male | - | - | - | - | - | - | - | - | - | - | - | - |
| Female | - | - | - | - | - | - | - | - | - | - | - | - |
| Transgender Male | - | - | - | - | - | - | - | - | - | - | - | - |
| Transgender Female | - | - | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Race/ethnicity*** | | | | | | | | | | | | |
| White | - | - | - | - | - | - | - | - | - | - | - | - |
| Black | - | - | - | - | - | - | - | - | - | - | - | - |
| Hispanic | - | - | - | - | - | - | - | - | - | - | - | - |
| Native American | - | - | - | - | - | - | - | - | - | - | - | - |
| Asian/Pacific Islander | - | - | - | - | - | - | - | - | - | - | - | - |
| Other | - | - | - | - | - | - | - | - | - | - | - | - |
| Unknown | - | - | - | - | - | - | - | - | - | - | - | - |
| Age | | | | | | | | | | | | |
| 0-4 | - | - | - | - | - | - | - | - | - | - | - | - |
| 5-9 | - | - | - | - | - | - | - | - | - | - | - | - |
| 10-14 | - | - | - | - | - | - | - | - | - | - | - | - |
| 15-19 | - | - | - | - | - | - | - | - | - | - | - | - |
| 20-24 | - | - | - | - | - | - | - | - | - | - | - | - |
| 25-29 | - | - | - | - | - | - | - | - | - | - | - | - |
| 30-34 | - | - | - | - | - | - | - | - | - | - | - | - |
| 35-39 | - | - | - | - | - | - | - | - | - | - | - | - |
| 40-44 | - | - | - | - | - | - | - | - | - | - | - | - |
| 45-49 | - | - | - | - | - | - | - | - | - | - | - | - |
| 50+ | - | - | - | - | - | - | - | - | - | - | - | - |
| Region of residence | | | | | | | | | | | | |
| Northeastern | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE |
| Northern | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE |
| Southeastern | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE |
| Southern | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE |
| Western | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE | FALSE |

*Demographic breakdown not available for diseases with less than 5 cases reported.

**Cases per 100,000 population.

*** Race/ethnicity breakdowns are not shown if the cell in the subgroup has either a value of <5 or an estimated population size (denominator) <100.

