

	Syph./HIV co-infect	All syphilis	Co_infect/ all syph.
Sex	<u>Cases</u>	<u>Cases</u>	<u>Percent</u>
Male	100	244	41.0%
Female	0	33	0.0%
Transgender	<u>1</u>	<u>2</u>	50.0%
TOTAL	101	<u>2</u>	<b>36.2%</b>
IOIAL	101	2,3	30.270
Age			
10-14	0	1	0.0%
15-19	1	12	8.3%
20-24	22	70	31.4%
25-29	24	59	40.7%
30-34	11	40	27.5%
35-39	9	28	32.1%
40-44	10	17	58.8%
45-49	14	21	66.7%
50+	<u>10</u>	<u>31</u>	32.3%
TOTAL	101	279	36.2%
Race/Ethnicity			
White	32	97	33.0%
Afr. Amer.	61	136	44.9%
Hispanic	7	35	20.0%
Amer. Indian	0	3	0.0%
Asian/Pac. Is.	0	5	0.0%
Other (Mult.)	1	2	50.0%
Unknown	<u>0</u>	<u>1</u>	0.0%
TOTAL	101	279	36.2%
Pagion of			
Region of Residence			
Northeastern	6	19	31.6%
Northern	0	19	0.0%
Southeastern	82	191	42.9%
Southern	12	49	42.9% 24.5%
Western	1	<u>19</u>	5.3%
TOTAL	101	1 <u>9</u> 279	36.2%
· OIAL	101	_, ,	30.2/0

<sup>\*</sup>All syphilis cases are diagnosed for the current year of the report; however, the co-infected cases are those with a new or pre-existing HIV diagnosis.

## Syphilis Stages Among Syphilis/HIV Co-infected cases Syph./HIV ΑII Co\_infect/ co-infect syphilis all syph. Cases Cases Percent Primary syph. 4 22 18.2% Secondary syph. 21 60 35.0% Early latent syph. 41 96 42.7% Late syphilis 35 101 34.7% **TOTAL** 101 279 36.2%





	Syphilis		Syphilis/HIV	
		Percent		Percent
	Cases	of Total	Cases	of Total
Men who have				
sex w/men (MSM)	<u>183</u>	65.6%	<u>101</u>	100%
TOTAL	279		101	



DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases
STD Control Section
P-01097(08/2015)

<sup>•</sup> The source of the information for this report was the STD Control Section Performance Measure Database