

Reported Syphilis Cases With and Without HIV Co-infection,* Wisconsin 2015

Sex	Syphilis/ HIV Co-infected Cases	All Syphilis Cases	Co-infected/ All Syphilis Percent
Male	92	241	38.2%
Female	2	28	7.1%
Transgender	1	1	100.0%
TOTAL	95	270	35.2%
TOTAL	33	270	33.270
Age			
10-14	0	0	0.0%
15-19	2	11	18.2%
20-24	13	50	26.0%
25-29	22	56	39.9%
30-34	14	37	37.8%
35-39	9	33	27.3%
40-44	9	27	33.3%
45-49	7	17	41.2%
50+	19	39	48.7%
TOTAL	95	270	35.2%
Race/ Ethnicity			
White	41	127	32.3%
Afr. Amer.	39	106	36.8%
Hispanic	10	28	35.7%
Amer. Indian	1	1	100.0%
Asian/Pac. Is.	1	4	25.0%
Other (Mult.)	2	3	66.7%
Unknown	1	1	100.0%
TOTAL	95	270	35.2%
Region of Residence			
Northeastern	6	29	20.7%
Northern	2	3	66.7%
Southeastern	71	183	38.8%
Southern	10	37	27.0%

Syphilis Stages Among Syphilis/HIV Co-infected cases								
	Syphilis/HIV Co-infected	All Syphilis	Co-infected/ All Syphilis					
	Cases	Cases	Percent					
Primary syphilis	3	21	14.3%					
Secondary syphilis	28	61	45.9%					
Early latent syphilis	47	103	45.4%					
Late syphilis	17	85	20.0%					
TOTAL	95	270	35.1%					





	All Syphilis		Syphilis/HIV	
	Cases	Percent of Total	Cases	Percent of Total
Men who have sex with men (MSM)	185	68.5%	84	88.4%
TOTAL	270		95	

6

95

Western

TOTAL

WISCONSIN DEPARTMENT OF HEALTH SERVICES
Division of Public Health
Bureau of Communicable Diseases
STD Control Section
P- (03/2016)

Note: The source of the information for this report was the STD Control Section Performance Measure Database.

33.3%

35.2%

18

270

^{*}All syphilis cases are diagnosed for the current report year; however, co-infected cases are those with a new or pre-existing HIV diagnosis.