

## Reported Syphilis Cases With and Without HIV Co-infection,\* Wisconsin 2016

Sex	Syphilis/HIV Co-infected Cases	All Syphilis Cases	Co-infected/ All Syphilis Percent 38.4%	
Male	141	367		
Female	1	56	1.8%	
Transgender	2	4	50.0%	
TOTAL	144	427	33.7%	
Age				
0-14	0	2	0.0%	
15-19	2	18	11.1%	
20-24	21	90	23.3%	
25-29	33	87	37.9%	
30-34	23	54	42.6%	
35-39	18	41	43.9%	
40-44	14	35	40.0%	
45-49	16	34	47.1%	
50+	18	66	27.3%	
TOTAL	144	427	33.7%	
Page (Ethnicity				
<b>Race/Ethnicity</b> White	66	198	33.3%	
Afr. Amer.	66	190	36.3%	
Hispanic Amer.	11	36	30.6%	
Indian Asian/	0	1	0.0%	
Pac. Is. Other	0	7	0.0%	
(Mult.)	0	, 1	0.0%	
Unknown	1	2	50.0%	
TOTAL	144	427	33.7%	
Region of				
Residence		- <b>-</b>	24 604	
Northeastern	14	57	24.6%	
Northern	1	9	11.1%	
Southeastern	100 21	253	39.5%	
Southern Western	21	81 27	25.9% 29.6%	
vvestelli	ŏ	۷۲	29.0%	

## Syphilis Stages Among Syphilis/HIV Co-infected cases Syphilis/HIV All Co-infected/ **Co-infected Syphilis All Syphilis** Cases Cases Percent imary syphilis 13 49 26.5% condary syphilis 33 84 39.3% rly latent syphilis 68 154 44.2% ite syphilis 30 139 21.6% OTAL 33.7% 144 427





	All Syphilis		Syphilis/HIV	
	<b>C</b>	Percent	<b>C</b>	Percent
Men who have sex	Cases	of Total	Cases	of Total
with men (MSM)	289	67.7%	137	95.1%
TOTAL	427		144	



WISCONSIN DEPARTMENT OF HEALTH SERVICES Division of Public Health Bureau of Communicable Diseases STD Control Section P-01097(05/2017)

\*All syphilis cases are diagnosed for the current report year; however, co-infected cases are those with a new or pre-existing HIV diagnosis.

Note: The source of the information for this report was the STD Control Section Performance Measure Database.