

PATIENT RED FLAG CHECKLIST



Be the trusted
resource patients
need you to be.

Know the red flags
to watch for when
dispensing opioids.



Wisconsin Pharmacy
Examining Board



Wisconsin
Department of Health Services

PATIENT RED FLAG CHECKLIST



Be the trusted
resource patients
need you to be.

Know the red flags
to watch for when
dispensing opioids.



Wisconsin Pharmacy
Examining Board



Wisconsin
Department of Health Services

PATIENT RED FLAG CHECKLIST



Be the trusted
resource patients
need you to be.

Know the red flags
to watch for when
dispensing opioids.



Wisconsin Pharmacy
Examining Board



Wisconsin
Department of Health Services



Red Flags

Are you about to fill a prescription for opioids?

Pay special attention to whether any of the following apply to those receiving this prescription:

- Known respiratory disorder, e.g., COPD, asthma
- History of smoking
- Renal dysfunction or hepatic disease
- Prescription for:
 - Benzodiazepine
 - SSRI or TCA antidepressant
 - High dose of opioid (higher than 90 mg morphine daily)
- Receiving methadone, Suboxone[®] or Vivitrol[®]
- Never been prescribed opioids

If any of these apply, this patient may also need naloxone.

Red Flags

Are you about to fill a prescription for opioids?

Pay special attention to whether any of the following apply to those receiving this prescription:

- Known respiratory disorder, e.g., COPD, asthma
- History of smoking
- Renal dysfunction or hepatic disease
- Prescription for:
 - Benzodiazepine
 - SSRI or TCA antidepressant
 - High dose of opioid (higher than 90 mg morphine daily)
- Receiving methadone, Suboxone[®] or Vivitrol[®]
- Never been prescribed opioids

If any of these apply, this patient may also need naloxone.

Red Flags

Are you about to fill a prescription for opioids?

Pay special attention to whether any of the following apply to those receiving this prescription:

- Known respiratory disorder, e.g., COPD, asthma
- History of smoking
- Renal dysfunction or hepatic disease
- Prescription for:
 - Benzodiazepine
 - SSRI or TCA antidepressant
 - High dose of opioid (higher than 90 mg morphine daily)
- Receiving methadone, Suboxone[®] or Vivitrol[®]
- Never been prescribed opioids

If any of these apply, this patient may also need naloxone.