You would do anything to protect your family.



Take steps to protect yourself and those you love against Zika and other diseases.



IF YOU OR YOUR PARTNER ARE PREGNANT OR PLANNING TO BECOME PREGNANT, DO NOT TRAVEL TO AN AREA WHERE ZIKA IS SPREADING.



ZIKA CAN BE PASSED THROUGH SEX.

You can spread Zika to your sexual partners. Sex includes vaginal, anal, and oral sex, as well as the sharing of sex toys.



ZIKA MAY BE IN PLACES YOU DON'T EXPECT. CHECK IF THE AREA YOU ARE TRAVELING TO IS AN AREA WHERE ZIKA IS SPREADING:



USE A CONDOM. If you or your sexual partner lives in or travels to an area with Zika, you should use condoms from start to finish every time you have sex during pregnancy.

www.cdc.gov/zika/geo/



ZIKA CAN CAUSE BIRTH DEFECTS.

Zika can be passed from a pregnant woman to her baby anytime during pregnancy. Approximately 1 in 12 Zika-infected moms have babies with birth defects.



ZIKA CAN BE SPREAD EVEN IF SYMPTOMS ARE NOT PRESENT.

80 percent of people with Zika have no symptoms. Zika may be spread for weeks to months after infection.

