




WISCONSIN DEPARTMENT
of HEALTH SERVICES



RESPIRATORY VIRUS SURVEILLANCE REPORT

Week 3, Ending January 22, 2022


Wisconsin Department of Health Services | Division of Public Health
Bureau of Communicable Diseases | Communicable Diseases Epidemiology Section

www.dhs.wisconsin.gov/dph/bcd.htm | dhsdphbcd@dhs.wi.gov




INFLUENZA LIKE ILLNESS (ILI) ACTIVITY


STATE OF WISCONSIN



REGION V OF US (WI, MN, IL, MI, OH, IN)



UNITED STATES



● ILI: HIGH LEVELS ● ILI: MODERATE LEVELS ● ILI: BELOW BASELINE ● ILI: INSUFFICIENT DATA

AT-A-GLANCE:

■ Predominant Viruses of the Week:

Seasonal Coronaviruses are the predominant viruses this week.

Current Alerts:

Influenza-like illness activity has been decreasing for the previous 5 weeks.

Additional data on SARS-CoV-2 (the virus causing COVID-19) trends in Wisconsin can be found at:

<https://www.dhs.wisconsin.gov/covid-19/data.htm>

INFLUENZA-ASSOCIATED PEDIATRIC DEATHS REPORTED:

	Week 3, 2022	October 1, 2021 to present
Wisconsin	0	0
Nationwide	0	5


For National US influenza surveillance statistics visit: www.cdc.gov/flu/weekly/



INFLUENZA AND PNEUMONIA-ASSOCIATED MORTALITY

Influenza and Pneumonia Deaths, Wisconsin

Influenza- and pneumonia-associated deaths by influenza season year and week, Wisconsin



Influenza- and pneumonia-associated deaths by most recent 3 week period.


Influenza season week	Influenza-associated deaths (I)	Pneumonia-associated deaths (P)	Percent I+P of all deaths	Baseline I+P of all deaths	Threshold I+P of all deaths
1	2	177	11.5%	5.4%	6.2%
2	1	176	13.8%	5.5%	6.3%
3 Preliminary Data	4	153	14.0%	5.6%	6.4%

Data source: [DPH, Office of Health Informatics](#)



WISCONSIN LABORATORY SURVEILLANCE FOR RESPIRATORY VIRUSES BY PCR

Wisconsin positive influenza results and subtypes by PCR



Cumulative number of positive influenza PCR tests by subtype


October 9, 2021 to present

	A (2009 H1N1)	Influenza A: A (H3N2)	98% A (Unknown)	B (Victoria)	Influenza B: B (Yamagata)	2% B (Unknown)	Total
Total positive (n)	12	769	4,331	1	0	90	5,203
% of total positive	0%	15%	83%	0%	0%	2%	100%



WISCONSIN LABORATORY SURVEILLANCE FOR RESPIRATORY VIRUSES BY PCR

Trends in respiratory virus activity by PCR



Week 3 Ending on January 22, 2022

Respiratory virus	Tested	Positive (n)	Positive (%)	Influenza A			Influenza B		
				H3N2	2009	Unknown	Victoria	Yamagata	Unknown
Influenza	13355	330	2.5%	26	0	296	0	0	8

Respiratory virus	Tested	Positive (n)	Positive (%)	Parainfluenza 1	Parainfluenza 2	Parainfluenza 3	Parainfluenza 4
Parainfluenza	1038	15	1.4%	0	4	10	1


Respiratory virus	Tested	Positive (n)	Positive (%)	CoV 229E	CoV OC43	CoV NL63	CoV HKU1
Coronavirus (seasonal)	211	10	4.7%	2	8	0	0

Respiratory virus	Tested	Positive (n)	Positive (%)
RSV	5309	122	2.3%
Human metapneumovirus	1043	50	4.8%
Rhino-enterovirus	1009	65	6.4%
Adenovirus	211	5	2.4%



WISCONSIN STATE SUMMARY

ILI activity trend analysis by influenza season, Wisconsin




Influenza-associated hospitalizations, Wisconsin Electronic Disease Surveillance System

October 1, 2021 to present

Age group (years)	Total reported (n)	Influenza subtype					Admitted to ICU	Required mechanical ventilation	Pregnant	Postpartum (≤6 weeks)
		A (2009 H1N1)	A (H3N2)	A (Unknown)	B	Not reported				
<1	3	0	0	3	0	0	0	0		
1-4	8	1	0	6	1	0	4	0		
5-17	11	1	0	10	0	0	2	1		
18-49	42	0	4	37	1	0	6	0	5	0
50-64	25	0	1	22	2	0	4	0		
65+	125	0	5	108	12	0	5	2		
Total	214	2	10	186	16	0	21	3	5	0

Reported cumulative influenza-associated hospitalizations by influenza season, Wisconsin




* The 2020 - 2021 influenza season was unusually low due much in part to the ongoing COVID-19 pandemic. As such, numbers for that season are substantially different than previous seasons and should be considered an anomaly.




ILI ACTIVITY TREND ANALYSIS


Wisconsin



Northeastern Region




Northern Region




ILI ACTIVITY TREND ANALYSIS


Southeastern Region



Southern Region




Western Region



SEASONAL INFLUENZA VACCINATION

Cumulative percentage of Wisconsin residents who received 1 or more doses of influenza vaccine, 2021-2022 influenza season



Data source: All influenza vaccination rates presented were calculated using data from the Wisconsin Immunization Registry (numerator) and Wisconsin population estimates (denominator).

Influenza vaccine composition 2021-2022:

Egg-based vaccines are recommended to contain:

- A/Victoria/2570/2019 (H1N1) pdm09-like virus
- A/Cambodia/e0826360/2020 (H3N2)-like virus
- B/Washington/02/2019- like virus (B/Victoria lineage)
- B/Phuket/3073/2013-like virus (B/Yamagata lineage)


Cell- or recombinant-based vaccines are recommended to contain:

- A/Wisconsin/588/2019 (H1N1) pdm09-like virus
- A/Cambodia/e0826360/2020 (H3N2)-like virus
- B/Washington/02/2019- like virus (B/Victoria lineage)
- B/Phuket/3073/2013-like virus (B/Yamagata lineage)



SEASONAL INFLUENZA VACCINATION

Percentage of Wisconsin residents who received one or more doses of influenza vaccine, by age group and influenza season



Each season includes doses administered during the same time period (August 1 through May 3).

Percentage of Wisconsin residents who received one or more doses of influenza vaccine, by race and ethnicity and region, 2021-2022 influenza season

