

Wisconsin EMS Protocols

FIELD VERSION 1
JANUARY 2021

The complete "[Wisconsin Protocols for Emergency Medical Services](#)" is also available on the Internet. Protocols are occasionally amended during the year; notice and publication will be communicated through DHS. Please check the DHS website to be sure you have the most up-to-date version. The edition date appears on the lower portion of the page

P-02875A (06/2021)

Contents

Introduction	5
Purpose and Notes	5
Acknowledgements	6
Universal Care Guideline	7
Functional Needs	14
Patient Refusals	17
Cardiovascular	19
Adult and Pediatric Syncope and Presyncope	19
ACUTE CORONARY SYNDROME (ACS) OR ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION (STEMI)	22
Bradycardia	24
Implantable Ventricular Assist Devices	29
Tachycardia with a Pulse	32
Suspected Stroke/Transient Ischemic Attack	37
General Medical	39
Abdominal Pain	39
Abuse and Maltreatment	42
Agitated or Violent Patient/Behavioral Emergency	46
Anaphylaxis and Allergic Reaction	50
Altered Mental Status	53
Back Pain	55
End-of-Life Care/Palliative Care	57
Hyperglycemia	59
Hypoglycemia	61
Nausea-Vomiting	64
Pain Management	66
Seizures	73
Shock	76
Sickle Cell Pain Crisis	79
Resuscitation	81
Cardiac Arrest: VF, VT, Asystole, and PEA	81
Adult Post-ROSC (Return of Spontaneous Circulation) Care	87
Determination of Death/Withholding Resuscitative Efforts	89
Do Not Resuscitate Status, Advance Directives, and Health Care Power of Attorney (POA) Status	91
Termination of Resuscitative Efforts	93
Pediatric-Specific Guidelines	97
Brief Resolved Unexplained Event (BRUE)	97
Pediatric Respiratory Distress (Bronchiolitis)	100
Pediatric Respiratory Distress (Croup)	103
Neonatal Resuscitation	106

Top 10 Take-Home Messages for Neonatal Life Support	108
Pediatric Resuscitation	111
OB/GYN	113
Childbirth	113
Eclampsia/Pre-Eclampsia	117
Obstetrical and Gynecological Conditions	120
Respiratory	122
Airway Management	122
Bronchospasm (due to Asthma and Obstructive Lung Disease)	127
Pulmonary Edema	130
Trauma	133
General Trauma Management	133
Blast Injuries	138
Burns	141
Crush Injury	144
Extremity Trauma and External Hemorrhage Management	146
Facial and/or Dental Trauma	150
Head Injury	152
High Threat Considerations, Active Shooter Scenario	155
Spinal Care	157
Toxins and Environmental	160
Poisoning or Overdose Universal Care	160
Acetylcholinesterase Inhibitors (Carbamates, Nerve Agents, Organophosphates) Exposure	167
Radiation Exposure	180
Topical Chemical Burn	183
Stimulant Poisoning or Overdose	187
Cyanide Exposure	190
Beta Blocker Poisoning or Overdose	193
Bites and Envenomation	196
Calcium Channel Blocker Poisoning or Overdose	198
Carbon Monoxide or Smoke Inhalation	201
Opioid Poisoning or Overdose	204
Airway Respiratory Irritants	208
Riot Control Agents	215
Hyperthermia and Heat Exposure	217
Hypothermia and Cold Exposure	221
Drowning	227
Dive (SCUBA) Injury or Accidents	230
Conducted Electrical Weapon Injury (e.g. TASER®)	232
Electrical Injuries	234
Lightning or Lightning Strike Injury	237
APPENDICES	240
I. Universal Documentation Guideline	240
II. 2021 Scope of Practice Medications	254

III. Medications.....	259
IV. Approved Abbreviations.....	276
V. Burn Body Surface Area.....	280
VI. Neurologic Status Assessment.....	283
VII. Abnormal Vital Signs	284
VIII. Evidence-Based Guidelines: GRADE Methodology.....	285
IX. 2011 Guidelines for Field Triage of Injured Patients.....	286

Introduction

The Wisconsin Model EMS Clinical Guidelines are not mandatory. The focus of these guidelines is primarily clinical; operational decisions must be determined at the agency level as community resources vary considerably throughout the state. These guidelines were adapted from the National Association of State EMS Officials' [National Model Clinical Guidelines](#) and are aligned with the metrics developed through the EMS Compass Project and the National EMS Quality Alliance.

Purpose and Notes

The Wisconsin Model EMS Clinical Guidelines are intended to help Emergency Medical Services (EMS) systems ensure a more standardized approach to the practice of patient care and are to be used in conjunction with the DHS-approved Wisconsin [EMS Scope of Practice](#). In the event of error in this clinical guideline document, the Wisconsin EMS Scope of Practice is the arbiter and defines scope of care at the EMS Provider (Agency) and Professional (Individual) levels.

These clinical guidelines will be maintained by the Wisconsin Department of Health Services, Division of Public Health, Office of Preparedness and Emergency Health, EMS Section and will be reviewed and updated annually with input from the Physician Advisory Committee and the Wisconsin EMS Board among other partners and stakeholders.

Key:

EMS provider skills, equipment or medications, based on the DHS-approved Wisconsin EMS Scope of Practice and authorized for use by the service EMS medical director, are located in the **Patient Management** and **Notes and Education Pearls** sections and highlighted in an *[italicized bracket]* followed by an "R" or "O" to designate when the intervention is a required or optional skill for the EMS provider practice level.

For example, if the clinical guideline directs oxygen delivery for shortness of breath, it will be followed by *[EMR-O; EMT-R]* which directs that oxygen is an optional skill for the EMR practice level but a required skill for EMT and higher. If the clinical guideline directs splinting with a pelvic compression device for suspected pelvic fracture, it will be followed by *[EMR-O; PARA-R]* which directs that a pelvic compression device is an optional skill for EMR through INT practice levels, but required at the paramedic level. Finally, if the clinical guideline directs isopropyl alcohol for nausea, it will be followed by *[ALL EMS SERVICE PROVIDERS-O]* since isopropyl alcohol is an optional intervention for all service levels.

Universal Care and Poisoning/Overdose Universal Care guidelines are included to reduce the need for extensive reiteration of basic assessment and other considerations in every guideline.

The appendices contain material such as neurologic status assessment and burn assessment tools, to which many guidelines refer, to increase consistency in internal standardization and to reduce duplication.

While some specific guidelines have been included for **pediatric patients**, considerations of patient age and size (pediatric, geriatric, and bariatric) have been interwoven in the guidelines

throughout the document.

Generic medication names are utilized throughout the guidelines. A complete list of these, along with respective brand names, may be found in [Appendix IV—Medications](#).

[Acknowledgements](#)

The EMS Section recognizes the valuable input and support from the EMS Physician Advisory Committee, the Wisconsin EMS Board, the Wisconsin EMS Technical Colleges systems and EMS Service and Medical Directors.

Universal Care

Universal Care Guideline

Aliases

Patient assessment, patient history, physical assessment, primary survey, secondary survey

Patient Care Goals

Facilitate appropriate initial assessment and management of any EMS patient and link to appropriate specific guidelines as dictated by the findings within the **Universal Care** guideline.

Patient Presentation

Inclusion Criteria

All patient encounters with and care delivery by EMS personnel

Exclusion Criteria

None

Patient Management

Assessment

1. Assess scene safety.
 - a. Evaluate for hazards to EMS personnel, patient, bystanders.
 - b. Ensure appropriately trained and equipped rescuers extricate patient from any dangerous environment (examples include hazardous material, water, combustible, electrical, confined space).
 - c. Determine number of patients.
 - d. Determine mechanism of injury.
 - e. Request additional resources if needed and weigh the benefits of waiting for additional resources against rapid transport to definitive care.
 - f. Consider declaration of mass casualty incident if needed.
2. Use appropriate personal protective equipment (PPE).
3. Wear high-visibility, retro-reflective apparel when deemed appropriate (e.g. operations at night or in darkness, on or near roadways).
4. Consider cervical spine stabilization , spinal motion restriction [*EMR-O; EMT-R*] and/or cervical collar [*EMR-O; EMT-R*] if trauma.
5. Conduct a primary survey.

(Airway, breathing, circulation is cited below; although there are specific circumstances where circulation, airway, breathing may be indicated such as cardiac arrest or major arterial bleeding.)

 - a. Airway (assess for patency and open the airway as indicated)
 - i. Patient is unable to maintain airway patency—open airway
 1. Perform head tilt chin lift **or**
 2. Perform jaw thrust
 3. Perform suction
 4. Consider use of the appropriate airway management adjuncts and devices: oral airway, nasal airway, non-visualized airways (supraglottic or extraglottic) [*EMR-O; EMT-R*], endotracheal

tube *[INT-O; PARA-R]*

Notes:

- For **patients with laryngectomies or tracheostomies**: Remove all objects or clothing that may obstruct the opening of these devices, maintain the flow of prescribed oxygen *[EMR-O; EMT-R]*, and reposition the head and/or neck.
- For **obstructed airway, laryngectomy, or tracheostomy**, go to [Airway Management guideline](#).

b. Breathing

- i. Evaluate rate, breath sounds, accessory muscle use, retractions, patient positioning.
- ii. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.

Note: For apnea (not breathing), go to [Airway Management guideline](#).

c. Circulation

- i. Control any major external bleeding [see [Extremity Trauma/External Hemorrhage Management guideline](#)].
- ii. Assess pulse.
 1. If none, go to [Cardiac Arrest guideline](#).
 2. Assess rate and quality of carotid and radial pulses.
- iii. Evaluate perfusion by assessing skin color and temperature.
 1. Evaluate capillary refill.

d. Disability

- i. Evaluate patient responsiveness: Glasgow Coma Scale or AVPU scale (Alert, Verbal, Pain, Unresponsive).
- ii. Evaluate gross motor and sensory function in all extremities.
- iii. Check blood glucose *[EMR-O; EMT-R]* in patients with altered mental status.
- iv. If acute stroke suspected, go to [Suspected Stroke/Transient Ischemic Attack guideline](#).

e. Expose patient as appropriate to complaint.

- i. Be considerate of patient modesty.
- ii. Keep patient warm.

6. Conduct a secondary survey.

The performance of the secondary survey should not delay transport in critical patients. See also secondary survey specific to individual complaints in other protocols. Secondary surveys should be tailored to patient presentation and chief complaint. The following are suggested considerations for secondary survey assessment:

a. Head

- i. Pupils
- ii. Naso-oropharynx
- iii. Skull and scalp

b. Neck

- i. Jugular venous distension
- ii. Tracheal position
- iii. Spinal tenderness

c. Chest

- i. Retractions
 - ii. Breath sounds
 - iii. Chest wall deformity
 - d. Abdomen and Back
 - i. Flank and abdominal tenderness or bruising
 - ii. Abdominal distension
 - e. Extremities
 - i. Edema
 - ii. Pulses
 - iii. Deformity
 - e. Neurologic
 - i. Mental status and orientation
 - ii. Motor and sensory
- 7. Obtain Baseline Vital Signs (an initial full set of vital signs is required: pulse, blood pressure, respiratory rate, neurologic status assessment).
 - a. Neurologic status assessment [see [Appendix VII](#)] involves establishing a baseline and then trending any change in patient neurologic status.
 - i. Use Glasgow Coma Score (GCS) **or**
 - ii. Use AVPU (**A**lert, **V**erbal, **P**ainful, **U**nresponsive)
 - b. Patients with cardiac or respiratory complaints
 - i. Use pulse oximetry [*EMR-O; EMT-R*]
 - ii. 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*] should be obtained early in patients with cardiac or suspected cardiac complaints
 - iii. Conduct continuous ECG cardiac monitoring [*Acquisition EMT-O; Interpretation INT-R*], if available
 - iv. Consider waveform capnography essential for patients who require invasive airway management [*Required INT-R*] or capnometry [*Required with Non-visualized Airway EMR-R*]
 - c. Patient with altered mental status
 - i. Check blood glucose [*EMR-O; EMT-R*]
 - ii. Consider waveform capnography (essential for patients who require invasive airway management) [*Required INT-R*] or capnometry [*Required with Non-visualized Airway EMR-R*]
 - d. Stable patients should have at least two sets of pertinent vital signs. Ideally, one set should be taken shortly before arrival at receiving facility.
 - e. Critical patients should have pertinent vital signs frequently monitored.
- 8. Obtain OPQRST history:
 - a. **O**nset of symptoms
 - b. **P**rovocation—location; any exacerbating or alleviating factors
 - c. **Q**uality of pain
 - d. **R**adiation of pain
 - e. **S**everity of symptoms—pain scale
 - f. **T**ime of onset and circumstances around onset
- 9. Obtain SAMPLE history:
 - a. **S**ymptoms
 - b. **A**llergies—medication, environmental, and foods
 - c. **M**edications—prescription and over-the-counter; bring containers to ED if possible
 - d. **P**ast medical history

- i. Look for medical alert tags, portable medical records, advance directives.
- ii. Look for medical devices or implants (some common ones may be dialysis shunt, insulin pump, pacemaker, central venous access port, gastric tubes, urinary catheter).
- e. **Last oral intake**
- f. **Events leading up to the 911 call**

Treatment and Interventions

1. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
2. Place appropriate monitoring equipment as dictated by assessment; these may include:
 - a. Continuous pulse oximetry *[EMR-O; EMT-R]*.
 - b. ECG cardiac rhythm monitoring, *[Acquisition EMT-O; Interpretation INT-O/PARA-R]*.
 - c. Waveform capnography *[Interpretation INT-R]*.
 - d. Carbon monoxide assessment *[ALL PRACTICE LEVELS-O]*.
3. Establish vascular access *[AEMT-R]* if indicated or in patients who are at risk for clinical deterioration.
 - a. If IO is to be used for a conscious patient, consider lidocaine with slow push through IO needle *[INT-O; PARA-R]* to mitigate pain from IO medication administration.
4. Monitor pain scale if appropriate.
5. Reassess patient.

Patient Safety Considerations

- **Sirens:** Avoid routine use of lights and sirens.
- **Speeds:** Always limit speeds to a level that is safe for the emergency vehicle being driven and road conditions on which it is being operated, even when lights and sirens are in use.
- **Legal:** Be aware of legal issues and patient rights as they pertain to and impact patient care (e.g. patients with functional needs or children with special health care needs).
- **Age and comorbidities:** Be aware of potential need to adjust management based on patient age and comorbidities, including medication dosages.
- **Pediatric dose:** Do not exceed the maximum adult, weight-based dose of medication in pediatric patients except where specifically stated in a patient care guideline.
- **Medical control:** Contact online medical control when mandated or as needed.
- **Air medical transport:** Consider [air medical transport](#)

Notes and Educational Pearls

Key Considerations

Pediatrics:

- Use a weight-based assessment tool (length-based tape or other system) to estimate patient weight, and guide medication therapy and adjunct choice.
- Use a weight of 40kg and less to define a pediatric patient.
- Consider using the pediatric assessment triangle (appearance, work of breathing, circulation) when first approaching a child to help with

assessment.

Geriatrics:

- Define geriatric patients as those who are 65 years old or more.
- Assess reduced dosage needs. In these patients, as well as all adult patients, reduced medication dosages may apply to patients with renal disease (i.e. on dialysis or a diagnosis of chronic renal insufficiency) or hepatic disease (i.e. severe cirrhosis or end-stage liver disease)

Co-morbidities: Reduced medication dosages may apply to patients with renal disease (i.e. on dialysis or a diagnosis of chronic renal insufficiency) or hepatic disease (i.e. severe cirrhosis or end-stage liver disease).

Vital signs:

- Oxygen
 - Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- Normal vital signs (see chart below)
 - Hypotension is considered a systolic blood pressure less than the lower limit on the chart.
 - Tachycardia is considered a pulse above the upper limit on the chart.
 - Bradycardia is considered a pulse below the lower limit on the chart.
 - Tachypnea is considered a respiratory rate above the upper limit on the chart.
 - Bradypnea is considered a respiratory rate below the lower limit on the chart.
- Hypertension (although abnormal, may be an expected finding in many patients)
 - Document the hypertension, **but**
 - Do not use an intervention unless it is specifically suggested based on the patient's complaint or presentation.
 - Look for the occurrence of symptoms (e.g. chest pain, dyspnea, vision change, headache, focal weakness or change in sensation, altered mental status) in patients with hypertension. These should be considered concerning, and care should be provided appropriate with the patient's complaint or presentation.

Secondary survey: may not be completed if patient has critical primary survey problems

Critical patients:

- Proactive patient management should occur simultaneously with assessment.
- Ideally, one provider should be assigned to exclusively monitor and facilitate patient- focused care.
- Treatment and interventions should be initiated as soon as practical, but should not impede extrication or delay transport to definitive care.

Air medical transport: [Wisconsin Helicopter EMS Utilization Guidelines](#)

Pertinent Assessment Findings

This guideline is too broad to list all possible findings.

- Document all patient interventions and responses.

Normal Vital Signs

Age	Pulse	Respiratory Rate	Systolic BP
Preterm less than 1 kg	120–160	30–60	36–58
Preterm 1 kg	120–160	30–60	42–66
Preterm 2 kg	120–160	30–60	50–72
Newborn	120–160	30–60	60–70
Up to 1 year	100–140	30–60	70–80
1–3 years	100–140	20–40	76–90
4–6 years	80–120	20–30	80–100
7–9 years	80–120	16–24	84–110
10–12 years	60–100	16–20	90–120
13–14 years	60–90	16–20	90–120
15 years or older	60–90	14–20	90–130

Glasgow Coma Scale

ADULT GLASGOW COMA SCALE		PEDIATRIC GLASGOW COMA SCALE	
Eye Opening (4)		Eye Opening (4)	
Spontaneous	4	Spontaneous	4
To speech	3	To speech	3
To pain	2	To pain	2
None	1	None	1
Best Motor Response (6)		Best Motor Response (6)	
Obeys commands	6	Spontaneous movement	6
Localizes pain	5	Withdraws to touch	5
Withdraws from pain	4	Withdraws from pain	4
Abnormal flexion	3	Abnormal flexion	3
Abnormal extension	2	Abnormal extension	2
None	1	None	1
Verbal Response (5)		Verbal Response (5)	
Oriented	5	Coos, babbles	5
Confused	4	Irritable cry	4
Inappropriate	3	Cries to pain	3
Incomprehensible	2	Moans to pain	2
None	1	None	1
Total		Total	

Functional Needs

Aliases

Developmental delay, disabled, handicapped, impaired, mental illness, mental retardation, special needs

Patient Care Goals

To meet and maintain the additional support required for patients with functional needs during the delivery of prehospital care

Patient Presentation

Inclusion Criteria

These are patients who are identified by the World Health Organization's International Classification of Functioning, Disability, and Health that have experienced a decrement in health resulting in some degree of disability. According to the U.S. Department of Health and Human Services, this includes, but is not limited to, individuals with physical, sensory, mental health, and cognitive and/or intellectual disabilities affecting their ability to function independently without assistance

Functional needs may also include cultural impacts on health care such as language barriers and religious, cultural or ethnic customs that may influence how patients understand health concepts, how they take care of their health and how they make decisions related to their health.

Exclusion Criteria

None

Patient Management

Assessment

1. Identify the functional need by means of information from the patient, the patient's family, bystanders, medic alert bracelets or documents, or the patient's adjunct assistance devices.
2. Avoid intentionally abbreviating the physical examination, although the manner in which the exam is performed may need to be modified to accommodate the specific needs of the patient.

Treatment and Interventions

Do not intentionally reduce or abbreviate medical care during the triage, treatment, and transport of patients with functional needs; however, the manner in which the care is provided may need to be modified to accommodate the specific needs of the patient.

Patient Safety Considerations

For patients with communication barriers (language or sensory), it may be desirable to obtain secondary confirmation of pertinent data (e.g. allergies) from the patient's family, interpreters, or written or electronic medical records. The family members can be an excellent source of information and the presence of a family member can have a calming influence on some of these patients.

Notes and Educational Pearls

Key Considerations

- Communication barriers
 - Language barriers
 - Expressive and/or receptive aphasia
 - Nonverbal
 - Fluency in a different language than that of the EMS professional

Note: Examples of tools to overcome language barriers include:

- Transport of an individual who is fluent in the patient's language along with the patient to the hospital.
- Medical translation cards.
- Telephone-accessible services with live language interpreters.
- Methods through which the patient augments his/her communication skills (e.g. eye blinking, nodding) should be noted, utilized as able, and communicated to the receiving facility.
- Electronic applications for translation.

Sensory barriers:

- Visual impairment
- Auditory impairment
- Examples of tools to overcome sensory barriers include:
 - Braille communication card
 - Sign language
 - Lip reading
 - Hearing aids
 - Written communication

Physical barriers:

- Ambulatory impairment (e.g. limb amputation, bariatric)
- Neuromuscular impairment

Cognitive barriers:

- Mental illness
- Developmental challenge or delay

Pertinent Assessment Findings

- Assistance Adjuncts.
Examples of devices that facilitate the activities of daily living for the patient with functional needs include, but are not limited to:
 - Extremity prostheses
 - Hearing aids
 - Magnifiers
 - Tracheostomy speaking valves
 - White or sensory canes
 - Wheelchairs or motorized scooters

- Service Animals

As defined by the American Disabilities Act, "any guide dog, signal dog, or other animal individually trained to do work or perform tasks for the benefit of an individual with a disability, including, but not limited to guiding individuals with impaired vision, alerting individuals with impaired hearing to intruders or sounds, providing minimal protection or rescue work, pulling a wheelchair, or fetching dropped items."

 - a. Service animals are not classified as a pet and should, by law, always be permitted to accompany the patient with the following exceptions:
 - i. A public entity may ask an individual with a disability to remove a service animal from the premises if:
 - 1. The animal is out of control and the animal's handler does not take effective action to control it; or
 - 2. The animal is not housebroken
 - b. Service animals are not required to wear a vest or a leash. It is illegal to make a request for special identification or documentation from the service animal's partner. EMS providers may only ask the patient if the service animal is required because of a disability and the form of assistance the animal has been trained to perform.
 - c. EMS providers are not responsible for the care of the service animal. If the patient is incapacitated and cannot personally care for the service animal, a decision can be made whether or not to transport the animal in this situation.
 - d. Animals that solely provide emotional support, comfort, or companionship do not qualify as service animals

Patient Refusals

Aliases

Against medical advice, refusal of treatment, refusal of transport

Patient Care Goals/Patient Presentation (Overview)

If an individual (or the parent or legal guardian of the individual) refuses care and/or ambulance transport after prehospital providers have been called to the scene, providers should determine the patient's capacity to make decisions.

Patient Management

Assessment

Decision-Making Capacity

- An individual who is alert, oriented, and has the ability to understand the circumstances surrounding his/her illness or impairment, as well as the possible risks associated with refusing treatment and/or transport, and can communicate their decision typically is considered to have decision-making capacity
- The individual's judgment must also not be significantly impaired by illness, injury or drugs/alcohol intoxication. Individuals who have attempted suicide, verbalized suicidal intent, or have other factors that lead EMS providers to suspect suicidal intent, should not be regarded as having decision-making capacity and may not decline transport to a medical facility

Treatment and Interventions

1. Obtain a complete set of vital signs and complete an initial assessment, paying particular attention to the individual's neurologic and mental status.
2. Determine the individual's capacity to make a valid judgment concerning the extent of his/her illness or injury; if the EMS provider has doubts about whether the individual has the mental capacity to refuse or if the patient lacks capacity, the EMS provider should contact on-line medical control.
3. If patient has capacity, clearly explain to the individual and all responsible parties the possible risks and overall concerns with regards to refusing care.
4. Perform appropriate medical care with the consent of the individual.
5. Complete the patient care report clearly documenting the initial assessment findings and the discussions with all involved individuals regarding the possible consequences of refusing additional prehospital care and/or transportation.

Notes and Educational Pearls

Key Considerations

- An adult or emancipated minor who has demonstrated possession of sufficient mental capacity for making decisions has the right to determine the course of his or her medical care, including the refusal of care. Emancipated minors can make decisions regarding their health care. An "emancipated minor" means a minor who is or has been married; a minor who has previously given birth; or a minor who has been legally freed from the care, custody and control of her parents, with little likelihood of returning to the care, custody and control prior to marriage or prior to reaching the age of majority. These individuals must be advised of the risks and consequences resulting from refusal of medical care.

- An individual determined to lack decision-making capacity by EMS providers should not be allowed to refuse care against medical advice or to be released at the scene. Mental illness, drugs, alcohol intoxication, or physical or mental impairment may significantly impair a patient's decision-making capacity. Individuals who have attempted suicide, verbalized suicidal intent, or have other factors that lead EMS providers to suspect suicidal intent, should not be regarded as having demonstrated sufficient decision-making capacity.
- The determination of decision-making capacity may be challenged by communication barriers or cultural differences.
- EMS providers should not put themselves in danger by attempting to treat and/or transport an individual who refuses care.
- Always act in the best interest of the patient and contact on-line medical control when appropriate.

Special Considerations—Minors

- It is preferable for minors to have a parent or legal guardian who can provide consent for treatment on behalf of the child.
- All states allow health care providers to provide emergency treatment when a parent is not available to provide consent. This is known as the emergency exception rule or the doctrine of implied consent. For minors, this doctrine means that the prehospital professional can presume consent and proceed with appropriate treatment and transport if the following six conditions are met:
 - i. The child is suffering from an emergent condition that places his or her life or health in danger
 - ii. The child's legal guardian is unavailable or unable to provide consent for treatment or transport
 - If a minor is injured or ill and no parent contact is possible, the provider may contact on-line medical control for additional instructions
 - iii. Treatment or transport cannot be safely delayed until consent can be obtained
 - iv. The prehospital professional administers only treatment for emergency conditions that pose an immediate threat to the child

Note: As a general rule, when the prehospital professional's authority to act is in doubt, EMS providers should always do what they believe to be in the best interest of the minor

Cardiovascular

Adult and Pediatric Syncope and Presyncope

Aliases

Loss of consciousness, passed out, fainted

Patient Care Goals

1. Stabilize and resuscitate when necessary
2. Initiate monitoring and diagnostic procedures
3. Transfer for further evaluation

Patient Presentation

Syncope is heralded by both the loss of consciousness and the loss of postural tone and resolves spontaneously without medical interventions. Syncope typically is abrupt in onset and resolves equally quickly. EMS providers may find the patient awake and alert on initial evaluation. Presyncope is defined as the prodromal symptoms of syncope. It usually lasts for seconds to minutes and may be described by the patient as “nearly blacking out” or “nearly fainting.”

Inclusion Criteria

1. Abrupt loss of consciousness with loss of postural tone
2. Prodromal symptoms of syncope

Exclusion Criteria

Conditions other than the above, including:

1. Patients with alternate and obvious cause of loss of consciousness (e.g. trauma—go to [Head Injury guideline](#)).
2. Patients with ongoing mental status changes or coma (go to [Altered Mental Status guideline](#)).

Patient Management

Assessment

1. Pertinent history
 - a. Review the patient’s past medical history, including a history of:
 - i. Cardiovascular disease (e.g. cardiac disease/stroke)
 - ii. Seizure
 - iii. Recent trauma
 - iv. Anticoagulation
 - v. Dysrhythmia
 - vi. Congestive heart failure (CHF)
 - vii. Syncope
 - viii. Current Medications
 - b. History of Present Illness, including:
 - i. Conditions leading to the event.
 - ii. Patient complaints before or after the event including prodromal symptoms.

- iii. History from others on scene, including seizures or shaking, presence of pulse and breathing (if noted), duration of the event, and events that lead to the resolution of the event.
 - c. Review of systems:
 - i. Occult blood loss (GI/GU)
 - ii. Fluid losses (nausea/vomiting/diarrhea) and fluid intake
- 2. Pertinent Physical exam including:
 - a. Attention to vital signs as well as evaluation for trauma.
 - b. Detailed neurologic exam (including stroke screening and mental status).
 - c. Heart, lung, abdominal and extremity exam.
 - d. Additional evaluation:
 - i. ECG cardiac monitoring
 - ii. Ongoing vital signs
 - iii. 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*]

Treatment and Interventions:

Should be directed at abnormalities discovered in the physical exam or on additional examination and may include management of cardiac dysrhythmias, cardiac ischemia or infarct, hemorrhage, shock, and the like.

- a. Manage airway as indicated.
- b. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- c. Evaluate for hemorrhage and treat for shock if indicated.
- d. Establish IV/IO access [*AEMT-R*].
- e. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]) if clinically appropriate.
- f. Apply ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*].
- g. Apply 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*].
- h. Monitor for and treat arrhythmias (if present refer to appropriate guideline).

Patient Safety Considerations:

- 1. Patients suffering syncope due to arrhythmia may suffer recurrent arrhythmia and should therefore be placed on a ECG cardiac monitor.
- 2. Geriatric patients suffering falls from standing may sustain significant injury and should be diligently screened for trauma; go to [General Trauma Management guideline](#).

Notes and Educational Pearls

Key Considerations

- By being most proximate to the scene and to the patient's presentation, EMS providers are commonly in a unique position to identify the cause of syncope. Consideration of potential causes, ongoing monitoring of vitals and cardiac rhythm as well as detailed exam and history are essential pieces of information to pass onto hospital providers.
- All patients suffering from syncope deserve hospital level evaluation, even if they appear normal with few complaints on scene

- High risk causes of syncope include the following:
 - a. Cardiovascular
 - i. Myocardial infarction
 - ii. Aortic stenosis
 - iii. Hypertrophic cardiomyopathy
 - iv. Pulmonary embolus
 - v. Thoracic aortic dissection
 - vi. Lethal dysrhythmia
 - b. Neurovascular
 - i. Intracranial hemorrhage
 - ii. Transient ischemic attack or stroke
- Consider high risk 12-lead ECG features including, but not limited to:
 - a. Evidence of QT prolongation (generally over 500ms)
 - b. Delta waves
 - c. Brugada syndrome (incomplete RBBB pattern in V1/V2 with ST segment elevation)
 - d. Hypertrophic obstructive cardiomyopathy

Pertinent Assessment Findings

- Evidence of trauma
- Evidence of cardiac dysfunction (e.g. evidence of CHF, arrhythmia)
- Evidence of hemorrhage
- Evidence of neurologic compromise
- Evidence of alternate etiology, including seizure
- Initial and ongoing cardiac rhythm
- 12-lead ECG findings

ACUTE CORONARY SYNDROME (ACS) OR ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION (STEMI)

Aliases

Heart attack, myocardial infarction (MI)

Patient Care Goals

1. Identify STEMI quickly.
2. Determine the time of symptom onset.
3. Activate hospital-based STEMI system of care.
4. Monitor vital signs and cardiac rhythm and be prepared to provide CPR and defibrillation if needed.
5. Administer appropriate medications.
6. Transport to appropriate facility.

Patient Presentation

Inclusion Criteria

1. Patient presents with chest pain or discomfort in other areas of the body (e.g. arm, jaw, epigastrium) of suspected cardiac origin; shortness of breath; sweating; nausea; vomiting; and dizziness. Atypical or unusual symptoms are more common in women, the elderly and diabetic patients. May also present with CHF, syncope and/or shock.
2. Some patients will present with likely non-cardiac chest pain and otherwise have a low likelihood of ACS (e.g. blunt trauma to the chest of a child). For these patients, defer the administration of aspirin and nitrates per the [Pain Management guideline](#).

Exclusion Criteria

None recommended

Patient Management

Assessment, Treatment, and Interventions

1. Evaluate for signs and symptoms that include chest pain, congestive heart failure, syncope, shock, symptoms similar to a patient's previous MI.
2. Assess the patient's cardiac rhythm; treat pulseless rhythms, tachycardia, or symptomatic bradycardia [see [Cardiovascular](#) and [Resuscitation](#) guidelines]
3. If the patient is dyspneic, hypoxemic, or has obvious signs of heart failure, EMS providers should administer oxygen as appropriate with a target of achieving 94-98% saturation [see [Universal Care guideline](#)]
4. Use the 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*]; it is the primary diagnostic tool that identifies a STEMI. It is imperative that EMS providers routinely acquire a 12-lead ECG within 10 minutes for all patients exhibiting signs and symptoms of ACS.
 - a. The ECG may be transmitted for remote interpretation by a physician or screened for STEMI by properly trained EMS providers with or without the assistance of computer- interpretation.

- b. Advance notification should be provided to the receiving hospital for patients identified as having STEMI.
 - c. Deploy serial ECGs as clinically indicated for concerns of evolving acute coronary syndrome.
 - d. All ECGs should be made available to treating personnel at the receiving hospital, whether brought in or transmitted from the field.
5. Administer aspirin *[EMR-O; EMT-R]*.
6. Establish IV access *[AEMT-R]*.
7. Nitroglycerin *[AEMT-R]*
 - a. Avoid the use of nitrates in any patient who has used a phosphodiesterase inhibitor within the past 48 hours. Examples are:
 - Sildenafil (Viagra®, Revatio®)
 - Vardenafil (Levitra®, Staxyn®)
 - Tadalafil (Cialis®, Adcirca®) which are used for erectile dysfunction and pulmonary hypertension.
 - b. Avoid use in patients receiving intravenous epoprostenol (Flolan®) or treprostenil (Remodulin®) which is used for pulmonary hypertension.
 - c. Administer nitrates with extreme caution, if at all, to patients with inferior-wall STEMI or suspected right ventricular (RV) involvement, because these patients require adequate RV preload.
 - d. Analgesia is indicated in STEMI when chest discomfort is unresponsive to nitrates
8. Transport and destination decisions should be based on local resources and system of care.

Patient Safety Considerations

1. Observe for signs of clinical deterioration: dysrhythmias, CP, SOB, decreased LOC/syncope, or other signs of shock/hypotension.
2. Perform serial 12-lead ECGs (especially any time clinical changes noted).

Notes and Educational Pearls

Key Considerations

Acute coronary syndrome may present with atypical pain, vague or only generalized complaints.

Pertinent Assessment Findings

A complete medication list should be obtained from each patient. It is especially important for the treating physician to be informed if the patient is taking beta-blockers, calcium channel blockers, clonidine, digoxin, blood thinners (anticoagulants), or medications for the treatment of erectile dysfunction or pulmonary hypertension.

Bradycardia

Aliases

Heart block, junctional rhythm

Patient Care Goals

1. Maintain adequate perfusion.
2. Treat underlying cause:
 - a. Hypoxia
 - b. Shock
 - c. Second or third-degree AV block
 - d. Toxin exposure (beta-blocker, calcium channel blocker, organophosphates, digoxin)
 - e. Electrolyte disorder
 - f. Hypoglycemia
 - g. Increased intracranial pressure (ICP)
 - h. Other

Patient Presentation

Inclusion Criteria

1. Heart rate less than 50 beats per minute with either symptoms (AMS, CP, CHF, seizure, syncope, shock, pallor, diaphoresis) or evidence of hemodynamic instability
2. The major ECG rhythms classified as bradycardia include:
 - a. Sinus bradycardia
 - b. Second-degree AV block
 - i. Type I —Wenckenbach/Mobitz I
 - ii. Type II —Mobitz II
 - c. Third-degree AV block complete block
 - d. Ventricular escape rhythms
3. See additional inclusion criteria, below, for pediatric patients

Exclusion Criteria

No recommendations

Patient Management


Assessment, Treatment, and Interventions

Adult Management

- a. Manage airway as necessary.
 - i. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- b. Initiate monitoring and perform 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
- c. Establish IV access *[AEMT-R]*.
- d. Check blood glucose and treat hypoglycemia *[EMR-O; EMT-R]* per the [Hypoglycemia](#) and [Hyperglycemia](#) guidelines.

- e. Consider the following additional therapies if symptomatic bradycardia and hemodynamic instability continue:
- Atropine **[INT-R]**
 - Vasopressor medications (in order of preference)
 - Epinephrine infusion titrated to a MAP greater than 65 mmHg **[INT-R]**
or
 - Epinephrine by push dose (dilute boluses) **[INT-R]** every 3-5 minutes titrated MAP greater than 65mmHg
 - Norepinephrine infusion titrated to a MAP greater than 65 mmHg **[PARA-O]**
 - Transcutaneous pacing **[INT-R]**; if pacing is performed, consider sedation or pain control

Adult Bradycardia Algorithm



© 2020 American Heart Association


2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/hghlghts_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

Pediatric Management

Treatment is only indicated for patients who are symptomatic (pale and/or cyanotic, diaphoretic, altered mental status, hypoxic)

- a. Initiate chest compressions for heart less than 60 and signs of poor perfusion (altered mental status, hypoxia, hypotension, weak pulse, delayed capillary refill, cyanosis).
- b. Manage airway and assist ventilations as necessary with minimally interrupted chest compressions using a compression to ventilation ratio 15:2 (30:2 if single provider is present).
- c. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- d. Initiate ECG monitoring and perform 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
- e. Establish IV access *[AEMT-R]*.
- f. Check blood glucose *[EMR-O; EMT-R]* and treat hypoglycemia per the [Hypoglycemia guideline](#).
- g. Consider the following additional therapies if bradycardia and symptoms or hemodynamic instability continue:
 - i. Epinephrine by push dose *[INT-R]* every 3-5 minutes titrated to MAP greater than 65mmHg
 - ii. Atropine *[INT-R]* if increased vagal tone or cholinergic drug toxicity (maximum total dose of 3 mg)
 - iii. Transcutaneous pacing *[INT-R]*; if pacing is performed, consider sedation or pain control
 - iv. Epinephrine *[INT-R]* for bradycardia and poor perfusion unresponsive to ventilation and oxygenation
 - a. **Note:** It is reasonable to administer atropine for bradycardia caused by increased vagal tone or cholinergic drug toxicity

Figure 12. Pediatric Bradycardia With a Pulse Algorithm.



eccguidelines.heart.org

17

2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/hghlghts_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

Patient Safety Considerations

If pacing is performed, consider sedation or pain control.

Notes and Educational Pearls

Key Considerations

- Observe for signs of decreased end-organ perfusion: chest pain (CP), shortness of breath (SOB), decreased level of consciousness, syncope or other signs of shock/hypotension.
- Be aware that patients who have undergone cardiac transplant will not respond to atropine.

- Consider potential culprit medications including beta-blockers, calcium channel blockers, sodium channel blockers/anti-depressants, digoxin, and clonidine.
Note: If medication overdose is considered, refer to appropriate guideline in the [Toxins and Environmental](#) section
- Consider whether the differential diagnosis includes the following: MI, hypoxia, pacemaker failure, hypothermia, sinus bradycardia, athletes, head injury with increased ICP, stroke, spinal cord lesion, sick sinus syndrome, AV blocks, overdose, cholinergic nerve agents.
- Consider hyperkalemia in the patient with wide complex bradycardia.
- Manage bradycardia via the least invasive manner possible, escalating care as needed.
 - Third-degree heart block or the denervated heart (as in cardiac transplant) may not respond to atropine and in these cases, proceed quickly to chronotropic agents (such as epinephrine or dopamine), or transcutaneous pacing.
 - Dopamine is not indicated for pediatric patients.
 - In cases of impending hemodynamic collapse, proceed directly to transcutaneous pacing.
- Be aware of acute coronary syndrome as a cause of bradycardia in adult patients.
- When dosing medications for pediatric patients, dose should be weight-based for non-obese patients and based on ideal body weight for obese patients.
- Although dopamine is often recommended for the treatment of symptomatic bradycardia, recent research suggests that patients in cardiogenic or septic shock treated with norepinephrine have a lower mortality rate compared to those treated with dopamine.
- **Caution:** Norepinephrine can theoretically cause reflex bradycardia.

Pertinent Assessment Findings

No recommendations

Implantable Ventricular Assist Devices

Aliases

Ventricular assist device (VAD), left ventricular assist device (LVAD), right ventricular assist device (RVAD), biventricular assist device (BiVAD)

Patient Care Goals

1. Rapid identification of, and interventions for, cardiovascular compromise in patients with VADs
2. Rapid identification of, and interventions for VAD-related malfunctions or complications

Patient Presentation

Inclusion Criteria

1. Adult patients that have had an implantable ventricular assist device (VAD), including a left ventricular assist device (LVAD), right ventricular assist device (RVAD), or biventricular-assist device (BiVAD), and have symptoms of cardiovascular compromise
2. Patients with VADs that are in cardiac arrest
3. Patients with VADs that are experiencing a medical or injury-related event not involving the cardiovascular system or VAD malfunction

Exclusion Criteria

Adult patients who do not have a VAD in place

Patient Management

Assessment

1. Assess for possible pump malfunction.
 - a. Assess for alarms.
 - b. Auscultate for pump sound "hum."
 - c. Look for signs of hypoperfusion including pallor, diaphoresis, or altered mental status.
2. Assess the functionality of the VAD pump. If it has malfunctioned:
 - a. Utilize available resources to troubleshoot potential VAD malfunctions and to determine appropriate corrective actions to restore normal VAD function:
 - i. Contact the patient's VAD-trained companion, if available.
 - ii. Contact the patient's VAD coordinator, using the phone number on the device.
 - iii. Check all the connections to system controller.
 - iv. Change VAD batteries, and/or change system controller if indicated.
 - v. Have patient stop all activity and assess for patient tolerance.
 - vi. Follow appropriate cardiovascular condition-specific protocol(s) as indicated.

Treatment and Interventions

1. Manage airway as indicated.
2. Apply ECG cardiac monitoring [*Acquisition EMT-O; Interpretation INT-R*].
3. Establish IV access
4. Acquire 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*]

5. Assess patient for VAD-related complications or cardiovascular problems. If present, expedite transport to the medical facility where VAD was placed if patient's clinical condition and time allows.
 - a. Non-cardiovascular-related problem with functioning VAD: Transport to a facility that is appropriate for the patient's main presenting problem without manipulating the device
 - b. Hypofusing with functioning VAD:
 - i. Administer normal saline bolus 20ml/kg IV/IO [**AEMT-R**] maximum of 1 liter) over less than 15 minutes, using a push-pull method of drawing up the fluid in a syringe and pushing it through the IV/IO or pressure bag.
 - ii. Repeat up to 3 times based on patient's condition and clinical impression for a total cumulative dose not to exceed 3 L.
6. If patient is in full cardiac arrest:
 - a. Do not perform CPR if there is any evidence the pump is still functioning, the decision whether to perform CPR should be made based upon best clinical judgment in consultation with the patient's VAD-trained companion and the VAD coordinator (or on-line medical control if VAD coordinator unavailable)
 - b. Initiate CPR **only** where:
 - i. You have confirmed the pump has stopped and troubleshooting efforts to restart it have failed, **and**
 - ii. The patient is unresponsive and has no detectable signs of life.

Notes and Educational Pearls

- You do not need to disconnect the controller or batteries in order to:
 - Defibrillate or cardiovert [**INT-R**].
 - Acquire a 12-lead ECG.
- Automatic non-invasive cuff blood pressures [**ALL EMS PRACTICE LEVELS-O**] may be difficult to obtain due to the narrow pulse pressure created by the continuous flow pump.
- Flow through many VAD devices is not pulsatile and patients may not have a palpable pulse or accurate pulse oximetry.
- The blood pressure, if measurable, may not be an accurate measure of perfusion.
- Ventricular fibrillation, ventricular tachycardia, or asystole/PEA may be the patient's "normal" underlying rhythm. Evaluate clinical condition and provide care in consultation with VAD coordinator.
- The patient's travel bag should accompany them at all times with back-up controller and spare batteries.
- If feasible, bring the patient's power module, cable, and display module to the hospital.
- All patients should carry a spare pump controller with them.
- The most common cause for VAD alarms are low batteries or battery failures.
- Although automatic non-invasive blood pressure cuffs are often ineffective in measuring systolic and diastolic pressure, if they do obtain a measurement, the MAP is usually accurate.

- Other VAD complications:
 - Infection
 - Stroke/TIA
 - Bleeding
 - Arrhythmias
 - Cardiac tamponade
 - CHF
 - Aortic insufficiency

Tachycardia with a Pulse

Aliases

Supraventricular tachycardia (SVT), ventricular tachycardia (VT), multifocal atrial tachycardia (MAT), torsades, atrial fibrillation (A-FIB), atrial flutter

Patient Care Goals

1. Maintain adequate oxygenation, ventilation, and perfusion.
2. Control ventricular rate.
3. Restore regular sinus rhythm in unstable patient.
4. Search for underlying cause:
 - a. Medications (caffeine, diet pills, thyroid, decongestants)
 - b. Drugs (cocaine, amphetamines)
 - c. History of dysrhythmia
 - d. CHF

Patient Presentation

Patients will manifest elevated heart rate for age and may or may not also present with associated symptoms such as palpitations, dyspnea, chest pain, syncope/near-syncope, hemodynamic compromise, altered mental status, or other signs of end organ malperfusion.

Inclusion Criteria

Heart rate greater than 150 bpm in adults or relative tachycardia in pediatric patients (220 infant/ 180 child)

Exclusion Criteria

Sinus tachycardia

Patient Management


Assessment, Treatments, and Interventions

Adult Management

- a. Manage airway as necessary.
- b. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- c. Initiate ECG monitoring *[Acquisition EMT-O; Interpretation INT-R]* and perform 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
- d. Establish IV access *[AEMT-R]*.
- e. Check blood glucose *[EMR-O; EMT-R]* and treat hypoglycemia per the [Hypoglycemia guideline](#).
- f. Consider the following additional therapies if tachycardia and symptoms or hemodynamic instability continue:
 - i. **Regular Narrow Complex Tachycardia—Stable (SVT)**
 1. Perform vagal maneuvers.
 2. Administer adenosine *[INT-R]* (proximal site) followed by 10 mL fluid bolus.
 - a. If tachycardia continues, repeat adenosine.
 3. Administer diltiazem *[PARA-O]*.
 - a. After 15 minutes, a second dose of diltiazem may be given if needed.

4. Administer metoprolol *[PARA-O]*. May repeat as needed every 5 minutes for a total of 3 doses.
- ii. **Regular Narrow Complex Tachycardia—Unstable**
 1. Deliver a synchronized shock *[INT-R]* based on manufacturer's recommendations.
 2. Consider sedation and analgesia for responsive patients.
- iii. **Irregular Narrow Complex Tachycardia—Stable** (atrial fibrillation, atrial flutter, multifocal atrial tachycardia), administer:
 1. Diltiazem *[PARA-O]*, *or*
 2. Metoprolol *[PARA-O]*
- iv. **Irregular Narrow Complex Tachycardia—Unstable**
 1. Deliver a synchronized shock *[INT-R]* based on manufacturer's recommendation
 2. Consider sedation for responsive patients.
- v. **Regular Wide Complex Tachycardia—Stable** (ventricular tachycardia, supraventricular tachycardia, atrial fibrillation/flutter with aberrancy, accelerated idioventricular rhythms, pre-excited tachycardias with accessory pathways), administer:
 1. Amiodarone *[PARA-O]* *or*
 2. Procainamide *[PARA-O]* *or*
 3. Lidocaine *[INT-O; PARA-R]*.
 4. Adenosine, if monomorphic tachycardia *[INT-R]*.
- vi. **Regular Wide Complex Tachycardia—Unstable**
 1. Deliver a synchronized shock *[INT-R]* based on manufacturer's recommendation
 2. Consider sedation for responsive patients.
- vii. **Irregular Wide Complex Tachycardia—Stable** (atrial fibrillation with aberrancy, pre-excited atrial fibrillation (i.e. atrial fibrillation using an accessory pathway), MAT or polymorphic VT/torsades de pointes.
 1. Amiodarone *[PARA-O]* *or*
 2. Procainamide *[PARA-O]* *or*
 3. Magnesium sulfate if torsades de pointes *[PARA-O]*
- viii. **Irregular Wide Complex Tachycardia—Unstable**
 1. Deliver a synchronized shock *[INT-R]* based on manufacturer's recommendation.
 2. Consider sedation for responsive patients.

Adult Tachycardia With a Pulse Algorithm




2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/hghlights_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

2. Pediatric Management

- Manage airway as necessary.
- Administer oxygen **[EMR-O; EMT-R]** as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- Initiate monitoring and perform 12-lead ECG **[Acquisition EMT-O; Interpretation INT-R]**.
- Establish IV/IO access **[AEMT-R]**.
- Check blood glucose **[EMR-O; EMT-R]** and treat hypoglycemia per the [Hypoglycemia guideline](#).
- Consider the following additional therapies if tachycardia and symptoms or hemodynamic instability continue:
 - Regular Narrow Complex Tachycardia—Stable (SVT)**
 - Perform vagal maneuvers **[EMT-O; ALL OTHER EMS PROVIDERS-R]**.
 - Administer adenosine **[INT-R]**.
 - If unsuccessful, you may repeat.
 - Regular Narrow Complex Tachycardia—Unstable**
 - Deliver a synchronized shock **[INT-R]**: 0.5-1 J/kg for the first dose.
 - Use 2 J/kg for repeat doses.
 - Regular, Wide Complex Tachycardia—Stable**
 - Consider adenosine **[INT-R]** for SVT with aberrancy.

2. Otherwise give amiodarone over 10 minutes *[INT-O; PARA-R]*
- iv. **Regular, Wide Complex Tachycardia – Unstable**
 1. Synchronized cardioversion 0.5-1.0 J/kg *[INT-R]*

Figure 13. Pediatric Tachycardia With a Pulse Algorithm.



2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/highlights_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

Notes and Educational Pearls

Key Considerations

- Causes:
 - Hypovolemia
 - Hypoxia
 - Hydrogen (acidosis)
 - Myocardial infarction
 - Hypokalemia or hyperkalemia
 - Hypoglycemia
 - Hypothermia
 - Toxins or Overdose

- Tamponade
- Tension pneumothorax
- Thrombus—central or peripheral
- Trauma
- Hyperthyroidism
- Atrial fibrillation rarely requires cardioversion in the field. As it is difficult to ascertain onset of rhythm, risk of stroke needs to be considered prior to cardioversion.
- A wide-complex irregular rhythm should be considered pre-excited atrial fibrillation; extreme care must be taken in these patients.
 - Characteristic ECG findings include a short PR interval and, in some cases, a delta wave.
 - Avoid AV nodal blocking agents such as adenosine, calcium channel blockers, digoxin, and possibly beta-blockers in patients with pre-excitation atrial fibrillation (e.g. Wolff- Parkinson-White Syndrome, Lown-Ganong-Levine Syndrome) because these drugs may cause a paradoxical increase in the ventricular response.
 - Blocking the AV node in some of these patients may lead to impulses that are transmitted exclusively down the accessory pathway, which can result in ventricular fibrillation.
 - Amiodarone or procainamide may be used as an alternative.
- Amiodarone or procainamide can be used as a rate-controlling agent for patients who are intolerant of or unresponsive to other agents, such as patients with CHF who may not otherwise tolerate diltiazem or metoprolol.
- Biphasic waveforms have been proven to convert atrial fibrillation at lower energies and higher rates of success than monophasic waveforms.

Strategies include dose escalation (70, 120, 150, 170 J for biphasic or 100, 200, 300, 360 J for monophasic) versus beginning with single high energy/highest success rate for single shock delivered.
- Studies in infants and children have demonstrated the effectiveness of adenosine for the treatment of hemodynamically stable or unstable SVT.
- Adenosine should be considered the preferred medication for stable SVT.
 - Verapamil may be considered as alternative therapy in older children but should not be routinely used in infants.
 - Procainamide or amiodarone given by a slow IV infusion with careful hemodynamic monitoring may be considered for refractory SVT.

Pertinent Assessment Findings

No recommendations

Patient Safety Considerations

- Only use one antidysrhythmic at a time.
- Patients who receive metoprolol and diltiazem are at significant risk for hypotension and bradycardia.
- If using cardioversion, consider sedation and pain control.
- With irregular wide complex tachycardia (atrial fibrillation with aberrancy such as Wolff- Parkinson-White and Lown-Ganong Levine), avoid use of AV nodal blocking agents (e.g. adenosine, calcium channel blockers, beta blockers).
- Patients with Wolff-Parkinson-White should be given procainamide prior to amiodarone.

Suspected Stroke/Transient Ischemic Attack

Aliases

Cerebrovascular accident (CVA), TIA

Patient Care Goals

1. Detect neurological deficits
2. Determine eligibility for transport to a stroke center

Patient Presentation

1. Neurologic deficit such as facial droop, localized weakness, gait disturbance, slurred speech, altered mentation
2. Hemiparesis or hemiplegia
3. Dysconjugate gaze, forced or crossed gaze (if patient is unable to voluntarily respond to exam, makes no discernible effort to respond, or is unresponsive)
4. Severe headache, neck pain/stiffness, difficulty seeing

Inclusion Criteria

1. Patient has signs and symptoms consistent with stroke or transient ischemic attack (TIA)

Exclusion Criteria

1. If glucose less than 60 mg/dL [*EMR-O; EMT-R*], treat per the [Hypoglycemia guideline](#)
2. If trauma and GCS less than or equal to 13, treat per the [Head Injury](#) and [General Trauma Management](#) guidelines

Patient Management

Assessment

1. Use a validated prehospital stroke scale that may include, but is not limited to:
 - a. Facial smile/grimace—ask patient to smile
 - b. Arm—close eyes and hold out arms for count of 10 seconds
 - c. Speech—“You can’t teach an old dog new tricks”
2. Pertinent historical data includes:
 - a. History—“last known well” and source of that information
 - b. Neurologic status assessment [see [Appendix VII](#)]
 - c. Medication—patient is taking warfarin or any anticoagulant medication
3. Evaluate for the presence of stroke mimics including:
 - a. Hypoglycemia
 - b. Seizure
 - c. Sepsis
 - d. Migraine
 - e. Intoxication

Treatment and Interventions

1. Determine “last known well” time.
2. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
3. Treat seizures per [Seizures guideline](#), if seizure activity present.

4. Check blood glucose level [*EMR-O; EMT-R*].
 - a. Treat only if glucose less than 60 mg/dL.
5. Acquire 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*], if possible.
6. Use the local stroke plan for hospital notification

Patient Safety Considerations

1. Prevent aspiration—elevate head of stretcher 15–30 degrees if systolic BP greater than 100 mm Hg.
 - a. Maintain head and neck in neutral alignment, without flexing the neck.
2. Protect paralyzed limbs from injury.
3. Avoid multiple IV attempts.

Notes and Educational Pearls

Key Considerations

Adult care

- Base transport and destination decisions on local resources and stroke system of care.
 - Destinations hospitals may include:
 - Stroke Ready
 - Primary Stroke Center
 - Comprehensive Stroke Center
- Do not treat hypertension.
- Place on ECG cardiac monitor.

Pediatric care

- Treatment principles remain the same
- Although rare, pediatric patients can have strokes
- Stroke scales are not validated for pediatric patients

General Medical

Abdominal Pain

Aliases

None

Patient Care Goals

1. Improve patient comfort.
2. Identify life-threatening causes of abdominal pain.

Patient Presentation

Inclusion Criteria

Abdominal pain or discomfort related to a non-traumatic cause

Exclusion Criteria

1. Abdominal pain due to trauma [see [General Trauma Management guideline](#)]
2. Abdominal pain due to or related to pregnancy [see [OB/GYN guidelines](#)]

Patient Management

Assessment

1. Perform airway assessment and management per the [Airway Management guideline](#).
2. Obtain vital signs including pulse, respiratory rate, pulse oximetry, and blood pressure.
3. Provide evaluation and management of pain per the [Pain Management guideline](#).
4. Obtain vascular access [*AEMT-R*] as necessary to provide analgesia and/or fluid resuscitation.
5. Assess for life-threatening causes of abdominal pain, which may include:
 - a. Ischemic, necrotic, or perforated bowel.
 - i. Severe tenderness
 - ii. Abdominal pain with motion or “jiggling” of the abdomen
 - iii. Fever
 - iv. Bloody stool
 - v. Nausea and vomiting
 - vi. Possible absence of passage of stool or gas
 - vii. Abdominal distention, with possible tympany to percussion
 - b. Dissecting or ruptured abdominal aortic aneurysm (AAA).
 - i. Unequal femoral or distal lower extremity pulses
 - ii. “Pulsatile” abdominal mass
 - iii. Associated back pain and/or chest pain
 - iv. Known history of abdominal aortic aneurysm
 - c. Ruptured ectopic pregnancy.
 - i. Vaginal bleeding
 - ii. Recently diagnosed pregnancy
 - d. Recent missed period/menstrual cycle in women of childbearing age.

- e. Appendicitis.
 - i. Focal right lower quadrant tenderness, possibly with rebound and guarding
 - ii. Right lower quadrant tenderness noted during palpation of the left lower quadrant (positive Rovsing's sign)
 - iii. Peri-umbilical or diffuse abdominal tenderness with palpation or "jiggling" of the abdomen/pelvis
 - iv. Fever
 - v. Nausea, vomiting
 - vi. Lack of appetite
 - f. Acute Cholecystitis.
 - i. Right upper quadrant or epigastric tenderness
 - ii. Fever
 - iii. Nausea and vomiting
 - iv. Possible history of gallstones
 - g. Pyelonephritis.
 - i. Fever
 - ii. Nausea, vomiting
 - iii. Urinary frequency/urgency
 - iv. Dysuria
 - v. Hematuria
 - vi. Back/flank pain
 - vii. Costovertebral angle tenderness to percussion
6. Assess for signs of shock.
- a. If shock is present, provide treatment per appropriate [Shock guideline](#)
7. Assess for other non-life-threatening causes of abdominal pain.
- a. Kidney stone
 - i. Unilateral flank pain
 - ii. Nausea, vomiting
 - iii. Possible Hematuria

Treatment and Interventions

Medication Administration:

1. Provide analgesia per the [Pain Management guideline](#).
2. Administer antiemetics per the [Nausea-Vomiting guideline](#).
3. Provide transport to an appropriate receiving facility. Consider specialty destination centers for conditions such as suspected abdominal aortic aneurysm.
4. Reassess vital signs and response to therapeutic interventions throughout transport.

Patient Safety Considerations

None recommended

Notes and Educational Pearls

Key Considerations

- Assess for life-threatening causes of abdominal pain.
- Provide appropriate treatment for pain, vomiting, and shock.
- Consider transport to a trauma center if aortic aneurysm is suspected.

Pertinent Assessment Findings

- Rebound tenderness
- Guarding
- Abdominal distension
- Abdominal tympany to percussion
- Tenderness focal to a specific abdominal quadrant
- Presence of “pulsatile” abdominal mass
- Absence of or significant inequality of femoral or distal arterial pulses in lower extremities
- Hyper or hypothermia
- Rectal bleeding, hematemesis (character), vaginal bleeding

Abuse and Maltreatment

Aliases

Maltreatment of vulnerable populations

Definitions

- **Abuse and Maltreatment:** Abuse and Maltreatment is any act or series of acts of commission or omission by a caregiver or person in a position of power over the patient that results in harm, potential for harm, or threat of harm to a patient
- **Child Maltreatment and Abuse:** Child maltreatment includes any act or series of acts of commission or omission by a parent or other caregiver that results in harm, potential for harm, or threat of harm to a child. An act of commission (child abuse) is the physical, sexual or emotional maltreatment or neglect of a child or children. An act of omission (child neglect) includes, but is not limited to, failure to provide for the child's needs (e.g. physical, emotional, medical/dental, and educational neglect) and failure to supervise (e.g. inadequate supervision or safety precautions, lack of appropriate car seat use, and exposure to violent or dangerous environments).
- **Human Trafficking:** Human trafficking is when people are abducted or coerced into service and often transported across international borders. Signs may include, but are not limited to: patient with branding/tattoos and environmental clues such as padlocks and/or doorknobs removed on interior doors, and intact windows that are boarded up

Patient Care Goals

1. Recognize any act or series of acts of commission or omission by a caregiver or person in a position of power over the patient that results in harm, potential for harm, or threat of harm to a patient.
2. Take appropriate steps to protect the safety of the responders as well as bystanders.
3. Get the patient out of immediate danger.
4. Assess any patient injuries that may be the result of acute or chronic events.
5. Attempt to preserve evidence whenever possible; however, the overriding concern should be providing appropriate emergency care to the patient.

Patient Presentation

1. Clues to abuse or maltreatment can vary with age group of the patient and type of abuse.
2. Not all abuse or maltreatment is physical.
3. EMS role is to:
 - a. Document concerns.
 - b. Assess potentially serious injuries.
 - c. Disclose concerns to appropriate authorities.
 - d. Initiate help to get the patient into a safe situation.
 - e. Not investigate or intervene beyond the steps above.
 - f. Leave further intervention to law enforcement personnel.

Inclusion/Exclusion Criteria

Absolute inclusion/exclusion criteria are not possible in this area. Rather, clues consistent with different types of abuse/maltreatment should be sought:

- Potential clues to abuse/maltreatment from caregivers or general environment:
 - Apathy by caregiver about patient's current situation
 - Overreaction by caregiver to questions about situation
 - Inconsistencies from caregivers or bystanders regarding history of what happened
 - Inconsistencies between injury patterns and information provided by caregivers or patient
 - Injuries not appropriate for patient's age or physical abilities (e.g. infants with injuries usually associated with ambulatory children, elders who have limited mobility with injury mechanisms inconsistent with their capabilities)
 - Controlling behavior by caregiver (e.g., not allowing adult patient to speak for themselves—pay special attention to patients who cannot communicate due to young age or language and/or cultural barriers)
 - Inadequate safety precautions or facilities where the patient lives and/or evidence of security measures that appear to confine the patient inappropriately
- Potential clues to abuse or maltreatment that can be obtained from the patient:
 - Multiple bruises in various stages of healing
 - Age-inappropriate behavior (e.g. adults who are submissive or fearful, children who act in a sexually inappropriate way)
 - Pattern burns, bruises, or scars suggestive of specific weaponry used
 - Evidence of medical neglect for injuries or infections
 - Unexplained trauma to genitourinary systems or frequent infections to this system
 - Evidence of malnourishment and/or serious dental problems
- Have a high index of suspicion for abuse in children presenting with a Brief Resolved Unexplained Event (BRUE) [see [BRUE guideline](#)]

Patient Management

Assessment

1. Start with a primary survey and identify any potentially life-threatening issues.
2. Document thorough secondary survey to identify clues suggesting potential abuse/maltreatment:
 - a. Inability to communicate due to developmental age, language and/or cultural barrier
 - b. Multiple bruises in various stages of healing
 - c. Age-inappropriate behavior (e.g. adults who are submissive or fearful, children who act in a sexually inappropriate way)
 - d. Pattern burns, bruises, or scars suggestive of specific weaponry used
 - e. Evidence of medical neglect for injuries or infections
 - f. Unexplained trauma to genitourinary systems or frequent infections to this system
 - g. Evidence of malnourishment and/or serious dental problems
3. Assess physical issues and avoid extensive investigation of the specifics of abuse or maltreatment, but document any statements made spontaneously by patient.
 - a. Avoid asking directed questions of a child.

Treatment and Interventions

1. Address life-threatening issues.
2. Remove the patient to a safe place even if no medical indication for transport.
 - a. Report concerns about potential abuse and/or maltreatment to law enforcement immediately, in accordance with [state law](#), if caregivers are impeding your ability to assess and/or transport patient, or refuse care for the patient.
3. For patients transported, report concerns to hospital and/or law enforcement personnel per mandatory [reporting laws](#).

Patient Safety Considerations

- If no medical emergency exists, the next priority is safe patient disposition/removal from the potentially abusive situation.
- Do not confront suspected perpetrators of abuse and/or maltreatment. This can create an unsafe situation for EMS and for the patient.

Notes and Educational Pearls

Key Considerations

- All states have specific mandatory reporting laws that dictate which specific crimes such as suspected abuse or maltreatment must be reported and to whom they must be reported. It is important to be familiar with the specific laws in your state including specifically who must make disclosures, what the thresholds are for disclosures, and to whom the disclosures must be made.
- Clues to abuse or maltreatment can vary depending on the age group of the patient and on the nature of the abuse. Remember that not all abuse or maltreatment involves physical harm. It is important to realize that the job of EMS is to document their concerns, assess the patient for potentially serious injuries, make sure that their concerns are disclosed to the appropriate legal authorities, and work towards getting the patient into a safe situation. EMS personnel should not take it upon themselves to investigate, interview, or intervene above and beyond those concepts and should leave further intervention to the appropriate law enforcement personnel.
- It is very important to have a high index of suspicion for abuse in children presenting with a Brief Resolved Unexplained Event (BRUE). Of the very serious causes of BRUE, child abuse has been found in as many as 11% of cases. One retrospective review noted that a call to 911 for BRUE was associated with an almost 5 times greater odds of abusive head trauma being diagnosed as the cause of the BRUE, clearly emphasizing the high index of suspicion EMS providers must have when responding to these calls.
- Abuse and maltreatment can happen to patients of all ages.
- Patients may be unwilling or unable to disclose abuse or maltreatment so the responsibility falls on EMS personnel to assess the situation, document appropriately, and take appropriate action to secure a safe place for the patient.
- Document findings by describing what you see and not ascribing possible causes (e.g. "0.5- inch round burn to back" as opposed to "burn consistent with cigarette burn").
- Providers should be knowledgeable about mandatory reporting statutes in their area, especially regarding adults (domestic violence, elder abuse).

Pertinent Assessment Findings

As noted above

Agitated or Violent Patient/Behavioral Emergency

Aliases

Acute psychosis, patient restraint

Patient Care Goals

1. Provision of emergency medical care to the agitated, violent, or uncooperative patient
2. Maximizing and maintaining safety for the patient, EMS personnel, and others

Patient Presentation

Inclusion Criteria

Patients of all ages who are exhibiting agitated, violent, or uncooperative behavior or who are a danger to self or others

Exclusion Criteria

Patients exhibiting agitated or violent behavior due to medical conditions including, but not limited to:

- Head trauma
- Metabolic disorders (e.g. hypoglycemia, hypoxia)

Patient Management

Assessment

1. Note medications/substances on scene that may contribute to the agitation, or may be relevant to the treatment of a contributing medical condition.
2. Maintain and support airway.
3. Note respiratory rate and effort—if possible, monitor pulse oximetry *[EMR-O; EMT-R]* and/or capnography *[Acquisition EMT-O; Interpretation INT-O/PARA-R]*.
4. Assess circulatory status:
 - a. Blood pressure (if possible)
 - b. Pulse rate
 - c. Capillary refill
5. Assess mental status:
 - a. Check blood glucose *[EMR-O; EMT-R]* (if possible).
6. Obtain temperature (if possible).
7. Assess for evidence of traumatic injuries.
8. Use a validated risk assessment tool such as RASS (Richmond Agitation Sedation Score), AMSS (Altered Mental Status Score), or BARS (Behavioral Activity Rating Scale) to risk stratify violent patients to help guide interventions.

Treatment and Interventions

1. Establish patient rapport:
 - a. Attempt verbal reassurance and calm patient prior to use of pharmacologic and/or physical management devices.
 - b. Engage family members/loved ones to encourage patient cooperation if their presence does not exacerbate the patient's agitation.
 - c. Provide continued verbal reassurance and calming of patient following use of chemical/physical management devices.

2. Pharmacologic management
 - a. Notes:
 - i. Selection of medications for pharmacologic management should be based upon the patient's clinical condition, current medications, and allergies in addition to EMS resources and on-line medical control.
 - ii. The medications are annotated to indicate when they are preferred for patients that are particularly high risk for violence as assessed by a validated scale (note that the dosing can be adjusted to achieve different levels of sedation).
 - iii. The ordering of medications below is not intended to indicate a hierarchy/preference of administration.
 - b. Benzodiazepines choices
 - i. Diazepam *[INT-O]*
 - ii. Lorazepam *[INT-O]*
 - iii. Midazolam *[INT-O; PARA-R]*
 - c. Antipsychotics choices
 - i. Haloperidol *[PARA-O]*
 - ii. Olanzapine *[INT-O]*
(Note: Concurrent use of IM/IV benzodiazepines and olanzapine IM is not recommended as fatalities have been reported)
 - iii. Ziprasidone *[PARA-O]*
 - d. Dissociative agents (provide sedation and anesthesia)
 - i. Ketamine (option for high violence risk) *[PARA-O]*
 - e. Antihistamines choice
 - i. Diphenhydramine *[PARA-O]*
3. Physical management devices
 - a. Body
 - i. Stretcher straps should be applied as the standard procedure for all patients during transport.
 - ii. Physical management devices, including stretcher straps, should never restrict the patient's chest wall motion.
 - iii. If necessary, sheets may be used as improvised supplemental stretcher straps. Other forms of improvised physical management devices should be discouraged.
 - iv. Supplemental straps or sheets may be necessary to prevent flexion/extension of torso, hips, legs by being placed around the lower lumbar region, below the buttocks, and over the thighs, knees, and legs.
 - b. Extremities
 - i. Do not use devices that require a key to release them (use soft or leather devices).
 - ii. Secure all four extremities to maximize safety for patient, staff, and others.
 - iii. Secure all extremities to the stationary frame of the stretcher.
 - iv. Do not use multiple knots to secure a device.

Patient Safety Considerations

The management of violent patients requires a constant reevaluation of the risk/benefit balance for the patient and bystanders in order to provide the safest care for all involved. These are complex and high-risk encounters. There is no one size fits all solution for addressing these patients.

1. Don PPE.
2. Do not attempt to enter or control a scene where physical violence or weapons are present.
3. Dispatch law enforcement immediately to secure and maintain scene safety.
4. De-escalate patient agitation. This is imperative in the interest of patient safety as well as for EMS personnel and others on scene.
5. Avoid uncontrolled or poorly controlled patient agitation and physical violence. It can place the patient at risk for sudden cardiopulmonary arrest due to the following etiologies:
 - a. Excited delirium/exhaustive mania: A postmortem diagnosis of exclusion for sudden death thought to result from metabolic acidosis (most likely from lactate) stemming from physical agitation or physical control measures and potentially exacerbated by stimulant drugs (e.g. cocaine) or alcohol withdrawal
 - b. Positional asphyxia: Sudden death from restriction of chest wall movement and/or obstruction of the airway secondary to restricted head or neck positioning resulting in hypercarbia and/or hypoxia
6. Apply a ECG cardiac monitor as soon as possible, particularly when pharmacologic management medications have been administered.
7. Monitor closely all patients who have received pharmacologic management medications.
 - a. Monitor for the development of hypoventilation and over sedation.
 - b. Utilize capnography if available.
8. Monitor closely patients who have received antipsychotic medication for pharmacologic management. Monitor them for the potential development of:
 - a. Dystonic reactions (this can easily be treated with diphenhydramine/benzodiazepines).
 - b. Mydriasis (dilated pupils).
 - c. Ataxia.
 - d. Cessation of perspiration.
 - e. Dry mucous membranes.
 - f. Cardiac arrhythmias (particularly QT prolongation).
9. Place stretcher in sitting position to prevent aspiration. This also reduces the patient's physical strength by placing the abdominal muscles in the flexed position.
10. Physically secure patients who are more physically uncooperative. Secure with one arm above the head and the other arm below the waist, and both lower extremities individually secured.
11. The following techniques should be **expressly prohibited** by EMS providers:
 - a. Securing or transporting in a prone position with or without hands and feet behind the back (i.e., hobbling or "hog-tying")
 - b. "Sandwiching" patients between backboards
 - c. Using techniques that constrict the neck or compromise the airway
 - d. Using weapons as adjuncts in managing a patient
12. Avoid concurrent use of IM/IV benzodiazepines and olanzapine IM (it is not recommended as fatalities have been reported).

Notes and Educational Pearls

Key considerations

- Contact on-line medical control at any time for advice, especially when patient's level of agitation is such that transport may place all parties at risk.
- Avoid transport by air (it is not advised).
- Use stretchers with adequate foam padding, particularly around the head, as it facilitates patient's ability to self-position the head and neck to maintain airway patency.
- Consider the following options for patients with key-locking restraint devices (e.g. handcuffs), applied by another agency (e.g. police):
 - Invite the agency officer (who has the key) to accompany the patient in the ambulance.
 - Collaborate with agency to switch to a different restraint that does not require a key if agency officer is not able to accompany patient in ambulance.

Pertinent assessment findings

Continuous monitoring of:

1. Airway patency.
2. Respiratory status with pulse oximetry and/or capnography.
3. Circulatory status with frequent blood pressure measurements.
4. Mental status and trends in level of patient cooperation.
5. Cardiac status, especially if the patient has received pharmacologic management medication.
6. Extremity perfusion with capillary refill in patients in physical management device.

Anaphylaxis and Allergic Reaction

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

Anaphylactic Shock

Patient Care Goals

1. Provide timely therapy for potentially life-threatening reactions to known or suspected allergens to prevent cardiorespiratory collapse and shock.
2. Provide symptomatic relief for symptoms due to known or suspected allergens.

Patient Presentation

Inclusion Criteria

Patients of all ages with suspected allergic reaction and/or anaphylaxis

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Evaluate for patent airway and presence of oropharyngeal edema.
2. Auscultate for wheezing and assess level of respiratory effort.
3. Assess for adequacy of perfusion.
4. Assess for anaphylaxis:
 - a. Criteria either i or ii
 - i. Acute onset of an illness (minutes to several hours) with simultaneous involvement of the skin or mucosal tissue AND at least
 1. Respiratory compromise (dyspnea, wheeze, cough, stridor)
 2. Reduced BP or associated symptoms (syncope, collapse, incontinence)
 3. Severe gastrointestinal symptoms (e.g., severe crampy abdominal pain, repetitive vomiting), especially after exposure to non-food allergens
 - ii. Acute onset of hypotension, respiratory symptoms or laryngeal involvement after exposure to a known or highly probable allergen

Treatment and Interventions

1. If signs of anaphylaxis, administer epinephrine *[EMR-O; EMT-R]*.
2. For urticaria or pruritus, in addition to epinephrine, also administer diphenhydramine *[PARA-O]*.
 - a. The IV route is preferred for the patient in severe shock.
 - b. As a supplement to diphenhydramine given for urticaria, any H2-blocking antihistamine (e.g. famotidine, cimetidine) can be given IV or PO in conjunction with diphenhydramine.
3. If respiratory distress, in addition to epinephrine, consider administering
 - a. Albuterol *[EMR-O; EMT-R]*.

AND/OR

- b. Epinephrine nebulized *[EMR-O; EMT-R]*.
4. If signs of anaphylaxis persist following the first dose of epinephrine, additional IM epinephrine can be repeated every 5-15 minutes.
5. For signs of hypoperfusion, in addition to epinephrine, consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*) rapidly (over 15 minutes) and repeat as needed for ongoing hypoperfusion.
6. When cardiovascular collapse (hypotension with altered mental status, pallor, diaphoresis and/or delayed capillary refill) is present despite repeated IM doses of epinephrine in conjunction with at least 60 mL/kg isotonic fluid boluses, consider an epinephrine infusion *[PARA-R]*
7. Transport as soon as possible, and perform ongoing assessment as indicated.

Patient Safety Considerations

- Time to epinephrine delivery
- Concentration of epinephrine in relation to route
- Weight-based dosing of medications

Notes and Educational Pearls

Key Considerations

1. Allergic reactions and anaphylaxis are serious and potentially life-threatening medical emergencies. It is the body's adverse reaction to a foreign protein (e.g. food, medicine, pollen, insect sting or any ingested, inhaled, or injected substance). A localized allergic reaction (e.g. urticaria or angioedema that does not compromise the airway) may be treated with antihistamine therapy. When anaphylaxis is suspected, EMS personnel should always consider epinephrine as first-line treatment. Cardiovascular collapse may occur abruptly, without the prior development of skin or respiratory symptoms. Constant monitoring of the patient's airway and breathing is essential.
2. Contrary to common belief that all cases of anaphylaxis present with cutaneous manifestations, such as urticaria or mucocutaneous swelling, a significant portion of anaphylactic episodes may not involve these signs and symptoms on initial presentation. Moreover, most fatal reactions to food-induced anaphylaxis in children were not associated with cutaneous manifestations.
3. A thorough assessment and a high index of suspicion are required for all potential allergic reaction patients. Consider:
 - a. History of Present Illness
 - i. Onset and location
 - ii. Insect sting or bite
 - iii. Food allergy or exposure
 - iv. New clothing, soap, detergent
 - v. Past history of reactions
 - vi. Medication history
 - b. Signs and symptoms
 - i. Itching or urticaria
 - ii. Coughing, wheezing, or respiratory distress
 - iii. Chest tightness or throat constriction
 - iv. Hypotension or shock
 - v. Persistent gastrointestinal symptoms (nausea, vomiting, and diarrhea)

- vi. Altered mental status
- c. Other considerations
 - i. Angioedema (drug-induced)
 - ii. Aspiration or airway obstruction
 - iii. Vasovagal event
 - iv. Asthma or COPD
 - v. Heart failure
- 4. Gastrointestinal symptoms occur most commonly in food-induced anaphylaxis, but can occur with other causes.
 - a. Oral pruritus is often the first symptom observed in patients experiencing food-induced anaphylaxis.
 - b. Abdominal cramping is also common, but nausea, vomiting, and diarrhea are frequently observed as well.
- 5. Patients with asthma are at high risk for a severe allergic reaction.
- 6. There is no proven benefit to using steroids in the management of allergic reactions and/or anaphylaxis.
- 7. There is controversy among experts with very low quality evidence to guide management for the use of empiric IM epinephrine after exposure to a known allergen in asymptomatic patients with a history of prior anaphylaxis.

Pertinent Assessment Findings

- Presence or absence of angioedema
- Presence or absence of respiratory compromise
- Presence or absence of circulatory compromise
- Localized or generalized urticaria
- Response to therapy

Altered Mental Status

Aliases

Confusion, altered level of consciousness

Patient Care Goals

1. Identify treatable causes.
2. Protect patient from harm.

Patient Presentation

Inclusion Criteria

Impaired decision-making capacity

Exclusion Criteria

Traumatic brain injury

Patient Management

Assessment

Look for treatable causes of altered mental status:

1. Airway—make sure airway remains patent; reposition patient as needed.
2. Breathing—look for respiratory depression; check SPO₂ [*EMR-O; EMT-R*], ETCO₂, [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] and CO detector [*ALL EMS PRACTICE LEVELS-O*] readings.
3. Circulation—look for signs of shock.
4. Glasgow Coma Score and/or AVPU—document
5. Pupils—document findings
6. Neck—look for rigidity or pain with range of motion.
7. Stroke tool—document findings
8. Blood glucose level [*EMR-O; EMT-R*]— treat if applicable
9. ECG [*Acquisition EMT-O; Interpretation INT-R*]- Arrhythmia limiting perfusion
10. Breath odor—look for possible unusual odors such as alcohol, acidosis, or ammonia.
11. Chest/Abdominal—[insert complete, explicit directive as you have above] intra-thoracic hardware, assist devices, abdominal pain or distention.
12. Extremities/skin—look for track marks, hydration, edema, dialysis shunt; assess temperature to touch (or if able, use a thermometer).
13. Environment—survey for pills, paraphernalia, ambient temperature.

Treatment and Interventions

1. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients [see [Universal Care guideline](#)].
2. Assess blood glucose [*EMR-O; EMT-R*] [see [Hypoglycemia](#) or [Hyperglycemia](#) guidelines]
3. Consider naloxone [*EMR-O; EMT-R*] [see [Opioid Poisoning/Overdose guideline](#)]
4. Consider restraint: physical and chemical [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)]

5. Consider anti-dysrhythmic medication [*INT-O*] [see [Cardiovascular](#) section guidelines for specific dysrhythmia guidelines]
6. Active cooling or warming [see [Hypothermia/Cold Exposure](#) or [Hyperthermia/Heat Exposure](#) guidelines]
7. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]) [see fluid administration doses in [Shock](#) and [Hypoglycemia](#) or [Hyperglycemia](#) guidelines]
8. Vasopressors [*INT-R*] [see [Shock guideline](#)]

Patient Safety Considerations

- With depressed mental status, initial focus is on airway protection, oxygenation, ventilation, and perfusion.
- The violent patient may need pharmacologic and/or physical management to insure proper assessment and treatment.
- Hypoglycemic and hypoxic patients can be irritable and violent [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].

Notes and Educational Pearls

Key Considerations

- History from bystanders
- Age of the patient
- Environment where patient found
- Recent complaints (e.g. headache, chest pain, difficulty breathing, vomiting, fever)
- Pill bottles/medications:
 - Anticoagulants
 - Anti-depressants
 - Narcotic pain relievers
 - Benzodiazepines
- Medical alert tags and accessory medical devices
- Evaluate for reduced PO intake and/or vomiting and/or diarrhea or dehydration as a cause of AMS in the pediatric and geriatric populations
- Medications a child may have access to including but not limited to:
 - Antihypertensives
 - Oral hypoglycemic
 - Opioids
 - Benzodiazepines
 - Antiepileptics

Pertinent Assessment Findings

- Track marks
- Breath odor
- Skin temperature
- Location

Back Pain

Aliases

None

Patient Care Goals

1. Improve patient discomfort.
2. Identify life-threatening causes of back pain.

Patient Presentation

Inclusion Criteria

Back pain or discomfort related to a non-traumatic cause or when pain was due to non-acute trauma (e.g. chronic pain conditions)

Exclusion Criteria

1. Back pain from spinal trauma [see [Trauma](#) guidelines]
2. Back pain due to sickle cell pain crisis [see [Sickle Cell Pain Crisis guideline](#)]
3. Back pain from suspected labor [see [OB/GYN](#) guidelines]

Patient Management

Assessment

1. Perform airway assessment and management, per the [Airway Management guideline](#).
2. Obtain vital signs including pulse, respiratory rate, pulse oximetry [*EMR-O; EMT-R*], and blood pressure.
3. Provide evaluation and management of pain, per the [Pain Management guideline](#).
4. Obtain vascular access [*AEMT-R*] as necessary to provide analgesia and/or fluid resuscitation.
5. Assess for life-threatening causes of back pain, which may include:
 - a. Spinal cord compression (e.g. from spinal epidural abscess, malignancy, spinal epidural hematoma for patients on anticoagulants)
 - i. Urinary and/or bowel incontinence
 - ii. Inability to walk due to weakness
 - iii. New neurologic deficits in extremities
 - iv. Loss of sensation in saddle distribution
 - b. Aortic dissection or ruptured abdominal aortic aneurysm
 - i. Unequal femoral or distal lower extremity pulses
 - ii. "Pulsatile" abdominal mass
 - iii. Associated abdominal pain and/or chest pain
 - iv. Known history of abdominal aortic aneurysm or dissection
 - c. Pyelonephritis
 - i. Fever
 - ii. Nausea, vomiting
 - iii. Urinary frequency/urgency
 - iv. Dysuria
 - v. Hematuria
 - vi. Abdominal pain
 - vii. Costovertebral angle tenderness to percussion

6. Assess for signs of shock. If shock is present, provide treatment per appropriate [Shock guideline](#).
7. Assess for other non-life threatening causes of abdominal pain
 - a. Kidney stone
 - i. Unilateral flank pain
 - ii. Nausea, vomiting
 - iii. Possible hematuria
 - iv. History of kidney stones

Treatment and Interventions

1. Medication administration
 - a. Provide analgesia, per [Pain Management guideline](#).
 - b. Administer antiemetics, per [Nausea-Vomiting guideline](#).
 - c. Provide transport to an appropriate receiving facility—consider specialty destination centers for conditions such as suspected aortic emergency.
 - d. Reassess vital signs and response to therapeutic interventions throughout transport.

Patient Safety Considerations

No recommendations

Notes and Educational Pearls

Key Considerations

- Assess for life-threatening causes of back pain.
- Provide appropriate treatment for pain, vomiting, and shock.
- Consider transport to appropriate specialty center if aortic emergency suspected.
- Consider that back and abdominal pain can often coexist with similar disease processes.
- Identify patients on anticoagulants since they are higher risk for spinal epidural hematoma or retroperitoneal hemorrhage which can present as back pain.
- Identify patients with IVDA history and/or impaired immune system since they are higher risk for spinal epidural abscess.
- Identify patients with a history of cancer or with one suspicious for cancer—spinal metastases can cause spinal cord compression.

Pertinent Assessment Findings

- Midline back tenderness
- Back erythema or swelling
- Motor and/or sensory loss in arms or legs
- Loss of perianal sensation
- Absence of or significant inequality of femoral or distal arterial pulses in lower extremities
- Hyper or hypothermia
- Rectal bleeding or hematemesis

End-of-Life Care/Palliative Care

Aliases

None noted

Patient Care Goals

When providing care for a patient near end-of-life:

1. Provide relief from pain and other distressing symptoms.
2. Affirm dying as a normal process.
3. Integrate psychological and spiritual aspects of patient care.
4. Offer a support system to help the family cope during the patient's illness and in their own bereavement.

Patient Presentation

Inclusion Criteria

Patient enrolled in hospice or palliative care, or who has advance care directives, experiencing complaints related to the illness for which the patient is receiving those services

Exclusion Criteria

Complaints unrelated to the illness for which the patient is receiving those services

Patient Management

Assessment, Treatment, and Interventions

1. Perform general patient management.
2. Determine if the patient is able to communicate and has the capacity to make decisions regarding treatment and transport,
 - a. If yes, consult directly with the patient before treatment and/or transport.
 - b. If no, identify any advanced care planning in place for information relating to advanced care planning and consent for treatment, such as:
 - i. Advanced care directives.
 - ii. MOLST/POLST or similar forms.
 - iii. Guardian, power of attorney, or other accepted health care proxy.
3. Assess pain level; if the patient requires pain relief [see [Pain Management guideline](#)].
4. Assess breathing capacity; if the adult patient is experiencing severe respiratory distress, consider:
 - a. Midazolam *[INT-O; PARA-R]*
OR
 - b. Fentanyl *[INT-O]* or other analgesics
5. Assess for nausea; if the patient has nausea [see [Nausea and Vomiting guideline](#)].
6. Assess for secretions; if the patient has excessive secretions, provide suctioning.
7. Assess anxiety level; if the adult patient is anxious, consider:
 - a. Benzodiazepines *[INT-O]*
OR
 - b. Haldol *[PARA-O]*
OR
 - c. Ziprasidone *[PARA-O]*

8. Assess hydration; if the patient appears dehydrated:
 - a. Encourage PO fluid intake if patient is able to swallow.
 - b. Offer ice chips and swabs soaked in ice water, if available.
 - c. Consider administration of normal saline at 10 to 20 mL/kg IV *[AEMT-R]*.
9. Coordinate with guardian, power of attorney, or other accepted health care proxy, in collaboration with hospice or palliative care provider, if non-transport is considered.

Patient Safety Considerations

- Perform careful and thorough assessments to identify complaints not related to the illness for which the patient is receiving hospice or palliative care.
- Deliver care with the utmost patience and compassion.

Notes and Educational Pearls

Key Considerations

- Social interactions with family may affect end-of-life care.
- Scene safety should be considered when deciding on management.

Pertinent Assessment Findings

1. Vital signs
2. Pain score
3. Neurologic exam
4. Lung sounds

Hyperglycemia

Aliases

Diabetic ketoacidosis (DKA), hyperosmolar hyperglycemic state, hyperosmolar non-ketotic coma, diabetes

Patient Care Goals

1. Limit morbidity from hyperglycemia by:
 - a. Appropriate use of glucose monitoring.
 - b. Appropriate hydration for hyperglycemia.

Patient Presentation

Inclusion Criteria

1. Adult or pediatric patient with altered level of consciousness [see [Altered Mental Status guideline](#)]
2. Adult or pediatric patient with stroke symptoms (e.g. hemiparesis, dysarthria) [see [Suspected Stroke/Transient Ischemic Attack guideline](#)]
3. Adult or pediatric patient with seizure [see [Seizures guideline](#)]
4. Adult or pediatric patient with symptoms of hyperglycemia (e.g. polyuria, polydipsia, weakness, dizziness, abdominal pain, tachypnea)
5. Adult or pediatric patient with history of diabetes and other medical symptoms

Exclusion Criteria

Patient in cardiac arrest

Patient Management

Assessment

1. Monitor blood glucose level [*EMR-O; EMT-R*].
2. Conduct secondary survey pertinent to altered blood glucose level:
 - a. Constitutional: assess for tachycardia, hypotension, and tachypnea.
 - b. Eyes: assess for sunken eyes from dehydration.
 - c. Nose /mouth/ears: assess for dry mucus membranes or tongue bite from seizure.
 - d. Neurologic:
 - i. Assess GCS and mental status.
 - ii. Assess for focal neurologic deficit: motor and sensory.
3. Evaluate for possible concomitant sepsis and septic shock [see [Shock guideline](#)].
4. Obtain 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*] to assess for peaked T waves or other findings consistent with hyperkalemia.

Treatment and Interventions

1. If altered level of consciousness, stroke, or sepsis/septic shock, treat per [Altered Mental Status](#), [Suspected Stroke/Transient Ischemic Attack](#), or [Shock](#) guidelines accordingly.
2. If findings of hyperkalemia are present, administer Normal saline IV fluids 20 ml/kg IV/IO [*AEMT-R*] and consider administration of calcium gluconate (preferred) [*PARA-R*].

3. If findings of hyperkalemia, administer sodium bicarbonate *[PARA-R]* and consider albuterol nebulizer *[INT-R]*.
4. If glucose greater than 250 mg/dL with symptoms of dehydration, vomiting, abdominal pain, or altered level of consciousness:
 - a. Provide volume expansion with normal saline bolus *[AEMT-R]*
 - i. Adult: Normal saline 1 L bolus IV; reassess and rebolus 1L if indicated
 - ii. Pediatric: Normal saline 10 mL/kg bolus IV, reassess, and repeat up to 40 mL/kg total
5. Reassess patient.
 - a. Reassess vital signs, mental status, and signs of dehydration.
 - b. If mental status changes, reassess blood glucose level *[EMR-O; EMT-R]* and provide appropriate treatment if hypoglycemia has developed.
6. Implement disposition.
 - a. Transport to closest appropriate receiving facility.

Patient Safety Considerations

1. Overly aggressive administration of fluid in hyperglycemic patients may cause cerebral edema or dangerous hyponatremia.
 - a. Closely monitor for signs of altered mental status, increased intracranial pressure, and immediately discontinue IV fluids and elevate head of bed if signs of increased ICP develop.
 - b. Reassess and manage airway as needed.
2. Asymptomatic hyperglycemia poses no risk to the patient while inappropriately aggressive interventions to manage blood sugar can harm patients.

Notes and Educational Pearls

Key Considerations

- New onset diabetic ketoacidosis in pediatric patients commonly presents with nausea, vomiting, abdominal pain, and/or urinary frequency.
- Consider causes for hyperglycemia by thinking about the 3 I's:
 - Insulin—this refers to any medication changes for insulin or oral medications including poor compliance or malfunctioning insulin pump.
 - Ischemia—this refers to hyperglycemia sometimes being an indication of physiologic stress in a patient and can be a clue to myocardial ischemia in particular.
 - Infection—underlying infection can cause derangements in glucose control.

Pertinent Assessment Findings

- Concomitant trauma
- Abdominal pain, “fruity breath,” and rapid-deep respirations (Kussmaul’s respiration) may be associated with diabetic ketoacidosis.

Hypoglycemia

Aliases

Diabetic coma, insulin shock

Patient Care Goals

Limit morbidity from hypoglycemia by:

1. Describing appropriate use of glucose monitoring.
2. Treating symptomatic hypoglycemia.

Patient Presentation

Inclusion Criteria

1. Adult or pediatric patient with blood glucose less than 60 mg/dL with symptoms of hypoglycemia
2. Adult or pediatric patient with altered level of consciousness [see [Altered Mental Status guideline](#)]
3. Adult or pediatric patient with stroke symptoms (e.g. hemiparesis, dysarthria) [see [Suspected Stroke/Transient Ischemic Attack guideline](#)]
4. Adult or pediatric patient with seizure [see [Seizures guideline](#)]
5. Adult or pediatric patient with history of diabetes and other medical symptoms
6. Pediatric patient with suspected alcohol ingestion
7. Adult patient who appears to be intoxicated

Exclusion Criteria

Patient in cardiac arrest

Patient Management

Assessment

1. Monitoring:
 - a. Check blood glucose level [*EMR-O; EMT-R*]
2. Secondary survey pertinent to altered blood glucose level:
 - a. [Insert noun, like "Insulin pump" or "Diabetes"]: evaluate for presence of an automated external insulin delivery device (insulin pump).
 - b. Constitutional: assess for tachycardia and hypotension
 - c. Eyes: assess for sunken eyes from dehydration
 - d. Nose /mouth/ears: assess for dry mucus membranes or tongue bite from seizure
 - e. Neurologic:
 - i. Assess GCS and mental status.
 - ii. Assess for focal neurologic deficit: motor and sensory.

Treatment and Interventions

1. Assess state of consciousness; if altered level of consciousness or stroke, treat per [Altered Mental Status](#) or [Suspected Stroke/Transient Ischemic Attack](#) guidelines accordingly.
2. Assess glucose level; if blood glucose is 60 mg/dL or less administer one of the following:
 - a. Conscious patient with a patent airway:
 - i. Glucose, oral [*EMR-O; EMT-R*] (in form of glucose tablets, glucose gel, tube of cake icing, etc.)

- b. Unconscious patient, or patients who are unable to protect their own airway:
 - i. Dextrose IV *[AEMT-R]*—administer in incremental doses until mental status improves or maximum field dosing is reached
 - ii. Glucagon IM/IN *[EMT-O]*
 - iii. Remove or disable insulin pump if above treatments cannot be completed.
 - a. Patients with an insulin pump who are hypoglycemic with associated altered mental status (GCS <15):
 - i. Stop the pump, disconnect or remove at insertion site if patient cannot ingest oral glucose or ALS is not available.
 - ii. Leave the pump connected and running if able to ingest oral glucose or receive ALS interventions.
 2. Reassess patient.
 - a. Reassess vital signs and mental status.
 - b. Repeat check of blood glucose *[EMR-O; EMT-R]* level if previous hypoglycemia and mental status has not returned to normal.
 - i. It is not necessary to repeat blood sugar if mental status has returned to normal.
 - c. If maximal field dosage of dextrose solution does not achieve euglycemia and normalization of mental status:
 - i. Initiate transport to closest appropriate receiving facility for further treatment of refractory hypoglycemia.
 - ii. Evaluate for alternative causes of altered mental status.
 - iii. Continue treatment of hypoglycemia using dextrose solutions as noted above.
 3. Implement disposition.
 - a. If hypoglycemia with continued symptoms, transport to closest appropriate receiving facility.
 - b. If hypoglycemia with seizure transport to the hospital regardless of their mental status and response to therapy
 - c. If hypoglycemia resolves after treatment, consider release without transport **only** if **all** of the following are true:
 - i. Repeat glucose is greater than 80 mg/dL.
 - ii. Patient takes only short-acting insulin or metformin to control diabetes.
 - iii. Patient does not take oral antiglycemics.
 - iv. Patient returns to normal mental status, with no focal neurologic signs/symptoms after receiving glucose/dextrose.
 - v. Patient can promptly obtain and will eat a carbohydrate meal.
 - vi. Patient or legal guardian refuses transport and EMS providers agree transport not indicated.
 - vii. A reliable adult will be staying with patient.
 - viii. No major co-morbid symptoms exist, like chest pain, shortness of breath, seizures, intoxication.
 - ix. A clear cause of the hypoglycemia is identified (e.g. missed meal).

Patient Safety Considerations

1. Dextrose 10% can be safely used in all ages of patient.
2. Dextrose 50% can cause local tissue damage if it extravasates from vein, and may cause hyperglycemia. Dextrose 50% carries risk for little clinical gain. EMS systems may consider carrying no more than 25% concentration of dextrose for

- treating hypoglycemia in adults.
3. For children **less than** 1 years old, dextrose concentration of no more than 25% should be used.
 4. For neonates and infants **less than** 1 month of age, dextrose concentration of no more than 10-12.5% should be used.
 5. Sulfonylureas (e.g. glyburide, glipizide) have long half-lives ranging from 12–60 hours. Patients with corrected hypoglycemia who are taking these agents are at particular risk for recurrent symptoms and frequently require hospital admission.

Notes and Educational Pearls

Desired Dose	Fluid type	mL of fluid
0.5g/kg	25% dextrose	2mL/kg
	12.5% dextrose	4mL/kg
	10% dextrose	5mL/kg
1g/kg	25% dextrose	4mL/kg
	12.5% dextrose	8mL/kg
	10% dextrose	10mL/kg

Key Considerations

- Consider contribution of oral diabetic medications to hypoglycemia.
- Have family/patient turn off insulin pumps, if possible.
- Consider potential for intentional overdose of hypoglycemic agents.
- Avoid overshoot hyperglycemia when correcting hypoglycemia. Administer dextrose- containing IV fluids in small doses until either mental status improves or a maximum field dose is achieved.

Pertinent Assessment Findings

- Concomitant trauma
- Diaphoresis or hypothermia may be associated with hypoglycemia

Nausea-Vomiting

Aliases

Gastroenteritis, emesis

Patient Care Goals

Decrease discomfort secondary to nausea and vomiting.

Patient Presentation

Inclusion Criteria

Currently nauseated and/or vomiting

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Routine patient care (vital signs)
2. History and physical examination focused on potential causes of nausea and vomiting (e.g. gastrointestinal, cardiovascular, gynecologic, hypoglycemia, hyperglycemia)

Treatment and Interventions

1. Administer anti-emetic medication (optional, if available; any that can be given IV can be given IO):
 - a. Ondansetron *[INT-O]*
 - b. Metoclopramide *[PARA-O]*
 - c. Prochlorperazine *[PARA-O]*
 - d. Diphenhydramine *[PARA-O]*
 - e. Isopropyl alcohol *[ALL EMS PRACTICE LEVELS-O]*—allow patient to inhale vapor from isopropyl alcohol wipe 3 times every 15 minutes as tolerated.
1. If signs of hypovolemia, consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
 - a. May repeat as indicated.

Patient Safety Considerations

- For very young pediatric patients, ondansetron *[INT-O]* can be sedating.
- Dystonic and extrapyramidal symptoms are possible side effects of antiemetics—if encountered, consider diphenhydramine *[PARA-O]*.

Notes and Educational Pearls

Key Considerations

- Ondansetron is preferred in children for the treatment of nausea and vomiting.
- Metoclopramide has fewer adverse effects than Prochlorperazine in children.

- Prochlorperazine and metoclopramide (phenothiazines) have an increased risk of dystonic reactions.
 - Some phenothiazines also have an increased risk of respiratory depression when used with other medications that cause respiratory depression, and some phenothiazines can cause neuroleptic malignant syndrome.
 - Prochlorperazine carries a black box warning for children under 2 years old.
- IV form of ondansetron may be given PO in same dose.
- Nausea and vomiting are symptoms of illness—in addition to treating the patient's nausea and vomiting, a thorough history and physical are key to identifying what may be a disease in need of emergent treatment (e.g. bowel obstruction, myocardial infarction, pregnancy).
- While ondansetron has not been adequately studied in pregnancy to determine safety, it remains a treatment option for hyperemesis gravidum in pregnant patient.

Pertinent Assessment Findings

- Vital signs
- Risk factors for heart disease. Obtain ECG if applicable
- Pregnancy status
- Abdominal exam

Pain Management

(Incorporates elements of an evidence-based guideline for prehospital analgesia in trauma created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

Analgesia, pain control, acute pain, acute traumatic pain, acute atraumatic pain

Patient Care Goals

The practice of prehospital emergency medicine requires expertise in a wide variety of pharmacological and non-pharmacological techniques to treat acute pain resulting from myriad injuries and illnesses. Approaches to pain relief must be designed to be safe and effective in the dynamic prehospital environment. The degree of pain and the hemodynamic status of the patient will determine the urgency and extent of analgesic interventions.

Patient Presentation

Inclusion Criteria

Patients who are experiencing pain

Exclusion Criteria

1. Pregnancy with active labor
2. Dental pain
3. Patients with care-plans that prohibit use of parenteral analgesics by EMS
4. Patients with chronic pain who aren't part of a hospice and/or palliative care plan

Patient Management

Assessment, Treatment, and Interventions

1. Determine patient's pain score assessment using standard pain scale.
 - a. Less than 4 yo: observational scale (e.g. Faces, Legs, Arms, Cry, Consolability [FLACC] or Children's Hospital of Eastern Ontario Pain Scale (CHEOPS)
 - b. 4–12 yo: self-report scale (e.g. Wong Baker Faces, Faces Pain Scale [FPS], Faces Pain Scale Revised [FPS-R])
 - c. Greater than 12 yo: self-report scale (numeric rating scale [NRS])
2. Place patient on ECG cardiac monitor ***[Acquisition EMT-O; Interpretation INT-R]*** per patient assessment.
3. Consider use of non-pharmaceutical pain management techniques, if available.
 - a. Placement of the patient in a position of comfort.
 - b. Application of ice packs and/or splints for pain secondary to trauma.
 - c. Use of verbal reassurance to control anxiety.
4. If not improved and patient is experiencing moderate discomfort consider use of analgesics:
 - a. Acetaminophen ***[AEMT-R]***
 - b. Ibuprofen ***[AEMT-O]***
 - c. Fentanyl ***[INT-O]***
 - d. Ketorolac (one-time dose only) ***[AEMT-O]***:
 - e. Morphine sulfate ***[INT-O]***

- f. Ketamine: **[PARA-O]**
- g. Nitrous Oxide **[AEMT-O]**
5. Establish IV/IO of normal saline **[AEMT-R]**.
6. If the patient is experiencing severe pain, administer analgesics.
 - a. Ketorolac **[AEMT-O]**
 - b. Morphine sulfate **[INT-O] OR**
 - c. Fentanyl **[INT-O] OR**
 - d. Hydromorphone: **[PARA-O] OR**
 - e. Ketamine: **[PARA-O]**
7. Consider administration of oral, sublingual, or IV antiemetics to prevent nausea in high risk patients [see [Nausea/Vomiting guideline](#)].
8. If indicated based on pain assessment, and vital signs allow, repeat pain medication administration (excluding ketorolac) after 5 minutes of the previous dose.
9. Transport in position of comfort and reassess as indicated.

Adult nonverbal pain scale University of Rochester Medical Center			
Categories	0	1	2
Face	No particular expression or smile.	Occasional grimace, tearing, frowning, wrinkled forehead.	Frequent grimace, tearing, frowning, wrinkled forehead.
Activity (movement)	Lying quietly, normal position.	Seeking attention through movement or slow, cautious movement.	Restless, excessive activity and/or withdrawal reflexes.
Guarding	Lying quietly, no positioning of hands over areas of body.	Splinting areas of the body, tense.	Rigid, stiff.
Physiology (vital signs)	Stable vital signs	Change in any of the following: * SBP > 20 mm Hg. * HR > 20/minute.	Change in any of the following: * SBP > 30 mm Hg. * HR > 25/minute.
Respiratory	Baseline RR/SpO ₂ Compliant with ventilator	RR > 10 above baseline, or 5% ↓SpO ₂ mild asynchrony with ventilator	RR > 20 above baseline, or 10% ↓SpO ₂ severe asynchrony with ventilator

Abbreviations: HR, heart rate; RR, respiratory rate; SBP, systolic blood pressure; SpO₂, pulse oximetry. Instructions: Each of the 5 categories is scored from 0-2, which results in a total score between 0 and 10. Document total score by adding numbers from each of the 5 categories. Scores of 0-2 indicate no pain, 3-6 moderate pain, and 7-10 severe pain. Document assessment every 4 hours on nursing flow-sheet and complete assessment before and after intervention to maximize patient comfort. Sepsis, hypovolemia, hypoxia need to be excluded before interventions.

© Strong Memorial Hospital, University of Rochester Medical Center, 2004. Used with permission.

From: Odhner M, Wegman D, Freeland N, Ingersoll G. Evaluation of a newly developed non-verbal pain scale (NVPS) for assessment of pain in sedated critically ill patients. Available at: [http://www.aacn.org/AACN/NTIPoster.nsf/vwdoc/2004NTI Posters](http://www.aacn.org/AACN/NTIPoster.nsf/vwdoc/2004NTI%20Posters). Accessed July 18, 2017.

Universal Pain Assessment Tool

Verbal Descriptor Scale											
	0	1	2	3	4	5	6	7	8	9	10
	No Pain	Mild Pain	Moderate Pain		Severe Pain		Very Severe Pain		Excruciating Pain		

Wong - Baker FACE S®						
-----------------------------	--	--	--	--	--	--

Descriptive Scale	Alert Smiling	No Humor Serious, Flat	Furrowed Brow Pursed Lips Breath Holding	Wrinkled Nose Raised Upper Lip Rapid Breathing	Slow Blink Open Mouth	Eyes Closed Moaning Crying
--------------------------	---------------	---------------------------	--	--	--------------------------	----------------------------------

Activity Tolerance Scale	No Pain	Can be Ignored	Interferes with Tasks	Interferes with Concentration	Interferes with Basic Needs	Bed Rest Required
Spanish	Nada de Dolor	Un Poquito de Dolor	Un Dolor Leve	Dolor Fuerte	Dolor Desmesurado o Fuerte	Un Dolor Insoportable

Source: Hybrid of scales by authors. Wong-Baker FACES® Pain Scale Rating license grants this use. Reproduction of the Wong-Baker FACES® material requires licensing at www.wongbakerfaces.org.

Pediatric-Appropriate Pain Assessment Tools

Faces, Legs, Activity, Cry, Consolability (FLACC) Behavioral Scale

Appropriate age for use (per guideline): less than 4 years

Scoring			
Categories	0	1	2
Face	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested	Frequent to constant frown, clenched jaw, quivering chin
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid, or jerking
Cry	No cry (awake or asleep)	Moans or whimpers, occasional complaint	Crying steadily, screams or sobs, frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging, or being talked to, distractible	Difficult to console or comfort
<i>Each of the five categories (F) Face; (L) Legs; (A) Activity; (C) Cry; (C) Consolability is scored from 0-2, which results in a total score between zero and ten.</i>			

© 2002, The Regents of the University of Michigan. All Rights Reserved.

Instructions:

- **Patients who are awake:** Observe for at least 1–2 minutes. Observe legs and body uncovered. Reposition patient or observe activity, assess body for tenseness and tone. Initiate consoling interventions if needed.
- **Patients who are asleep:** Observe for at least 2 minutes or longer. Observe body and legs uncovered. If possible reposition the patient. Touch the body and assess for tenseness and tone.

Face

- Score 0 point if patient has a relaxed face, eye contact and interest in surroundings.
- Score 1 point if patient has a worried look to face, with eyebrows lowered, eyes partially closed, cheeks raised, mouth pursed.

- Score 2 points if patient has deep furrows in the forehead, with closed eyes, open mouth and deep lines around nose and/or lips.

Legs

- Score 0 points if patient has usual tone and motion to limbs (legs and arms).
- Score 1 point if patient has increased tone, rigidity, tense, intermittent flexion or extension of limbs.
- Score 2 points if patient has hyper tonicity, legs pulled tight, exaggerated flexion or extension of limbs, tremors.

Activity

- Score 0 points if patient moves easily and freely, normal activity or restrictions.
- Score 1 point if patient shifts positions, hesitant to move, guarding, tense torso, pressure on body part.
- Score 2 points if patient is in fixed position, rocking, side-to-side head movement, rubbing body part.

Cry

- Score 0 points if patient has no cry or moan awake or asleep.
- Score 1 point if patient has occasional moans, cries, whimpers, sighs.
- Score 2 points if patient has frequent/continuous moans, cries, grunts.

Consolability

- Score 0 points if patient is calm and does not require consoling.
- Score 1 point if patient responds to comfort by touch or talk in ½–1 minute.
- Score 2 points if patient requires constant consoling or is unconsolated after an extended time.

Whenever feasible, behavioral measurement of pain should be used in conjunction with self-report. When self-report is not possible, interpretation of pain behaviors and decision-making regarding treatment of pain requires careful consideration of the context in which the pain behaviors were observed.

Each category is scored on a 0–2 scale, which results in a total score of 0–10.


Assessment of Behavioral Score:

- 0 = Relaxed and comfortable
- 1–3 = Mild discomfort
- 4–6 = Moderate pain
- 7–10 = Severe discomfort/pain

© 2002, The Regents of the University of Michigan. All Rights Reserved.

Source: *The FLACC: A behavioral scale for scoring postoperative pain in young children*, by S Merkel and others, 1997, *Pediatr Nurse* 23(3), p. 293–297.

Faces Pain Scale – Revised (FPS-R)



In the following instructions, say "hurt" or "pain," whichever seems right for a particular child. "These faces show how much something can hurt. This face [point to face on far left] shows no pain. The faces show more and more pain [point to each from left to right] up to this one [point to face on far right]—it shows very much pain. Point to the face that shows how much you hurt [right now]."

Score the chosen face 0, 2, 4, 6, 8, or 10, counting left to right, so "0" = "no pain" and "10" = "very much pain." Do not use words like "happy" or "sad." This scale is intended to measure how children feel inside, not how their face looks.

Permission for Use. Copyright of the FPS-R is held by the International Association for the Study of Pain (IASP) ©2001. This material may be photocopied for non-commercial clinical, educational and research use. For reproduction of the FPS-R in a journal, book or web page, or for any commercial use of the scale, request permission from IASP online at www.iasp-pain.org/FPS-R.

Patient Safety Considerations

All patients should have drug allergies identified prior to administration.

- Administer opioids with caution to patients with GCS less than 15, hypotension, identified medication allergy, hypoxia (oxygen saturation less than 90%) after maximal supplemental oxygen therapy, or signs of hypoventilation.
- Opioids are contraindicated for patients who have taken monoamine oxidase inhibitors (MAOIs – e.g. Nardil®, Parnate®, Azilect®, Marplan®, Eldepryl®) during the previous 14 days.
- Avoid non-steroidal anti-inflammatory medications such as ketorolac in patients with NSAID allergy, aspirin-sensitive asthma, renal insufficiency, pregnancy, or known peptic ulcer disease.
- Ketorolac should not be used in patients with hypotension (due to renal toxicity).
- Use of splinting techniques and application of ice should be done to reduce the total amount of medication used to keep the patient comfortable.

Notes and Educational Pearls

Key Considerations

- Record pain severity (0–10) before and after analgesic medication administration and upon arrival at destination.
- Provide analgesic interventions for patients with acute abdominal pain. Use of analgesics for acute abdominal pain does not mask clinical findings or delay diagnosis.
- Recognize that opiates may cause a rise in intracranial pressure.

Pertinent Assessment Findings

- Mental status (GCS and pain level)
- Respiratory system (tidal volume, chest rigidity)
- Gastrointestinal (assess for tenderness, rebound, guarding, and nausea)

Seizures

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

Status epilepticus, febrile seizure, convulsions, eclampsia

Patient Care Goals

1. Prompt cessation of seizures in the prehospital setting
2. Minimizing adverse events in the treatment of seizures in the prehospital setting
3. Minimizing seizure recurrence during transport

Patient Presentation

Seizures due to trauma, pregnancy, hyperthermia, or toxic exposure should be managed according to those condition-specific guidelines.

Inclusion Criteria

Seizure activity upon arrival of prehospital personnel or new or recurrent seizure activity lasting greater than 5 minutes

Exclusion Criteria

None

Patient Management

Assessment

1. History
 - a. Duration of current seizure
 - b. Prior history of seizures, diabetes, or hypoglycemia
 - c. Typical appearance of seizures
 - d. Baseline seizure frequency and duration
 - e. Focality of onset, direction of eye deviation
 - f. Concurrent symptoms of apnea, cyanosis, vomiting, bowel/bladder incontinence, or fever
 - g. Bystander administration of medications to stop the seizure
 - h. Current medications, including anticonvulsants
 - i. Recent dose changes or non-compliance with anticonvulsants
 - j. History of trauma, pregnancy, heat exposure, or toxin exposure
2. Exam
 - a. Air entry and airway patency
 - b. Breath sounds, respiratory rate and effectiveness of ventilation
 - c. Signs of perfusion (pulses, capillary refill, color)
 - d. Neurologic status (GCS, nystagmus, pupil size, focal neurologic deficit or signs of stroke)

Treatment and Interventions

1. If signs of airway obstruction are present and a chin-lift, jaw thrust, positioning, and/or suctioning does not alleviate it, place oropharyngeal airway (if gag

- reflex is absent) or nasopharyngeal airway.
2. Place pulse oximeter [*EMR-O; EMT-R*] and/or waveform capnography [*Interpretation INT-R*] to monitor oxygenation/ventilation.
 3. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
 4. Assess perfusion.
 5. Assess neurologic status.
 6. Routes for treatment:
 - a. Use IN/IM routes are preferred over rectal (PR), IV, or IO routes, if within the provider's scope of practice
 - i. If none of these routes (IN/IM/IV/IO) of medication administration are in provider's scope of practice, diazepam 0.2 mg/kg PR [*INT-O*] (maximum dose 10 mg) is an acceptable route of administration.
 - b. IV placement is not necessary for treatment of seizures, but could be obtained if needed for other reasons.
 7. Anticonvulsant Treatment midazolam (preferred) [*INT-O*]
 8. Glucometry
 - a. If still actively seizing, check blood glucose level [*EMR-O; EMT-R*].
 - b. If less than 60 mg/dL, treat per the [Hypoglycemia guideline](#).
 8. Consider magnesium sulfate [*PARA-R*] in the presence of seizure in the third trimester of pregnancy or post-partum [see the [Eclampsia/Pre-eclampsia guideline](#)].
 9. For febrile seizures, consider the following interventions after stopping the seizure, since the following interventions provide symptomatic relief for fevers but do not stop the seizure:
 - a. Acetaminophen [*AEMT-R*]
AND/OR
 - b. Ketorolac IV/IO/IM OR Ibuprofen PO (if able to swallow) [*INT-O*]
AND/OR
 - c. Removing excessive layers of clothing
AND/OR
 - d. Applying cool compresses to the body
 10. Consider 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*] following cessation of seizure in patients without a history of seizure to determine possible cardiac cause.

Patient Safety Considerations

- Trained personnel should be able to give medication without contacting on-line medical control, however, more than two doses of benzodiazepines are associated with high risk of airway compromise.
 - Use caution, weigh risks and benefits of deferring treatment until hospital, and/or consider consultation with direct medical oversight if patient has received two doses of benzodiazepines by bystanders and/or prehospital providers.
- Hypoglycemic patients who are treated in the field for seizure should be transported to hospital, regardless of whether or not they return to baseline mental status after treatment.

Notes and Educational Pearls

Key Considerations

- Many airway/breathing issues in seizing patients can be managed without intubation or placement of an advanced airway. Reserve these measures for patients that fail less invasive maneuvers as noted above.
- For children with convulsive status epilepticus requiring medication management in the prehospital setting, trained EMS personnel should be allowed to administer medication without on-line medical control.
- For new onset seizures or seizures that are refractory to treatment, consider other potential causes including, but not limited to, trauma, stroke, electrolyte abnormality, toxic ingestion, pregnancy with eclampsia, hyperthermia.
- A variety of safe and efficacious doses for benzodiazepines have been noted in the literature for seizures.
 - The doses for anticonvulsant treatment noted above are those that are common to the forms and routes of benzodiazepines noted in this guideline.
 - One dose, rather than a range, has been suggested in order to standardize a common dose in situations when an EMS agency may need to switch from one type of benzodiazepine to another due to cost or resource limitations.
- Recent evidence supports the use of midazolam IM as an intervention that is at least as safe and effective as intravenous lorazepam for prehospital seizure cessation.

Pertinent Assessment Findings

The presence of fever with seizure in children less than 6 months old and greater than 6 yo is **not** consistent with a simple febrile seizure, and should prompt evaluation for meningitis, encephalitis or other cause.

Shock

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

None noted

Patient Care Goals

1. Initiate early fluid resuscitation and vasopressors to maintain/restore adequate perfusion to vital organs.
2. Differentiate between possible underlying causes of shock in order to promptly initiate additional therapy.

Patient Presentation

Inclusion Criteria

1. Signs of poor perfusion (due to a medical cause) such as one or more of the following:
 - a. Altered mental status
 - b. Delayed capillary refill
 - c. Hypoxia (pulse oximetry **less than 94%**) [*EMR-O; EMT-R*]
 - d. Decreased urine output
 - e. Respiratory rate greater than 20 in adults or elevated in children (see normal vital signs table)
 - f. Hypotension for age (lowest acceptable systolic blood pressure in mm Hg):
 - i. Less than 1 yo: 60
 - ii. 1–10 yo: (age in years) (2)+70
 - iii. Greater than 10 yo: 90
 - g. Tachycardia for age, out of proportion to temperature [see [Appendix VIII – Abnormal Vital Signs](#)]
 - h. Weak, decreased or bounding pulses
 - i. Cool/mottled or flushed or ruddy skin
2. Potential etiologies of shock:
 - a. Hypovolemia (poor fluid intake, excessive fluid loss (e.g. bleeding, SIADH, hyperglycemia excessive diuretics, vomiting, diarrhea)
 - b. Sepsis
 - i. Temperature instability:
 1. Less than 36°C or 96.8°F
 2. Greater than 38.5°C or 101.3°F**and/or**
 3. Tachycardia, warm skin, tachypnea
 - c. Anaphylaxis (urticaria, nausea/vomiting, facial edema, wheezing,)
 - d. Signs of heart failure (hepatomegaly, rales on pulmonary exam, extremity edema, JVD)

Exclusion Criteria

Shock due to suspected trauma [see [Trauma](#) section guidelines]

Patient Management

Assessment

1. History
 - a. History of GI bleeding
 - b. Cardiac problems
 - c. Stroke
 - d. Fever
 - e. Nausea and/or vomiting, diarrhea
 - f. Frequent or no urination
 - g. Syncopal episode
 - h. Allergic reaction
 - i. Immunocompromise (malignancy, transplant, asplenia)
 - j. Adrenal insufficiency
 - k. Presence of a central line or port
 - l. Other risk of infection (spina bifida or other genitourinary anatomic abnormality)
2. Exam
 - a. Airway and breathing (airway edema, rales, wheezing, pulse oximetry, respiratory rate)
 - b. Circulation (heart rate, blood pressure, capillary refill)
 - c. Abdomen (hepatomegaly)
 - d. Mucous membrane hydration
 - e. Skin (turgor, rash)
 - f. Neurologic (GCS, sensorimotor deficits)
3. Determination of type of shock
 - a. Cardiogenic
 - b. Distributive (neurogenic, septic, anaphylactic)
 - c. Hypovolemic
 - d. Obstructive (e.g. pulmonary embolism, cardiac tamponade, tension pneumothorax)

Treatment and Interventions

1. Check vital signs.
2. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
3. Apply ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]*
4. Pulse oximetry *[EMR-O; EMT-R]* and ETCO₂ *[Acquisition EMT-O; Interpretation INT-O/PARA-R]* (reading of less than 25 mmHg may be sign of poor perfusion)
5. Check blood sugar *[EMR-O; EMT-R]*, and correct if less than 60 mg/dl.
6. 12-Lead ECG *[Acquisition EMT-O; Interpretation INT-R]*
7. Check lactate, if available (greater than 2.0 mmol/L is abnormal).
8. Establish IV/IO access *[AEMT-R]*.
9. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*) over less than 15 minutes, using a push-pull method of drawing up the fluid in a syringe and pushing it through the IV/IO (preferred for pediatric patients). You may repeat up to 3 times based on patient's condition and clinical impression.
10. If there is a history of adrenal insufficiency or long-term steroid dependence, give:
 - a. Hydrocortisone succinate V/IM (preferred) *[PARA-O]*

OR

- b. Methylprednisolone *[PARA-O]*
- 11. Vasopressors (shock unresponsive to IV fluids)
 - i. Cardiogenic shock, hypovolemic shock, obstructive shock and distributive shock, give Norepinephrine infusion (preferred) *[PARA-O]*
- 12. For anaphylactic shock, treat per the [Anaphylaxis and Allergic Reaction guideline](#)
- 13. Provide advanced notification to the hospital
- 14. Antipyretics for fever
 - a. Acetaminophen *[AEMT-R]*
 - b. Ibuprofen *[AEMT-O]*

Patient Safety Considerations

Recognition of cardiogenic shock: If patient condition deteriorates after fluid administration, or rales or hepatomegaly develop, then consider cardiogenic shock and holding further fluid administration.

Notes and Educational Pearls

Key Considerations

- Early, aggressive IV fluid administration is essential in the treatment of suspected shock.
- Patients predisposed to shock include:
 - Immunocompromised (patients undergoing chemotherapy or with a primary or acquired immunodeficiency)
 - Adrenal insufficiency (Addison's disease, congenital adrenal hyperplasia, chronic or recent steroid use)
 - History of a solid organ or bone marrow transplant
 - Infants
 - Elderly
- In most adults, tachycardia is the first sign of compensated shock, and may persist for hours. Tachycardia can be a late sign of shock in children and a tachycardia child may be close to cardiovascular collapse.
- Hypotension indicates uncompensated shock, which may progress to cardiopulmonary failure within minutes.
- Hydrocortisone succinate, if available, is preferred over methylprednisolone and dexamethasone for the patient with adrenal insufficiency, because of its dual glucocorticoid and mineralocorticoid effects.
 - Patients with no reported history of adrenal axis dysfunction may have adrenal suppression due to their acute illness, and hydrocortisone should be considered for any patient showing signs of treatment-resistant shock.
 - Patients with adrenal insufficiency may have an emergency dose of hydrocortisone available that can be administered IV or IM.

Pertinent Assessment Findings

Decreased perfusion manifested by altered mental status, or abnormalities in capillary refill or pulses, decreased urine output (**less than** 1mL/kg/hr):

1. Cardiogenic, hypovolemic, obstructive shock: capillary refill greater than 2 seconds, diminished peripheral pulses, mottled cool extremities
2. Distributive shock: flash capillary refill, bounding peripheral pulses

Sickle Cell Pain Crisis

Aliases

None

Patient Care Goals

1. Identify potentially life-threatening complications of a sickle cell disease.
2. Improve patient comfort.

Patient Presentation

Inclusion Criteria

1. Patient with known sickle cell disease experiencing a pain crisis

Exclusion Criteria

1. Pain due to acute traumatic injury [see [Trauma](#) section guidelines]
2. Abdominal pain due to or related to pregnancy [see [OB/GYN](#) section guidelines]
3. Patients with sickle cell trait

Patient Management

Assessment

1. Perform airway assessment and management per the [Airway Management guideline](#).
2. Obtain vital signs (including pulse, respiratory rate), pulse oximetry [*EMR-O; EMT-R*], and blood pressure
3. Provide evaluation and management of altered mental status per the [Altered Mental Status guideline](#).
4. Provide evaluation and management of pain per the [Pain Management guideline](#).
5. Obtain vascular access as necessary to provide analgesia and/or fluid resuscitation.
6. Assess for potentially serious complications other than pain crisis which may include:
 - a. Acute chest syndrome
 - i. Hypoxia
 - ii. Chest pain
 - iii. Fever
 - b. Stroke [see [Suspected Stroke/Transient Ischemic Attack guideline](#)]
 - i. Focal neurologic deficits
 - c. Meningitis
 - i. Headache
 - ii. Altered mental status
 - iii. Fever
 - d. Septic arthritis
 - i. Severe pain in a single joint
 - ii. Fever
 - e. Splenic sequestration crisis (usually young pediatric patients)
 - i. Abdominal pain, LUQ
 - ii. Splenic enlargement (examine with care)
 - iii. Hypotension, tachycardia
7. Assess for signs of shock—if shock is present, treat per [Shock guideline](#)

Treatment and Interventions

1. Medication Administration:
 - a. Provide analgesia per the [Pain Management guideline](#).
 - b. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients
 - c. Start an IV and provide saline 10ml/kg normal saline bolus (up to 1L) *[AEMT-R]*.
 - d. Provide transport to an appropriate receiving facility.
 - e. Reassess vital signs and response to therapeutic interventions throughout transport.
2. Comfort measures:
 - a. Keep patient warm and dry.
 - b. Transport in a position of comfort unless clinical condition requires otherwise.

Patient Safety Considerations

None recommended

Notes and Educational Pearls

Key Considerations

- Assess for life-threatening complications of sickle cell disease—these patients have significantly higher risk of numerous complications in addition to pain crises.
- Provide appropriate treatment for pain, respiratory distress, and shock.
- These patients may have a higher tolerance to narcotic pain medications if they are taking them on a regular basis.
- These patients will tolerate acute blood loss poorly due to baseline anemia.
- Patients with sickle cell trait can have acute pain crises in extreme conditions (e.g. heat exhaustion, dehydration) and a number of college athlete deaths have been linked to sickle cell trait.

Pertinent Assessment Findings

- Lung exam and assessment of respiratory distress
- Altered mental status
- Focal neurologic deficits
- Inability to move a joint

Resuscitation

Cardiac Arrest: VF,VT, Asystole, and PEA

Aliases

Heart attack, arrest, cardiac arrest

Patient Care Goals

1. Return of spontaneous circulation (ROSC)
2. Preservation of neurologic function
3. High-quality chest compressions with minimal interruption from recognition of cardiac arrest until confirmation of ROSC or field termination of care

Patient Presentation

Inclusion Criteria

Patients with cardiac arrest

Exclusion Criteria

1. Patients suffering cardiac arrest due to severe hypothermia [see [Hypothermia/Cold Exposure guideline](#)]
2. Patients with identifiable Do Not Resuscitate (or equivalent such as POLST) order [see [Do Not Resuscitate Status/Advance Directive/Health Care Power of Attorney \(POA\) Status guideline](#)]
3. Patients in arrest due to traumatic etiology [see [General Trauma Management guideline](#)]


Patient Management

Assessment

1. The patient in cardiac arrest requires a prompt balance of treatment and assessment
2. In cases of cardiac arrest, assessments should be focused and limited to obtaining enough information to reveal the patient is pulseless
3. Once pulselessness is discovered, treatment should be initiated immediately and any further history must be obtained by bystanders while treatment is ongoing

Treatment and Interventions

Figure 4. Adult Cardiac Arrest Algorithm.



2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/hghlights_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

1. Consider reversible causes of cardiac arrest which include the following:
 - a. Hypothermia: Additions to care include attempts at active rewarming [see [Hypothermia/Cold Exposure guideline](#)].
 - b. The dialysis patient/known hyperkalemic patient: Additions to care include the following:
 - i. Calcium gluconate (preferred) [PARA-R]

- ii. Sodium bicarbonate *[PARA-R]*
- c. Tricyclic antidepressant overdose: Additions to care include sodium bicarbonate *[PARA-R]*
- d. Hypovolemia: Additions to care include normal saline 2 L IV *[AEMT-R]* (or 20 mL/kg, repeated up to 3 times for **pediatrics**)
- e. If the patient is intubated at the time of arrest, assess for tension pneumothorax and misplaced ETT
- f. If tension pneumothorax suspected, perform needle decompression *[INT-R]*. Assess ETT, if misplaced, replace ETT
- 2. If at any time during this period of resuscitation the patient regains return of spontaneous circulation, treat per [Adult Post-ROSC Care guideline](#)
- 3. If resuscitation remains ineffective, consider termination of resuscitation [see [Termination of Resuscitative Efforts guideline](#)]

Patient Safety Considerations

- Performing manual chest compressions in a moving vehicle may pose a provider safety concern.
- In addition, manual chest compressions during patient movement are less effective in regards to hands-on time, depth, recoil, and rate.
- Ideally, patients should be resuscitated as close to the scene as operationally possible.
- Risks and benefits should be considered before patient movement in cardiac arrest situations.

Notes and Educational Pearls

Key Considerations

1. Effective chest compressions and defibrillation are the most important therapies to the patient in cardiac arrest. Effective chest compressions are defined as:
 - a. A rate of greater than 100 and less than 120 compressions/minute.
 - b. Depth
 - i. Adults and children—at least 2 inches (5 cm) and less than 2.4 inches (6cm);
 - ii. Infants— or 1.5 inches (4 cm);
 - iii. Adolescents— patients who have entered puberty should receive the same depth of chest compressions as an adult
 - c. Allow for complete chest recoil (avoid leaning).
 - d. Minimize interruptions in compressions.
 - e. Avoid rescuer fatigue by rotating rescuers at least every 2 minutes. Some EMS pit crew approaches use a provider on either side of the chest, alternating compressions every minute or every 100 compressions to avoid fatigue.
2. Avoid excessive ventilation and consider delayed airway management. If no advanced airway, consider:
 - a. Passive ventilation using an NRB with 3–4 cycles of uninterrupted chest compressions (for arrests of suspected cardiac etiology). Consider BVM ventilation or advanced airway after 3–4 cycles.
 - b. BVM ventilation every 10–15 compressions with cycles of uninterrupted chest compressions. Upstroke ventilation between compressions. 30:2 ventilation to compression ratio for adults, and 15:2 for children when 2 rescuers are present.
 - c. If an advanced airway is placed, ventilations should not exceed 10

breaths/minute (1 breath every 6 seconds or 1 breath every 10 compressions) in adults. **Pediatric Consideration:** For children with an advanced airway, 1 breath every 3–5 seconds is recommended (equivalent to 12–20 breaths/minute)

3. Quantitative end-tidal CO₂ should be used to monitor effectiveness of chest compressions.
 - a. If ETCO₂ less than 10 mmHg during the initial phases of resuscitation, attempt to improve chest compression quality.
 - b. Consider additional monitoring with biometric feedback which may improve compliance with suggested [Resuscitation](#) section guidelines.

Chest compressions are usually the most rapidly applied therapy for the patient in cardiac arrest and should be applied as soon as the patient is noted to be pulseless. If the patient is being monitored with pads in place at the time of arrest, immediate defibrillation should take precedence over all other therapies, however, if there is any delay in defibrillation (for instance, in order to place pads), chest compressions should be initiated while the defibrillator is being applied. There is no guidance on how long these initial compressions should be applied; however, it is reasonable to either complete between 30 seconds and 2 minutes of chest compressions in cases of no bystander chest compressions **or** to perform defibrillation as soon as possible after chest compressions initiated in cases of witnessed arrest.

4. Extracorporeal CPR (ECPR) for patients with cardiac arrest may be considered for select cardiac arrest patients for whom the suspected etiology of the cardiac arrest is potentially reversible during a limited period of mechanical cardiorespiratory support; ECPR systems of care should drive this intervention strategy.
5. Chest compressions should be reinitiated immediately after defibrillation as pulses, if present, are often difficult to detect and rhythm and pulse checks interrupt compressions.
6. Continue chest compressions between completion of AED analysis and AED charging.
7. Effectiveness of chest compressions decreases with any movements.
 - a. Patients should therefore be resuscitated as close to the point at which they are first encountered and should only be moved if the conditions on scene are unsafe or do not operationally allow for resuscitation.
 - b. Chest compressions are also less effective in a moving vehicle.
 - c. It is also dangerous to EMS providers, patients, pedestrians, and other motorists to perform chest compressions in a moving ambulance.
 - d. For these reasons and because in most cases the care provided by EMS providers is equivalent to that provided in emergency departments, resuscitation should occur on scene.
8. The maximum setting on the defibrillator should be used for initial and subsequent defibrillation attempts. Defibrillation dosing should follow manufacturer's recommendation in the case of biphasic defibrillators. If the manufacturer's recommendation is unknown, use highest setting possible. In the case of monophasic devices, the setting should be 360 J (or 4 J/kg for children).
9. Place IV or IO access without interrupting chest compressions.
10. Administer epinephrine during the first or second round of compressions.
11. At present, the most effective mechanism of airway management is uncertain due to some systems managing the airway aggressively and others managing the airway with basic measures, and both types of systems finding excellent outcomes. Regardless of the airway management style, consider the following principles:

- a. Airway management should not interrupt chest compressions.
 - b. Carefully follow ventilation rate and prevent hyperventilation.
 - c. Consider limited tidal volumes.
 - d. There is uncertainty regarding the proper goals for oxygenation during resuscitation:
 - i. Current recommendations suggest using the highest flow rate possible through NRB or BVM.
 - ii. This should not be continued into the post-resuscitation phase in which the goal should be an oxygen saturation of greater than 93%.
 - e. **Pediatric Considerations:** Special attention should be applied to the pediatric population and airway management and respiratory support. Given that the most likely cause of cardiac arrest is respiratory, airway management may be considered early in the patient's care.
 - i. However, the order of Circulation-Airway-Breathing is still recommended as the order of priority by the American Heart Association for pediatric resuscitation in order to ensure timely initiation of chest compressions to maintain perfusion, regardless of the underlying cause of the arrest.
 - ii. In addition, conventional CPR is preferred in children, since it is associated with better outcomes when compared to compression-only CPR.
12. Special Circumstances in Cardiac Arrest
- a. Trauma: Treat per the [General Trauma Management guideline](#).
 - b. Pregnancy
 - i. The best hope for fetal survival is maternal survival.
 - ii. Position the patient in the supine position with a second rescuer performing manual uterine displacement to the left in an effort to displace the gravid uterus and increase venous return by avoiding aorto-caval compression.
 - iii. If manual displacement is unsuccessful, the patient may be placed in the left lateral tilt position at 30°. This position is less desirable than the manual uterine displacement as chest compressions are more difficult to perform in this position.
 - iv. Chest compressions should be performed slightly higher on the sternum than in the non-pregnant patient to account for elevation of the diaphragm and abdominal contents in the obviously gravid patient.
 - v. Defibrillation should be performed as in non-pregnant patients.
 - c. Arrests of respiratory etiology (including drowning): In addition to the above, consider early management of the patient's airway. Passive ventilation with a NRB is not indicated for these patients.
13. Application of the "pit crew" model of resuscitation **[ALL EMS PRACTICE LEVELS-O]**
- a. Ideally, providers in each EMS agency will use a "pit crew" approach when using this protocol to ensure the most effective and efficient cardiac arrest care. Training should include teamwork simulations integrating first responders, BLS, and ALS crewmembers who regularly work together. High-performance systems should practice teamwork using "pit crew" techniques with predefined roles and crew resource management principles. For example (the Pennsylvania State EMS Model for Pit Crew):
 - i. Rescuer 1 and 2 set up on opposite sides of patient's chest and perform continuous chest compressions, alternating after every 100 compressions to avoid fatigue.

- ii. Use a metronome or CPR feedback device to ensure that compression rate is 100–120/minute.
 - iii. Chest compressions are only interrupted during rhythm check (AED analysis or manual) and defibrillation shocks. Continue compressions when AED/ defibrillator is charging.
 - iv. Additional rescuer obtains IV/Oaccess and gives epinephrine. For IO access:
 - 1. The proximal humerus is the preferred site for adults.
 - 2. The tibial site is preferred for infants and children.
 - v. During the first four cycles of compressions and defibrillation, (approximately 10 minutes) avoid advanced airway placement.
 - vi. One responding provider assumes code leader position overseeing the entire response.
 - vii. Use a CPR checklist to ensure that all best practices are followed during CPR.
 - b. For efficient “pit crew” style care, the EMS agency medical director should establish the options that will be used by providers functioning within the EMS agency. Options include establishing:
 - i. The airway/ventilation management, if any, that will be used.
 - ii. The initial route of vascular access.
14. The EMS agency must perform a QI review of care and outcome, overseen by the agency medical director, for every patient that receives CPR.
- a. The QI should be coordinated with local receiving hospitals to include hospital admission, discharge, and condition information. This EMS agency QI can be accomplished by participation in an organized cardiac arrest registry.
 - b. The QI should be coordinated with local PSAP or dispatch centers to review opportunities to assure optimal recognition of possible cardiac arrest cases and provision of dispatch-assisted CPR (including hands-only CPR when appropriate).

Adult Post-ROSC (Return of Spontaneous Circulation) Care

Aliases

None noted

Patient Care Goals

Out-of-hospital cardiac arrest in the U.S. has a mortality rate greater than 90% and results in excess of 300,000 deaths per year. Many of those who do survive suffer significant neurologic morbidity. Current research has demonstrated that care of patients with return of spontaneous circulation (ROSC) at specialized centers is associated with both decreased mortality and improved neurologic outcomes.

The goal is to optimize neurologic and other function following a return of spontaneous circulation following resuscitated cardiac arrest.

Patient Presentation

Inclusion Criteria

Patient returned to spontaneous circulation following cardiac arrest resuscitation

Exclusion Criteria

None recommended

Patient Management

Assessment, Treatment, and Interventions

1. Perform general patient management.
2. Support life-threatening problems associated with airway, breathing, and circulation. Monitor closely for reoccurrence of cardiac arrest.
3. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
4. Do not hyperoxygenate.
5. Do not hyperventilate. Maintain a ventilation rate of 6–8 per minute and ETCO₂ of 35–45 *[Acquisition EMT-O; Interpretation INT-O/PARA-R]* mmHg
6. For hypotension (SBP less than 90 mmHg or MAP less than 65):
 - a. Norepinephrine infusion titrated to a MAP greater than 65 mmHg *[PARA-O]*
or
 - b. Epinephrine infusion (if bradycardic) titrated to a MAP greater than 65 mmHg *[PARA-O]*
7. Perform 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
8. Check blood glucose *[EMR-O; EMT-R]*.
 - a. If hypoglycemic, treat per [Hypoglycemia guideline](#).
 - b. If hyperglycemic, notify hospital on arrival.
9. If patient seizes, treat per [Seizures guideline](#).
10. Post-cardiac arrest patients with evidence or interpretation consistent with ST elevation myocardial infarction (STEMI/Acute MI) should be transported to any hospitals which offer percutaneous coronary intervention in their cardiac catheterization laboratory.
11. Consider transport patients to facility which offers specialized post-resuscitative care.

12. Do not allow patient to become hyperthermic.

Patient Safety Considerations

1. Avoid hyperthermia
2. Prehospital initiation of therapeutic hypothermia is not routinely recommended

Notes and Educational Pearls**Key Considerations**

- Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided.
- Most patients, immediately post resuscitation, will require ventilatory assistance.
- The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. A significant percentage of post-ROSC patients will re-arrest.
- A moderate number of post-ROSC patients may have evidence of ST elevation MI on ECG.
- Common causes of post-resuscitation hypotension include hyperventilation, hypovolemia, and pneumothorax.

Pertinent Assessment Findings

Assess post-ROSC rhythm, lung sounds, and for signs of hypoperfusion.

Determination of Death/Withholding Resuscitative Efforts

Aliases

None noted

Patient Care Goals

All clinically dead patients will receive all available resuscitative efforts including cardiopulmonary resuscitation (CPR) unless contraindicated by one of the exceptions defined below.

Patient Presentation

A clinically dead patient is defined as any unresponsive patient found without respirations and without a palpable carotid pulse.

Inclusion/Exclusion Criteria:

1. Resuscitation should be started on all patients who are found apneic and pulseless unless the following conditions exist (does not apply to victims of lightning strikes, drowning, or hypothermia):
 - a. Medical cause or traumatic injury or body condition clearly indicating biological death (irreversible brain death), limited to:
 - i. Decapitation: the complete severing of the head from the remainder of the patient's body
 - ii. Decomposition or putrefaction: the skin is bloated or ruptured, with or without soft tissue sloughed off. The presence of at least one of these signs indicated death occurred at least 24 hours previously.
 - iii. Transection of the torso: the body is completely cut across below the shoulders and above the hips through all major organs and vessels. The spinal column may or may not be severed.
 - iv. Incineration: 90% of body surface area with full thickness burns as exhibited by ash rather than clothing and complete absence of body hair with charred skin.
 - v. Injuries incompatible with life (such as massive crush injury, complete exsanguination, severe displacement of brain matter).
 - vi. Futile and inhuman attempts as determined by agency policy or protocol related to "compelling reasons" for withholding resuscitation.
 - vii. In blunt and penetrating trauma, if the patient is apneic, pulseless, and without other signs of life upon EMS arrival including, but not limited to, spontaneous movement, ECG activity, or pupillary response
 - viii. Nontraumatic arrest with obvious signs of death including dependent lividity or rigor mortis.
- OR**
- a. A valid DNR order (form, card, bracelet) or other actionable medical order (e.g. POLST/ MOLST form) present, when it:
 - i. Conforms to the state specifications for color and construction.
 - ii. Is intact: it has not been cut, broken or shows signs of being repaired.
 - iii. Displays the patient's name and the physician's name.

Patient Management

Assessment

Assess for dependent lividity with rigor mortis and/or other inclusion criteria.

Treatment and Interventions

1. If all the components above are confirmed, no CPR is required.
2. If CPR has been initiated but all the components above have been subsequently confirmed, CPR may be discontinued and on-line medical control contacted as needed.
3. If any of the findings are different than those described above, clinical death is not confirmed and resuscitative measures should be immediately initiated or continued. The [Termination of Resuscitative Efforts guideline](#) should then be implemented.
4. Do Not Resuscitate order (DNR/MOLST/POLST) with signs of life:
 - a. If there is a DNR bracelet or DNR transfer form and there are signs of life (pulse and respirations), provide standard appropriate treatment under existing protocols matching the patient's condition
 - b. To request permission to withhold treatment under these conditions for any reason, obtain on-line medical control.
 - c. If there is documentation of a Do Not Intubate (DNI/MOLST/POLST) advanced directive, the patient should receive full treatment per protocols with the exception of any intervention specifically prohibited in the patient's advanced directive.
 - d. If for any reason an intervention that is prohibited by an advanced directive is being considered, on-line medical control should be obtained.

Patient Safety Considerations

In cases where the patient's status is unclear and the appropriateness of withholding resuscitation efforts is questioned, EMS personnel should initiate CPR immediately and then contact on-line medical control.

Notes and Educational Pearls

Key Considerations

- For scene safety and/or family wishes, provider may decide to implement CPR even if all the criteria for death are met.
- At a likely crime scene, disturb as little potential evidence as possible.

Pertinent Assessment Findings

No recommendations

Do Not Resuscitate Status, Advance Directives, and Health Care Power of Attorney (POA) Status

Aliases

DNR, comfort care

Patient Care Goals

To acknowledge and maintain the variety of ways that patients can express their wishes about cardiopulmonary resuscitation or end of life decision making

Patient Presentation

Inclusion/Exclusion Criteria

1. Patients must have one of the following documents or a valid alternative (such as identification bracelet indicating wishes) immediately available. Note that some specifics can vary widely from state to state:
 - a. Physician Orders for Life Sustaining Treatment (POLST) or Medical Orders for Life Sustaining Treatment (MOLST): explicitly describes acceptable interventions for the patient in the form of medical orders, must be signed by a physician or other empowered medical provider to be valid
 - b. Do Not Resuscitate (DNR) order: identifies that CPR and intubation are not to be initiated if the patient is in arrest or peri-arrest; the interventions covered by this order and the details around when to implement them can vary widely
 - c. Advance directives: document that describes acceptable treatments under a variable number of clinical situations including some or all of the following: what to do for cardiac arrest, whether artificial nutrition is acceptable, organ donation wishes, dialysis, and other parameters; the directives frequently do not apply to emergent or potentially transient medical conditions
 - d. As specified from state to state, in the absence of formal written directions (MOLST, POLST, DNR, advanced directives), and in the presence of a person with power of attorney for health care or health care proxy, that person may prescribe limits of treatment
2. One of the documents above is valid when it meets all of the following criteria:
 - a. Conforms to the state specifications for color and construction
 - b. Is intact: it has not been cut, broken or shows signs of being repaired
 - c. Displays the patient's name and the physician's name
3. If there is question about the validity of the form or instrument, the best course of action is to proceed with the resuscitation until additional information can be obtained to clarify the best course of action
4. If a patient has a valid version of one of the above documents, it will be referred to as a "valid exclusion to resuscitation" for the purposes of this protocol

Patient Management

Assessment

1. If the patient has a valid exclusion to resuscitation then no CPR or airway management should be attempted, however this does not exclude comfort measures including medications for pain as appropriate.
2. If CPR has been initiated and a valid exclusion to resuscitation has been

subsequently verified, CPR may be discontinued and on-line medical control contacted as needed.

Treatment and Interventions

1. If there is a valid exclusion to resuscitation and there are signs of life (pulse and respirations), EMS providers should provide standard appropriate treatment under existing protocols according to the patient's condition.
 - a. If the patient has a MOLST or POLST, it may provide specific guidance on how to proceed in this situation.
 - b. Directives should be followed as closely as possible and on-line medical control contacted as needed.
2. The patient should receive full treatment per protocols with the exception of any intervention specifically prohibited in the patient's valid exclusion to resuscitation.
3. If for any reason an intervention that is prohibited by an advanced directive is being considered, on-line medical control should be obtained.

Patient Safety Considerations

In cases where the patient's status is unclear and the appropriateness of withholding resuscitation efforts is questioned, EMS personnel should initiate CPR immediately and contact direct medical oversight.

Notes and Educational Pearls

Key Considerations

- If there is a personal physician present at the scene who has an ongoing relationship with the patient, that physician may decide if resuscitation is to be initiated.
- If there is a registered nurse from a home health care or hospice agency present at the scene who has an ongoing relationship with the patient, and who is operating under orders from the patient's private physician, that nurse (authorized nurse) may decide if resuscitation is to be initiated.
- If the physician or nurse decides resuscitation is to be initiated, usual on-line medical control will be followed.
- Special Consideration: For scene safety and/or family wishes, provider may decide to implement CPR even if all the criteria for death are met.

Pertinent Assessment Findings

No recommendations

Termination of Resuscitative Efforts

Aliases

Call the code

Patient Care Goals

1. When there is no response to prehospital cardiac arrest treatment, it is acceptable and often preferable to cease futile resuscitation efforts in the field.
2. In patients with cardiac arrest, prehospital resuscitation is initiated with the goal of returning spontaneous circulation before permanent neurologic damage occurs. In most situations, ALS providers are capable of performing an initial resuscitation that is equivalent to an in-hospital resuscitation attempt, and there is usually no additional benefit to emergency department resuscitation in most cases.
3. CPR that is performed during patient packaging and transport is much less effective than CPR done at the scene. Additionally, EMS providers risk physical injury while attempting to perform CPR in a moving ambulance while unrestrained. In addition, continuing resuscitation in futile cases places other motorists and pedestrians at risk, increases the time that EMS crews are not available for another call, impedes emergency department care of other patients, and incurs unnecessary hospital charges. Lastly, return of spontaneous circulation is dependent on a focused, timely resuscitation. The patient in arrest should be treated as expeditiously as possible, including quality, uninterrupted CPR and timely defibrillation as indicated.
4. When cardiac arrest resuscitation becomes futile, the patient's family should become the focus of the EMS providers. Families need to be informed of what is being done, and transporting all cardiac arrest patients to the hospital is not supported by evidence and inconveniences the family by requiring a trip to the hospital where they must begin grieving in an unfamiliar setting. Most families understand the futility of the situation and accept cessations of resuscitation efforts in the field.

Patient Presentation

Patient in cardiac arrest.

Inclusion Criteria

1. Any cardiac arrest patient that has received resuscitation in the field but has not responded to treatment
2. When resuscitation has begun and it is found that the patient has a DNR order or other actionable medical order (e.g. POLST/MOLST form)

Exclusion Criteria

Consider continuing resuscitation for patients in cardiac arrest associated with medical conditions that may have a better outcome despite prolonged resuscitation, including hypothermia (although under certain circumstances, on-line medical control may order termination of resuscitation in these conditions).

Patient Management

Resuscitation may be terminated under the following circumstances:

1. Non-traumatic arrest
 - a. Patient is at least 18 years of age

- b. Patient is in cardiac arrest at the time of arrival of advanced life support
 - i. No pulse
 - ii. No respirations
 - iii. No evidence of meaningful cardiac activity (e.g. asystole or wide complex PEA less than 60 BPM, no heart sounds)
- c. Advanced life support resuscitation is administered appropriate to the presenting and persistent cardiac rhythm.
 - i. Resuscitation may be terminated in asystole and slow wide complex PEA if there is no return of spontaneous circulation after 20 minutes in the absence of hypothermia and the ETCO₂ is less than 20mmHg.
[Acquisition EMT-O; Interpretation INT-O/PARA-R]
 - ii. Narrow complex PEA with a rate above 40 or refractory and recurrent ventricular fibrillation or ventricular tachycardia:
 - 1. Consider resuscitation for up to 60 minutes from the time of dispatch.
 - 2. Termination efforts may be ceased before 60 minutes based on factors including but not limited to ETCO₂ less than 20mmHg
[Acquisition EMT-O; Interpretation INT-O/PARA-R], age, co- morbidities, distance from, and resources available at the closest hospital. Termination before this timeframe should be done in consultation with on-line medical control.
- d. There is no return of spontaneous pulse and no evidence of neurological function (non- reactive pupils, no response to pain, no spontaneous movement).
- e. There is no evidence or suspicion of hypothermia.
- f. All EMS personnel involved in the patient's care agree that discontinuation of the resuscitation is appropriate.

Note: Consider on-line medical control before termination of resuscitative efforts.

- 2. Traumatic arrest
 - a. Patient is at least 18 years of age.
 - b. Resuscitation efforts may be terminated in any blunt trauma patient who, based on thorough primary assessment, is found apneic, pulseless, and asystolic on an ECG or ECG cardiac monitor ***[Acquisition EMT-O; Interpretation INT-R]*** upon arrival of emergency medical services at the scene.
 - c. Victims of penetrating trauma found apneic and pulseless by EMS should be rapidly assessed for the presence of other signs of life, such as pupillary reflexes, spontaneous movement, response to pain, and electrical activity on ECG.
 - i. Resuscitation may be terminated with on-line medical control if these signs of life are absent.
 - ii. If resuscitation is not terminated, transport is indicated.
 - d. Cardiopulmonary arrest patients in whom mechanism of injury does not correlate with clinical condition, suggesting a non-traumatic cause of arrest, should have standard ALS resuscitation initiated.
 - e. All EMS personnel involved in the patient's care agree that discontinuation of the resuscitation is appropriate.

Note: Consider on-line medical control before termination of resuscitative efforts.

Assessment

1. Pulse
2. Respirations
3. Neurologic status assessment [see [Appendix VII](#); purposeful movement, pupillary response]
4. Cardiac activity (including electrocardiography, cardiac auscultation and/or ultrasonography)
5. Quantitative capnography [*Interpretation INT-O/PARA-R*]

Treatment and Interventions

1. Focus on continuous, quality CPR that is initiated as soon as possible.
2. Focus attention on the family and/or bystanders. Explain the rationale for termination.
3. Consider support for family members such as other family, friends, clergy, faith leaders, or chaplains.
4. For patients that are less than 18 yo, consultation with on-line medical control is recommended.

Patient Safety Considerations

All patients who are found in ventricular fibrillation or whose rhythm changes to ventricular fibrillation should in general have full resuscitation continued on scene.

Notes and Educational Pearls

Key Considerations and Pertinent Assessment Findings

1. Recent evidence has shown that, in order to capture over 99% of potential survivors from medical cardiac arrest (especially VF and pulseless VT arrests), resuscitation should be continued for approximately 40 minutes. This does not imply, however, that all resuscitations should continue this long (e.g. asystolic rhythms).
2. In remote or wilderness situations, EMS providers should make every effort to contact on-line medical control, but resuscitation may be terminated in the field without on-line medical control when the following have occurred:
 - a. There has been no return of pulse despite greater than 30 minutes of CPR (this does not apply in the case of hypothermia)
 - b. Transport to an emergency department will take greater than 30 minutes (this does not apply in the case of hypothermia)
 - c. EMS providers are exhausted and it is physically impossible to continue the resuscitation
3. Logistical factors should be considered, such as collapse in a public place, family wishes, and safety of the crew and public.
4. Survival and functional neurologic outcomes are unlikely if ROSC is not obtained by EMS. It is dangerous to crew, pedestrians, and other motorists to attempt to resuscitate a patient during ambulance transport.
5. Quantitative end-tidal carbon dioxide measurements of less than 10 mmHg or falling greater than 25% despite resuscitation indicates a poor prognosis and provide additional support for termination.

Pediatric-Specific Guidelines

Brief Resolved Unexplained Event (BRUE)

Aliases

Apparent Life-Threatening Event, ALTE

Patient Care Goals

1. Recognize patient characteristics and symptoms consistent with a BRUE.
2. Promptly identify and intervene for patients who require escalation of care.
3. Choose proper destination for patient transport.

Patient Presentation

Inclusion Criteria

1. Suspected BRUE: An event in an infant less than 1 yo reported by a bystander as sudden, brief (less than 1 min), completely resolved upon EMS arrival that includes one or more of the following:
 - a. Absent, decreased, or irregular breathing
 - b. Color change (central cyanosis or pallor)
 - c. Marked change in muscle tone (hyper- or hypotonia)
 - d. Altered level of responsiveness

Exclusion Criteria

1. Any of the following present upon EMS evaluation:
 - a. Abnormal vital signs for age (including fever)
 - b. Vomiting
 - c. Signs of trauma
 - d. Noisy breathing
2. Identifiable cause for the event, which may include:
 - a. Gastric reflux (spitting up)
 - b. Swallowing dysfunction
 - c. Nasal congestion
 - d. Periodic breathing of the newborn
 - e. Breath-holding spell
 - f. Change in tone associated with choking, gagging, crying, feeding
 - g. Seizure (eye deviation, nystagmus, tonic-clonic activity)
3. History or exam concerning for child abuse or neglect
4. Color change that involved only redness (e.g. in the face) or isolated perioral or hand/feet cyanosis

Patient Management

Assessment

1. History
 - a. History of circumstances and symptoms before, during, and after the event, including duration, interventions done, and patient color, tone, breathing, feeding, position, location, activity, level of consciousness
 - b. Other concurrent symptoms (fever, congestion, cough, rhinorrhea, vomiting, diarrhea, rash, labored breathing, fussy, less active, poor sleep, poor feeding)
 - c. Prior history of BRUE
 - d. Past medical history (prematurity, prenatal and/or birth complications, gastric reflux, congenital heart disease, developmental delay, airway abnormalities, breathing problems, prior hospitalizations, surgeries, or injuries)
 - e. Family history of sudden unexplained death or cardiac arrhythmia in other children or young adults
 - f. Social history: who lives at home, recent household stressors, exposure to toxins, drugs, or sick contacts
 - g. Considerations for possible child abuse (multiple or changing versions of the story; reported mechanism of injury does not seem plausible, especially for child's developmental stage)
2. Exam
 - a. Full set of vital signs (per [Universal Care guideline](#), includes: T, P, RR, BP, O₂ sat) **[EMR-O; EMT-R]**
 - b. General assessment:
 - i. Signs of respiratory distress (grunting, nasal flaring, retracting)
 - ii. Color (pallor, cyanosis, normal)
 - iii. Mental status (alert, tired, lethargic, unresponsive, irritability)
 - a. Head to toe exam, including:
 - i. Physical exam for signs of trauma or neglect.
 - ii. Pupillary response.

Treatment and Interventions

1. Monitoring
 - a. ECG cardiac monitor **[Acquisition EMT-O; Interpretation INT-O/PARA-R]**
 - b. Continuous pulse oximetry **[EMR-O; EMT-R]**
 - c. Blood glucose check **[EMR-O; EMT-R]**
 - d. Serial observations during transport for change in condition
2. Airway
 - a. Administer oxygen **[EMR-O; EMT-R]** as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients. Escalate from a nasal cannula to a simple face mask to a non-rebreather mask as needed [see [Airway Management guideline](#)].
 - b. Suction the nose and/or mouth (via bulb, suction catheter) if excessive secretions are present.
3. Utility of IV placement and fluids
 - a. Routine IVs should not be placed on all BRUE patients.
 - b. IVs should only be placed in children for clinical concerns of shock, or when administering IV medications **[AEMT-R]**.

Patient Safety Considerations

1. Regardless of patient appearance, all patients with a history of signs or symptoms of BRUE should be transported for further evaluation.
2. Destination considerations
 - a. Consider transport to a facility with pediatric critical care capability for patients with high risk criteria present:
 - i. Less than 2 months of age
 - ii. History of prematurity (less than or equal to 32 weeks gestation or corrected gestational age less than or equal to 45 weeks)
 - iii. More than 1 BRUE, now or in the past
 - b. All patients should be transported to facilities with baseline readiness to care for children.

Notes and Educational Pearls**Key Considerations**

- BRUE is a group of symptoms, not a disease process.
- High risk BRUE patients may require ED or hospital intervention.
- All patients should be transported to an ED.
- Contact on-line medical control if parent or guardian is refusing medical care and/or transport, especially if any high-risk criteria are present (see above).

Pediatric Respiratory Distress (Bronchiolitis)

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

None noted

Patient Care Goals

1. Alleviate respiratory distress.
2. Promptly identify respiratory distress, failure, and/or arrest, and intervene for patients who require escalation of therapy.
3. Deliver appropriate therapy by differentiating other causes of pediatric respiratory distress.

Patient Presentation

Inclusion Criteria

Child less than 2 yo typically with diffuse rhonchi or an otherwise undifferentiated illness characterized by rhinorrhea, cough, fever, tachypnea, and/or respiratory distress

Exclusion Criteria

1. Anaphylaxis
2. Croup
3. Epiglottitis
4. Foreign body aspiration
5. Submersion
6. Drowning
7. Asthma

Patient Management

Assessment

1. History
 - a. Onset of symptoms
 - b. Concurrent symptoms (e.g. fever, cough, rhinorrhea, tongue/lip swelling, rash, labored breathing, foreign body aspiration)
 - c. Sick contacts
 - d. History of wheezing
 - e. Treatments given
 - f. Number of emergency department visits in the past year
 - g. Number of admissions in the past year
 - h. Number of ICU admissions ever
 - i. History of prematurity
 - j. Family history of asthma, eczema, or allergies
2. Exam
 - a. Full set of vital signs (T, BP, RR, P, O₂ saturation [*EMR-O; EMT-R*])
 - b. Air entry (normal vs. diminished)
 - c. Breath sounds (wheezes, crackles, rales, rhonchi, diminished, clear)

- d. Signs of distress (grunting, nasal flaring, retracting, stridor)
- e. Weak cry or inability to speak full sentences (sign of shortness of breath)
- f. Color (pallor, cyanosis, normal)
- g. Mental status (alert, tired, lethargic, unresponsive)
- h. Hydration status (+/- sunken eyes, delayed capillary refill, mucus membranes moist vs. tacky, fontanel flat vs. sunken)

Treatment and Interventions

1. Pulse oximetry [*EMR-O; EMT-R*] and end-tidal CO₂ (ETCO₂) [*Acquisition EMT-O; Interpretation INT-O/PARA-R*]: Use routinely as an adjunct to other forms of respiratory monitoring.
2. ECG [*Acquisition EMT-O; Interpretation INT-R*]: Perform only if there are no signs of clinical improvement after treating respiratory distress.
3. Airway
 - a. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients. Escalate from a nasal cannula to a simple face mask to a non-breather mask as needed, in order to maintain normal oxygenation.
 - b. Suction the nose and/or mouth (via bulb, Yankauer®, or suction catheter) if excessive secretions are present.
4. Inhaled medications: Administer nebulized epinephrine to children in severe respiratory distress with bronchiolitis (e.g. coarse breath sounds) in the prehospital setting if other treatments (e.g. suctioning, oxygen) fail to result in clinical improvement.
5. Utility of IV placement and fluids: Place IVs only in children with respiratory distress for clinical concerns of dehydration, or when administering IV medications [*AEMT-R*].
6. Steroids: Are generally not efficacious, and not given in the prehospital setting.
7. Improvement of oxygenation and/or respiratory distress with non-invasive positive pressure ventilation [*EMT-O; AEMT-R*] adjuncts:
 - a. Administer continuous positive airway pressure (CPAP), Bi-level positive airway pressure (BiPAP), or high flow nasal cannula (HFNC), when available, for severe respiratory distress.
 - b. Utilize bag-valve-mask ventilation in children with respiratory failure.
8. Supraglottic devices and intubation:
 - a. Utilize non-visualized airways [*EMR-O EMT-R*] and intubation [*INT-O; PARA-R*] only if bag-valve-mask ventilation fails.
 - b. Manage the airway in the least invasive way possible.

Patient Safety Considerations

Routine use of lights and sirens is not recommended during transport.

Notes and Educational Pearls

Key Considerations

- Suctioning can be a very effective intervention to alleviate distress, since infants are obligate nose breathers.
- Insufficient data exist to recommend the use of inhaled steam or nebulized saline.
- Though albuterol has previously been a consideration, the most recent evidence does not demonstrate a benefit in using it for bronchiolitis.

- Ipratropium and other anticholinergic agents should not be given to children with bronchiolitis in the prehospital setting.

Pertinent Assessment Findings

Frequent reassessment is necessary to determine if interventions have alleviated signs of respiratory distress or not.

Pediatric Respiratory Distress (Croup)

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

None noted

Patient Care Goals

1. Alleviate respiratory distress.
2. Promptly identify respiratory distress, respiratory failure, and respiratory arrest, and intervene for patients who require escalation of therapy.
3. Deliver appropriate therapy by differentiating other causes of pediatric respiratory distress.

Patient Presentation

Inclusion Criteria

Suspected croup (history of stridor or history of barking cough)

Exclusion Criteria

Presumed underlying cause that includes one of the following:

- Anaphylaxis
- Asthma
- Bronchiolitis (wheezing **less than 2 yo**)
- Foreign body aspiration
- Submersion
- Drowning
- Epiglottitis

Patient Management

Assessment

1. History
 - a. Onset of symptoms (history of choking)
 - b. Concurrent symptoms (fever, cough, rhinorrhea, tongue or lip swelling, rash, labored breathing, foreign body aspiration)
 - c. Sick contacts
 - d. Treatments given
 - e. Personal history of asthma, wheezing, or croup in past
2. Exam
 - a. Full set of vital signs (T, BP, RR, P, O₂ sat **[EMR-O; EMT-R]**)
 - b. Presence of stridor at rest or when agitated
 - c. Description of cough
 - d. Other signs of distress (grunting, nasal flaring, retracting)
 - e. Color (pallor, cyanosis, normal)
 - f. Mental status (alert, tired, lethargic, unresponsive)

Treatment and Interventions

1. Monitoring
 - a. Use pulse oximetry *[EMR-O; EMT-R]* and end-tidal CO₂ (ETCO₂) *[Acquisition EMT-O; Interpretation INT-O/PARA-R]* routinely as an adjunct to other forms of respiratory monitoring.
 - b. Perform ECG monitor *[Acquisition EMT-O; Interpretation INT-O/PARA-R]* only if there are no signs of clinical improvement after treating respiratory distress.
2. Airway
 - a. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
 - b. Suction the nose and/or mouth (via bulb, Yankauer®, or suction catheter) if excessive secretions are present.
3. Inhaled medications
 - a. Administer epinephrine *[INT-R]* nebulized, to all children with croup, in respiratory distress, with signs of stridor at rest. This medication should be repeated at this dose with unlimited frequency for ongoing distress.
 - b. Note: Humidified oxygen or mist therapy is **not** indicated.
4. Medications: Use dexamethasone for suspected croup *[PARA-O]*.
5. Utility of IV placement and fluids: Place IVs only in children with respiratory distress for clinical concerns of dehydration, or when administering IV medications *[AEMT-R]*.
6. Improvement of oxygenation and/or respiratory distress with non-invasive positive pressure ventilation adjuncts:
 - a. Administer continuous positive airway pressure (CPAP) or Bi-level positive airway pressure *[EMT-O; AEMT-R]* for severe respiratory distress.
 - b. Utilize bag-valve-mask ventilation in children with respiratory failure.
7. Use non-visualized airways *[EMR-O; EMT-R]* and intubation *[INT-O; PARA-R]* only if bag-valve-mask ventilation fails. The airway should be managed in the least invasive way possible.

Patient Safety Considerations

1. Routine use of lights and sirens is not recommended during transport.
2. Patients who receive inhaled epinephrine should be transported to definitive care.

Notes and Educational Pearls

Key Considerations

- Upper airway obstruction can have inspiratory, expiratory, or biphasic stridor.
- Foreign bodies can mimic croup, it is important to ask about a possible choking event.
- Impending respiratory failure is indicated by:
 - Change in mental status such as fatigue and listlessness.
 - Pallor.
 - Dusky appearance.
 - Decreased retractions.
 - Decreased breath sounds with decreasing stridor.
- Without stridor at rest or other evidence of respiratory distress, inhaled

medications may not be necessary.

Pertinent Assessment Findings

- Respiratory distress (retractions, wheezing, stridor)
- Decreased oxygen saturation
- Skin color
- Neurologic status assessment
- Reduction in work of breathing after treatment
- Improved oxygenation after breathing

Neonatal Resuscitation

Aliases

None noted

Patient Care Goals

1. Provide routine care to the newly born infant.
2. Perform a neonatal assessment.
3. Rapidly identify newly born infants requiring resuscitative efforts.
4. Provide appropriate interventions to minimize distress in the newly born infant.
5. Recognize the need for additional resources based on patient condition and/or environmental factors.

Patient Presentation

Inclusion Criteria

Newly born infants

Exclusion Criteria

Documented gestational age less than 20 weeks (usually calculated by date of last menstrual period); if any doubt about accuracy of gestational age, initiate resuscitation

Patient Management


Assessment

1. History
 - a. Date and time of birth
 - b. Onset of symptoms
 - c. Prenatal history (prenatal care, substance abuse, multiple gestation, maternal illness)
 - d. Birth history (maternal fever, presence of meconium, prolapsed or nuchal cord, maternal bleeding)
 - e. Estimated gestational age (may be based on last menstrual period)
2. Exam
 - a. Respiratory rate and effort (strong, weak, or absent; regular or irregular)
 - b. Signs of respiratory distress (grunting, nasal flaring, retractions, gasping, apnea)
 - c. Heart rate (fast, slow, or absent)
 - i. Precordium, umbilical stump or brachial pulse may be used
 - ii. Auscultation of chest is preferred since palpation of umbilical stump is less accurate
 - d. Muscle tone (poor or strong)
 - e. Color and appearance (central cyanosis, acrocyanosis, pallor, normal)
 - f. APGAR score (appearance, pulse, grimace, activity, respiratory effort) may be calculated for documentation, but not necessary to guide resuscitative efforts.
 - g. Estimated gestational age (term, late preterm, premature)
 - h. Pulse oximetry **[EMR-O; EMT-R]** should be considered if prolonged resuscitative efforts or if supplemental oxygen **[EMR-O; EMT-R]** is administered. Goal: oxygen saturation at 10 minutes is 85–95%

Treatment and Interventions

1. If immediate resuscitation is required and the newborn is still attached to the mother, clamp the cord in two places and cut between the clamps. If no resuscitation is required, warm, dry, and stimulate the newborn and then cut and clamp the cord after 60 seconds or the cord stops pulsating.
2. Warm, dry, and stimulate
 - a. Wrap infant in dry towel or thermal blanket to keep infant as warm as possible during resuscitation; keep head covered if possible.
 - b. If strong cry, regular respiratory effort, good tone, and term gestation, place infant skin-to-skin with mother and covered with dry linen.
3. If weak cry, signs of respiratory distress, poor tone, or preterm gestation then position airway (sniffing position) and clear airway as needed. If thick meconium or secretions present *and* signs of respiratory distress, suction mouth then nose.
4. If heart rate greater than 100 beats per minute
 - a. Monitor for central cyanosis; provide blow-by oxygen as needed **[EMR-O; EMT-R]**.
 - b. Monitor for signs of respiratory distress. If apneic or in significant respiratory distress:
 - i. Initiate bag-valve-mask ventilation with room air at 40–60 breaths per minute.
 - ii. Consider Non-visualized airway **[EMR-O; EMT-R]** *or* endotracheal intubation if bag-valve mask ventilation is ineffective.
5. If heart rate less than 100 beats per minute
 - a. Initiate bag-valve-mask ventilation with room air at 40–60 breaths per minute.
 - i. Primary indicator of effective ventilation is improvement in heart rate.
 - ii. Rates and volumes of ventilation required can be variable, only use the minimum necessary rate and volume to achieve chest rise and a change in heart rate.
 - b. If no improvement after 90 seconds, change oxygen delivery to 30% FiO₂ if blender available, otherwise 100% FiO₂ **[EMR-O; EMT-R]** until heart rate normalizes.
 - c. Consider Non-visualized airway **[EMR-O; EMT-R]** *or* endotracheal intubation **[INT-O; PARA-R]** if bag-valve-mask ventilation is ineffective.
6. If heart rate less than 60 beats per minute
 - a. Ensure effective ventilations with supplementary oxygen and adequate chest rise.
 - b. If no improvement after 30 seconds, initiate chest compressions; two-thumb-encircling-hands technique is preferred.
 - c. Coordinate chest compressions with positive pressure ventilation (3:1 ratio, 90 compressions and 30 breaths per minute).
 - d. Consider non-invasive airway **[EMR-O; EMT-R]** *or* endotracheal intubation **[INT-O; PARA-R]**.
 - e. Administer epinephrine **[INT-R]**.
7. Consider checking a blood glucose **[EMR-O; EMT-R]** for ongoing resuscitation, maternal history of diabetes, ill appearing or unable to feed.
8. Administer 20 mL/kg normal saline IV/IO **[AEMT-R]** for signs of shock or post-resuscitative care.

Neonatal Resuscitation Algorithm—2015 Update



Top 10 Take-Home Messages for Neonatal Life Support

1. Newborn resuscitation requires anticipation and preparation by providers who train individually and as teams.
2. Most newly born infants do not require immediate cord clamping or resuscitation and can be evaluated and monitored during skin-to-skin contact with their mothers after birth.
3. Inflation and ventilation of the lungs are the priority in newly born infants who need support after birth.
4. A rise in heart rate is the most important indicator of effective ventilation and response to resuscitative interventions.
5. Pulse oximetry is used to guide oxygen therapy and meet oxygen saturation goals.
6. Chest compressions are provided if there is a poor heart rate response to ventilation after appropriate ventilation corrective steps, which preferably include endotracheal intubation.
7. The heart rate response to chest compressions and medications should be monitored electrocardiographically.

8. If the response to chest compressions is poor, it may be reasonable to provide epinephrine, preferably via the intravenous route.
9. Failure to respond to epinephrine in a newborn with history or examination consistent with blood loss may require volume expansion.
10. If all these steps of resuscitation are effectively completed and there is no heart rate response by 20 minutes, redirection of care should be discussed with the team and family.

2020 American Heart Association. Available at <https://www.ahajournals.org/doi/10.1161/CIR.0000000000000902>. Accessed 11/29/2020

Patient Safety Considerations

1. Hypothermia is common in newborns and worsens outcomes of nearly all post-natal complications.
 - a. Ensure heat retention by drying the infant thoroughly, covering the head, and wrapping the baby in dry cloth.
 - b. When it does not encumber necessary assessment or required interventions, “kangaroo care” (i.e. placing the infant skin-to-skin directly against mother’s chest and wrapping them together) is an effective warming technique.
 - c. Newborn infants are prone to hypothermia which may lead to hypoglycemia, hypoxia and lethargy. Aggressive warming techniques should be initiated including drying, swaddling, and warm blankets covering body and head. Check blood glucose and follow [Hypoglycemia guideline](#) as appropriate.
2. During transport, neonate should be appropriately secured in seat or isolette and mother should be appropriately secured.

Notes and Educational Pearls

Key Considerations

- Approximately 10% of newly born infants require some assistance to begin breathing.
- Deliveries complicated by maternal bleeding (placenta previa, vas previa, or placental abruption) place the infant at risk for hypovolemia secondary to blood loss.
- Low birth weight infants are at high risk for hypothermia due to heat loss.
- If pulse oximetry is used as an adjunct, the preferred placement place of the probe is the right arm, preferably wrist or medial surface of the palm. Normalization of blood oxygen levels (SaO₂ 85-95%) will not be achieved until approximately 10 minutes following birth.
- Both hypoxia and excess oxygen administration can result in harm to the infant. If prolonged oxygen use is required, titrate to maintain an oxygen saturation of 85–95%.
- While not ideal, a larger facemask than indicated for patient size may be used to provide bag-valve-mask ventilation if an appropriately sized mask is not available. Avoid pressure over the eyes as this may result in bradycardia.
- Increase in heart rate is the most reliable indicator of effective resuscitative efforts.
- A multiple gestation delivery may require additional resources and/or providers.
- There is no evidence to support the routine practice of administering sodium bicarbonate for the resuscitation of newborns.

Pertinent Assessment Findings

- It is difficult to determine gestational age in the field; if there is any doubt as to viability, resuscitation efforts should be initiated.
- Acrocyanosis, a blue discoloration of the distal extremities, is a common finding in the newly born infant transitioning to extra uterine life; this must be differentiated from central cyanosis.

Aliases

Cardiac arrest

Patient Care Goals

- Return of spontaneous circulation (ROSC)
- Preservation of neurologic function
- High-quality chest compressions with minimal interruption from recognition of cardiac arrest until confirmation of ROSC or field termination of care

Patient Presentation

Inclusion Criteria

Patients with cardiac arrest

Exclusion Criteria


- Patients suffering cardiac arrest due to severe hypothermia [see [Hypothermia/Cold Exposure guideline](#)]
- Patients with identifiable Do Not Resuscitate (or equivalent such as POLST) order [see [Do Not Resuscitate Status/Advance Directive/Health Care Power of Attorney \(POA\) Status guideline](#)]
- Patients in arrest due to traumatic etiology [see [General Trauma Management guideline](#)]

Patient Management

Assessment

1. The patient in cardiac arrest requires a prompt balance of treatment and assessment.
2. In cases of cardiac arrest, assessments should be focused and limited to obtaining enough information to reveal the patient is pulseless.
3. Once pulselessness is discovered, treatment should be initiated immediately and any further history must be obtained by bystanders while treatment is ongoing.

Figure 11. Pediatric Cardiac Arrest Algorithm.



OB/GYN

Childbirth

Aliases

Labor, delivery, birth

Patient Care Goals

1. Recognize imminent birth.
2. Assist with uncomplicated delivery of term newborn.
3. Recognize complicated delivery situations.
4. Apply appropriate techniques when delivery complication exists.

Patient Presentation

Inclusion Criteria

Imminent delivery with crowning

Exclusion Criteria

1. Vaginal bleeding in any stage of pregnancy [see [Obstetrical/Gynecological Conditions guideline](#)]
2. Emergencies in first or second trimester of pregnancy [see [Obstetrical/Gynecological Conditions guideline](#)]
3. Seizure from eclampsia [see [Obstetrical/Gynecological Conditions](#) and [Eclampsia/Pre- Eclampsia](#) guidelines]

Patient Management

Assessment:

Signs of imminent delivery:

- Contractions
- Crowning
- Urge to push
- Urge to move bowels
- Membrane rupture
- Bloody show

Treatment and Interventions

1. If patient in labor but no signs of impending delivery, transport to appropriate receiving facility.
2. Conduct controlled delivery should so as to allow a slow controlled delivery of infant. This will prevent injury to mother.
 - a. Support the infant's head as needed.
3. Check for cord around the baby's neck.
 - a. If present, slip it over the head.
 - b. If unable to free the cord from the neck, double clamp the cord and cut between the clamps.
4. Do not routinely suction the infant's airway (even with a bulb syringe) during delivery.

5. Grasp the head with hand over the ears, and gently guide head down to allow delivery of the anterior shoulder.
6. Gently guide the head up to allow delivery of the posterior shoulder.
7. Slowly deliver the remainder of the infant.
8. After 1–3 minutes, clamp cord about 6 inches from the abdomen with 2 clamps; cut the cord between the clamps.
 - a. If resuscitation is needed, clamp cord and cut as soon as possible.
9. Record APGAR scores at 1 and 5 minutes.
 - a. After delivery of infant, suctioning (including suctioning with a bulb syringe) should be reserved for infants who have obvious obstruction to the airway or require positive pressure ventilation (follow [Neonatal Resuscitation guideline](#) for further care of the infant).
10. Dry and warm infant, wrap in towel and place on maternal chest unless resuscitation needed.
11. The placenta will deliver spontaneously, often within 5–15 minutes of the infant.
 - a. Do not force the placenta to deliver; do not pull on umbilical cord.
 - b. Contain all tissue in plastic bag and transport.
12. After delivery, massaging the uterus and allowing the infant to nurse will promote uterine contraction and help control bleeding.
 - a. Estimate maternal blood loss.
 - b. Treat for hypovolemia as needed.
13. Transport infant secured in seat or isolette unless resuscitation needed.
14. Keep infant warm during transport.
15. Most deliveries proceed without complications. If complications of delivery occur, the following are recommended:
 - a. Shoulder dystocia: If delivery fails to progress after head delivers, quickly attempt the following:
 - i. Hyperflex mother's hips to severe supine knee-chest position.
 - ii. Apply firm suprapubic pressure to attempt to dislodge shoulder.
 - iii. Apply high-flow oxygen to mother.
 - iv. Transport as soon as possible.
 - v. Contact on-line medical control and/or closest appropriate receiving facility for consultation and to prepare team.
 - b. Prolapsed umbilical cord
 - i. Placed gloved hand into vagina and gently lift head and body off of cord.
 1. Assess for pulsations in cord.
 2. Maintain until relieved by hospital staff.
 - ii. Consider placing mother in prone knee-chest position or extreme Trendelenburg.
 - iii. Apply high-flow oxygen to mother.
 - iv. Transport as soon as possible.
 - v. Contact on-line medical control as needed.
 - c. Breech birth
 - i. Place mother supine, allow the buttocks and trunk to deliver spontaneously, then support the body while the head is delivered.
 - ii. If head fails to deliver, place gloved hand into vagina with fingers between infant's face and uterine wall to create an open airway.

- iii. Apply high-flow oxygen to mother.
 - iv. Transport as soon as possible.
 - v. Contact on-line medical control and/or closest appropriate receiving facility for medical consultation and to prepare team.
 - vi. The presentation of an arm or leg through the vagina is an indication for immediate transport to hospital.
 - vii. Assess for presence of prolapsed cord and treat as above.
- d. Excessive bleeding during active labor may occur with placenta previa
 - i. Obtain history from patient.
 - ii. Placenta previa may prevent delivery of infant vaginally.
 - iii. C-Section needed: Transport urgently.
- e. Maternal cardiac arrest
 - i. Apply manual pressure to displace uterus from right to left.
 - ii. Treat per the [Cardiac Arrest guideline](#) for resuscitation care (defibrillation and medications should be given for same indications and doses as if non-pregnant patient).
 - iii. Transport as soon as possible if infant is estimated to be over 24 weeks gestation (perimortem Cesarean section at receiving facility is most successful if done within 5 minutes of maternal cardiac arrest).
 - iv. Contact on-line medical control.

Patient Safety Considerations

1. Supine Hypotension Syndrome:
 - a. If mother has hypotension before delivery, place patient in left lateral recumbent position or manually displace gravid uterus to the left if supine position necessary.
 - b. Knee-chest position may create safety issues during rapid ambulance transport.
2. Do not routinely suction the infant's airway (even with a bulb syringe) during delivery.
3. Newborns are very slippery, take care not to drop the infant.
4. Do not pull on the umbilical cord while the placenta is delivering.
5. If possible, transport between deliveries if mother is expecting twins.

Notes and Educational Pearls

- OB assessment:
 - Length of pregnancy
 - Number of pregnancies
 - Number of viable births
 - Number of non-viable births
 - Last menstrual period
 - Due date (gestational age)
 - Prenatal care
 - Number of expected babies (multiple gestations)
 - Drug use and maternal medication use
- Notify on-line medical control if:
 - Prepartum hemorrhage
 - Postpartum hemorrhage
 - Breech presentation
 - Limb presentation

- Nuchal cord (around neck)
- Prolapsed cord
- Some bleeding is normal with any childbirth.
 - Large quantities of blood or free bleeding are abnormal.

APGAR Score

Sign	0	1	2
Appearance	Blue, Pale	Body pink, Extremities blue	Completely pink
Pulse	Absent	Slow (less than 100)	≥ 100
Grimace	No response	Grimace	Cough or Sneeze
Activity	Limp	Some flexion	Active motion of extremities
Respirations	Absent	Slow, Irregular	Good, Crying

Eclampsia/Pre-Eclampsia

Aliases

Pregnant seizures, toxemia of pregnancy

Patient Care Goals

1. Recognize serious conditions associated with pregnancy and hypertension.
2. Prevention of eclampsia-related seizures.
3. Provide adequate treatment for eclampsia-related seizures.

Patient Presentation

Inclusion Criteria

1. Female patient, more than 20-weeks gestation, presenting with hypertension and evidence of end organ dysfunction, including renal insufficiency, liver involvement, neurological, or hematological involvement
2. May occur up to 4-weeks post-partum but is rare after 48 hours post-delivery
3. Severe features of pre-eclampsia include:
 - a. Severe hypertension (SBP **greater than** 160, DBP **greater than** 110).
 - a. Headache.
 - b. Mental confusion.
 - c. Vision changes.
 - d. Right upper quadrant or epigastric pain.
 - e. Pulmonary edema.
4. Eclampsia
 - a. Pre-eclampsia symptoms plus seizures
5. Eclampsia and pre-eclampsia associated with abruptio placenta and fetal loss

Exclusion Criteria

Chronic hypertension without end organ dysfunction

Patient Management

Assessment

1. Obtain history
 - a. Gestational age or recent post-partum
 - b. Symptoms suggestive of end organ involvement such as headache, confusion, visual disturbances, seizure, epigastric pain, right upper quadrant pain, nausea, and vomiting
 - c. Previous history of hypertension or known pre-eclampsia
2. Monitoring
 - a. Vital signs including repeat blood pressures every 10 min
3. Secondary survey pertinent to obstetric issues:
 - a. Constitutional: vital signs, orthostatic vital signs, skin color
 - b. Abdomen: distention, tenderness
 - c. Genitourinary: visible bleeding
 - d. Neurologic: mental status

Treatment and Interventions

1. Severe hypertension (SBP greater than 160 or DBP greater than 110) lasting more than 15 min with associated preeclampsia symptoms **[PARA-O]**
 - a. Labetalol 20mg IV over 2 min
 - i. May repeat every 10 min X 2 for persistent severe hypertension with preeclampsia symptoms
 - ii. Goal is to reduce MAP by 20–25% initially
 - iii. Ensure that HR is **greater than** 60 bpm prior to administration.
 - OR**
 - b. Hydralazine 5 mg IV
 - i. May repeat 10mg after 20 min for persistent severe hypertension with preeclampsia symptoms
 - ii. Goal is to reduce MAP by 20–25%
 - OR**
 - c. Nifedipine 10 mg. p.o.
 - i. May repeat 10–20 mg p.o. every 20 minutes X 2 for persistent severe hypertension with pre-eclampsia symptoms
 - ii. Goal is to reduce MAP by 20–25%
 - d. Magnesium sulfate **[PARA-R]**
 - e. Reassess vital signs every 10 min during transport
2. Seizures associated with pregnancy greater than 20-weeks gestation
 - a. Magnesium sulfate **[PARA-R]**
 - b. Benzodiazepine, per [Seizures guideline](#), for active seizure not responding to magnesium
Caution: respiratory depression
3. IV fluids:
 - a. Normal saline **[AEMT-R]** or Lactated Ringers **[AEMT-O]** at KVO rate but restrict maximum rate of fluids to 80 mL/hr
 - OR**
 - b. Saline lock
4. Disposition
 - a. Transport to closest appropriate receiving facility.
 - b. Transport patients in second or third trimester of pregnancy on left side or with uterus manually displaced to left if hypotensive.

Patient Safety Considerations

1. Magnesium toxicity (progression)
 - a. Hypotension followed by
 - b. Loss of deep tendon reflexes followed by
 - c. Somnolence, slurred speech followed by
 - d. Respiratory paralysis followed by
 - e. Cardiac arrest.
2. Treatment of magnesium toxicity
 - a. Stop magnesium drip.
 - b. Give calcium gluconate in cases of pending respiratory arrest **[PARA-R]**.
 - c. Support respiratory effort.

Notes and Educational Pearls

Key Considerations

- Delivery of the placenta is the only definitive management for pre-eclampsia and eclampsia.
- Early treatment of severe pre-eclampsia with magnesium and anti-hypertensive significantly reduces the rate of eclampsia. Use of magnesium encouraged if signs of severe pre-eclampsia present to prevent seizure.

Pertinent Assessment Findings

- Vital signs assessment with repeat blood pressure monitoring before and after treatment
- Assessment of deep tendon reflexes after magnesium therapy
- Examination for end organ involvement
- Evaluation of fundal height

Obstetrical and Gynecological Conditions

Aliases

None noted

Patient Care Goals

1. Recognize serious conditions associated with hemorrhage during pregnancy even when hemorrhage or pregnancy is not apparent (e.g. ectopic pregnancy, abruptio placenta, placenta previa)
2. Provide adequate resuscitation for hypovolemia

Patient Presentation

Inclusion Criteria

1. Female patient with vaginal bleeding in any trimester
2. Female patient with pelvic pain or possible ectopic pregnancy
3. Maternal age at pregnancy may range from 10 to 60 years of age

Exclusion Criteria

1. Childbirth and active labor [see [Childbirth guideline](#)]
2. Post-partum hemorrhage [see [Childbirth guideline](#)]

Differential Diagnosis

1. Abruptio placenta: Occurs in third trimester of pregnancy; placenta prematurely separates from the uterus causing intrauterine bleeding
 - a. Lower abdominal pain and uterine rigidity
 - b. Shock, with minimal or no vaginal bleeding
2. Placenta previa: placenta covers part or all of the cervical opening
 - a. Generally, late second or third trimester
 - b. Painless vaginal bleeding, unless in active labor
 - c. For management during active labor [See [Childbirth guideline](#)]
3. Ectopic pregnancy (ruptured)
 - a. First trimester
 - b. Abdominal/pelvic pain with or without minimal bleeding.
4. Spontaneous abortion (miscarriage)
 - a. Generally first trimester
 - b. Intermittent pelvic pain (uterine contractions) with vaginal bleeding

Patient Management

Assessment

1. Obtain history
 - a. Obstetrical history [see [Childbirth guideline](#)]
 - b. Abdominal pain – onset, duration, quality, radiation, provoking or relieving factors
 - c. Vaginal bleeding – onset, duration, quantity (pads saturated)
 - d. Syncope/lightheadedness
 - e. Nausea/vomiting
 - f. Fever

2. Monitoring
 - a. Monitor ECG if history of syncope or lightheadedness
 - b. Monitor pulse oximetry [*EMR-O*; *EMT-R*] if signs of hypotension or respiratory symptoms
3. Secondary survey pertinent to obstetric issues
 - a. Constitutional: vital signs, orthostatic vital signs, skin color
 - b. Abdomen: distention, tenderness, peritoneal signs
 - c. Genitourinary: visible bleeding
 - d. Neurologic: mental status

Treatment and Interventions

1. If signs of shock or orthostasis:
 - a. Position patient supine and keep patient warm
 - b. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*])
 - c. Reassess vital signs and response to fluid resuscitation
2. Disposition - transport to closest appropriate receiving facility

Patient Safety Considerations

1. Patients in third trimester of pregnancy should be transported on left side or with uterus manually displaced to left if hypotensive
2. Do not place hand/fingers into vagina of bleeding patient except in cases of prolapsed cord or breech birth that is not progressing

Notes and Educational Pearls

Key Considerations

Syncope can be a presenting symptom of hemorrhage from ectopic pregnancy or causes of vaginal bleeding.

Pertinent Assessment Findings

1. Vital signs to assess for signs of shock (e.g. tachycardia, hypotension)
2. Abdominal exam (e.g. distention, rigidity, guarding)
3. If pregnant, evaluate fundal height

Respiratory

Airway Management

Aliases

Asthma, upper airway obstruction, respiratory distress, respiratory failure, hypoxemia, hypoxia, hypoventilation, foreign body aspiration, croup, stridor, tracheitis, epiglottitis

Patient Care Goals

1. Provide effective oxygenation and ventilation
2. Recognize and alleviate respiratory distress
3. Provide necessary interventions quickly and safely to patients with the need for respiratory support
4. Identify a potentially difficult airway in a timely fashion

Patient Presentation

Inclusion Criteria

1. Children and adults with signs of severe respiratory distress/respiratory failure
2. Patients with evidence of hypoxemia or hypoventilation

Exclusion Criteria

1. Patients with tracheostomies
2. Chronically ventilated patients
3. Newborn patients
4. Patients in whom oxygenation and ventilation is adequate with supplemental oxygen alone, via simple nasal cannula or face mask

Patient Management

Assessment

1. History – Assess for:
 - a. Time of onset of symptoms
 - b. Associated symptoms
 - c. History of asthma or other breathing disorders
 - d. Choking or other evidence of upper airway obstruction
 - e. History of trauma
2. Physical Examination – Assess for:
 - a. Shortness of breath
 - b. Abnormal respiratory rate and/or effort
 - c. Use of accessory muscles
 - d. Quality of air exchange, including depth and equality of breath sounds
 - e. Wheezing, rhonchi, rales, or stridor
 - f. Cough
 - g. Abnormal color (cyanosis or pallor)
 - h. Abnormal mental status
 - i. Evidence of hypoxemia
 - j. Signs of a difficult airway (short jaw or limited jaw thrust, small thyromental space, upper airway obstruction, large tongue, obesity, large tonsils, large

neck, craniofacial abnormalities, excessive facial hair)

Treatment and Interventions

1. Non-invasive ventilation techniques
 - a. Maintain airway and Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients
For severe respiratory distress or impending respiratory failure, use Noninvasive positive pressure ventilation devices *[EMT-O; AEMT-R]*- Continuous positive airway pressure (CPAP), Bi-level positive airway pressure (BiPAP) or high flow nasal cannula (HFNC) should be administered
 - b. Use bag-valve mask (BVM) ventilation in the setting of respiratory failure or arrest. Two- person, two-thumbs-up BVM ventilation is more effective than one-person technique and should be used when additional providers are available
2. Oropharyngeal airways (OPA) and nasopharyngeal airways (NPA) - Consider the addition of an OPA and/or NPA to make BVM ventilation more effective, especially in patients with altered mental status
3. Non-visualized airways *[EMR-O; EMT-R]* - Consider the use of a Non-visualized airway if BVM is not effective in maintaining oxygenation and/or ventilation. This is especially important in children since endotracheal intubation is an infrequently performed skill in this age group and has not been shown to improve outcomes
4. Endotracheal intubation
 - a. When less-invasive airway methods are ineffective, use endotracheal intubation *[INT-O; PARA-R]* to maintain oxygenation and/or ventilation
 - b. Other indications may include potential airway obstructions, severe burns, multiple traumatic injuries, altered mental status or loss of normal protective airway reflexes
 - c. Monitor clinical signs, pulse oximetry *[EMR-O; EMT-R]*, cardiac rhythm, blood pressure, and capnography *[Interpretation INT-R]* for the intubated patient
 - d. Video laryngoscopy may enhance intubation success rates, and should be used when available. Consider using a bougie, especially when video laryngoscopy is unavailable and glottic opening is difficult to visualize with direct laryngoscope
5. Post-intubation management
 - a. Confirm placement of Non-visualized airway *[Interpretation EMR-R]* **or** endotracheal intubation with waveform capnography *[Interpretation INT-R]*, absent gastric sounds, and bilateral breath sounds
 - b. Continuously monitor placement with waveform capnography *[Interpretation INT-R]* during treatment and transport
 - c. Continuously secure tube manually until tube secured with tape, twill, or commercial device
 - i. Note measurement of tube at incisors or gum line and monitor frequently for tube movement/displacement
 - ii. Cervical collar *[EMR-O; EMT-R]* and/or cervical immobilization device *[EMR-O; EMT-R]* may help reduce neck movement and risk of tube displacement
 - d. Inflate endotracheal tube cuff with minimum air to seal airway *[INT-*

O; PARA-R] - an ETT cuff manometer can be used to measure and adjust the ETT cuff pressure to a recommended 20 cm H₂O pressure

- e. Ventilation
 - i. Tidal volume
 - 1. Ventilate with minimal volume to see chest rise, approximately 6- 7 mL/kg ideal body weight
 - 2. Over-inflation may be detrimental
 - ii. Rate
 - 1. Adult: 10-12 breaths/minute
 - 2. Child: 20 breaths/minute
 - 3. Infant: 30 breaths/minute
 - iii. Continuously monitor ETCO₂ ***[Interpretation INT-R]*** to maintain ETCO₂ of 35-40 mmHg - in head injury with signs of herniation (unilateral dilated pupil or decerebrate posturing), modestly hyperventilate to ETCO₂ 30 mmHg
- f. Consider sedation with sedative or opioid medications if agitated
- 6. Gastric decompression ***[OPTIONAL ALL EMS PRACTICE LEVELS; PARA-R]*** may improve oxygenation and ventilation, so it should be considered when there is obvious gastric distention
- 7. When patients cannot be oxygenated/ventilated effectively by previously mentioned interventions, the provider should consider cricothyroidotomy ***[PARA-O]*** if the risk of death for not escalating airway management seems to outweigh the risk of a procedural complication
- 8. Transport to the closest appropriate hospital for airway stabilization when respiratory failure cannot be successfully managed in the prehospital setting

Patient Safety Considerations

- 1. Avoid excessive pressures or volumes during BVM
- 2. Avoid endotracheal intubation, unless less invasive methods fail, since it can be associated with aspiration, oral trauma, worsening of cervical spine injury, malposition of the ET tube (right main stem intubation, esophageal intubation), or adverse effects of sedation, especially in children
- 3. Once a successful Non-visualized airway ***[EMR-O; EMT-R]*** placement or intubation has been performed, obstruction or displacement of the tube can have further deleterious effects on patient outcome
 - a. Tubes should be secured with either a commercial tube holder or tape
- 4. Providers who do not routinely use medications for rapid sequence intubation (RSI) should not use RSI on children, since the loss of airway protection with the use of RSI may increase complications
 - a. RSI should be reserved for specialized providers operating within a comprehensive program with ongoing training and quality assurance measures

Notes and Educational Pearls

Key Considerations

- 1. When compared to the management of adults with cardiac arrest, paramedics are less likely to attempt endotracheal intubation in children with cardiac arrest. Further, paramedics are more likely to be unsuccessful when intubating children in cardiac arrest and complications such as malposition of the ET tube or aspiration can be

- nearly three times as common in children as compared to adults.
2. Use continuous waveform capnography to detect end-tidal carbon dioxide (ETCO₂). This is an important adjunct in the monitoring of patients with respiratory distress, respiratory failure, and those treated with positive pressure ventilation. It should be used as the standard to confirm SGA, EGD, and endotracheal tube placement.
 3. Non-Invasive Positive Pressure Ventilation
 - a. Contraindications to these non-invasive ventilator techniques include intolerance of the device, severely impaired consciousness, increased secretions inhibiting a proper seal, or recent gastrointestinal and/or airway surgery
 4. Bag-valve-mask :
 - a. Appropriately-sized masks should completely cover the nose and mouth and maintain an effective seal around the cheeks and chin
 - b. Ventilation should be delivered with only sufficient volume to achieve chest rise
 - c. Ventilation rate:
 - i. During CPR, ventilation rate should be 10 breaths per minute, one breath every 10 compressions (or one breath every 6 seconds). When advanced airway is in place, ideally ventilations should be on upstroke between two chest compressions
 - ii. In adults who are not in cardiac arrest, ventilate at rate of 12 breaths per minute
 - iii. In children, ventilating breaths should be delivered over one second, with a two second pause between breaths (20 breaths/minute) in children
 5. Orotracheal intubation **[INT-O; PARA-R]**
 - a. Endotracheal tube sizes

Age	Size (mm) Uncuffed	Size (mm) Cuffed
Premature	2.5	
Term to 3 months	3.0	
3-7 months	3.5	3.0
7-15 months	4.0	3.5
15-24 months	4.5	3.5
2-15 years	$[\text{age}(\text{yr})/4]+4$	$[\text{age}(\text{yr})/4]+3.5$

- b. Approximate depth of insertion = (3) x (endotracheal tube size)
- c. In addition to preoxygenation, apneic oxygenation (high-flow oxygen by nasal cannula) may prolong the period before hypoxia during an intubation attempt

- d. Positive pressure ventilation after intubation can decrease preload and subsequently lead to hypotension - consider providing vasopressor support for hypotension
- e. Appropriate attention should be paid to adequate preoxygenation to avoid peri- intubation hypoxia and subsequent cardiac arrest
- f. Prompt suctioning of soiled airways before intubation attempt may improve first pass success
- g. Confirm successful placement with waveform capnography [*Interpretation INT-R*]. Less optimal methods of confirmation include bilateral chest rise, bilateral breath sounds, and maintenance of adequate oxygenation. Color change on end-tidal CO₂ is less accurate than clinical assessment, and wave-form capnography is superior. Misting observed in the tube is not a reliable method of confirmation. Visualization with video laryngoscopy, when available, may assist in confirming placement when unclear due to capnography failure or conflicting information.
- h. Ongoing education and hands-on practice is essential to maintain skills. This is especially true for children since pediatric intubation is an infrequently utilized skill for many prehospital providers.
- i. Video laryngoscopy may be helpful, if available, to assist with endotracheal intubation
- 6. Consideration should be made to dispatch the highest-level provider for an EMS system given the potential need for advanced airway placement for patients with severe respiratory distress or failure

Pertinent Assessment Findings

- 1. Ongoing assessment is critical when an airway device is in place
- 2. Acute worsening of respiratory status or evidence of hypoxemia can be secondary to displacement or obstruction of the airway device, pneumothorax or equipment failure

Bronchospasm (due to Asthma and Obstructive Lung Disease)

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

Asthma, respiratory distress, wheezing, respiratory failure, bronchospasm, obstructive lung disease, albuterol, levalbuterol, duoneb, nebulizer, inhaler

Patient Care Goals

1. Alleviate respiratory distress due to bronchospasm.
2. Promptly identify and intervene for patients who require escalation of therapy.
3. Deliver appropriate therapy by differentiating other causes of respiratory distress.

Patient Presentation

Inclusion Criteria

1. Respiratory distress with wheezing or decreased air entry in patients 2 yo or older, presumed to be due to bronchospasm from reactive airway disease, asthma, or obstructive lung disease. These patients may have a history of recurrent wheezing that improves with beta-agonist inhalers or nebulizers such as albuterol or levalbuterol.
 - a. Symptoms and signs may include:
 - i. Wheezing: will have expiratory wheezing unless they are unable to move adequate air to generate wheezes
 - ii. Signs of respiratory infection (e.g. fever, nasal congestion, cough, sore throat)
 - iii. Acute onset after inhaling irritant
 - b. This includes:
 - i. Asthma exacerbation
 - ii. Chronic obstructive pulmonary disease (COPD) exacerbation
 - iii. Wheezing from suspected pulmonary infection (e.g. pneumonia, acute bronchitis)

Exclusion Criteria

1. Respiratory distress due to a presumed underlying cause that includes one of the following:
 - a. Anaphylaxis
 - b. Bronchiolitis (wheezing less than 2 yo)
 - c. Croup
 - d. Epiglottitis
 - e. Foreign body aspiration
 - f. Submersion or drowning
 - g. Congestive heart failure
 - h. Trauma

Patient Management

Assessment

1. History
 - a. Onset of symptoms
 - b. Concurrent symptoms (fever, cough, rhinorrhea, tongue and/or lip swelling, rash, labored breathing, foreign body aspiration)
 - c. Usual triggers of symptoms (cigarette smoke, change in weather, upper respiratory infections)
 - d. Sick contacts
 - e. Treatments given
 - f. Previously intubated
 - g. Number of emergency department visits in the past year
 - h. Number of admissions in the past year
 - i. Number of ICU admissions
 - j. Family history of asthma, eczema, or allergies
2. Exam
 - a. Full set of vital signs (T, BP, RR, P, O₂ sat) - waveform capnography is a useful adjunct and will show a "sharkfin" waveform in the setting of obstructive physiology
 - b. Air entry (normal vs. diminished, prolonged expiratory phase)
 - c. Breath sounds (wheezes, crackles, rales, rhonchi, diminished, clear)
 - d. Signs of distress (grunting, nasal flaring, retracting, stridor)
 - e. Inability to speak full sentences (sign of shortness of breath)
 - f. Color (pallor, cyanosis, normal)
 - g. Mental status (alert, tired, lethargic, unresponsive)
 - h. Signs of distress include:
 - i. Apprehension, anxiety, combativeness
 - ii. Hypoxia (***less than*** 90% oxygen saturation)
 - iii. Intercostal, subcostal, or supraclavicular retractions
 - iv. Nasal flaring
 - v. Cyanosis

Treatment and Interventions

1. Monitoring
 - a. Use pulse oximetry and end-tidal CO₂ (ETCO₂) routinely [***Acquisition EMT-O; Interpretation INT-O/PARA-R***] as an adjunct to other forms of respiratory monitoring.
 - b. Check a 12-Lead ECG [***Acquisition EMT-O; Interpretation INT-R***] *only* if there are no signs of clinical improvement after treating respiratory distress.
2. Airway
 - a. Administer oxygen [***EMR-O; EMT-R***] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
 - b. Suction the nose and/or mouth (via bulb, Yankauer, suction catheter) if excessive secretions are present.
3. Inhaled Medications
 - a. Albuterol [***EMR-O; EMT-R***]: Administer to all patients in respiratory distress with signs of bronchospasm (e.g. known asthmatics, quiet wheezers) either by BLS or ALS providers. Repeat this medication at this dose with unlimited

- frequency for ongoing distress.
- b. Ipratropium nebulized: Administer up to 3 doses, in conjunction with albuterol **[EMT-O]**.
 4. Utility of IV Placement and Fluids: Place IVs when there are clinical concerns of dehydration in order to administer fluids, or when administering IV medications **[AEMT-R]**.
 5. Steroids **[PARA-O]** methylprednisolone or dexamethasone
 6. Magnesium sulfate **[PARA-R]**: Administer for severe bronchoconstriction and concern for impending respiratory failure.
 7. Epinephrine **[EMR-O]**: Administer *only* for impending respiratory failure as adjunctive therapy when there are no clinical signs of improvement.
 8. Improvement of oxygenation and/or respiratory distress with non-invasive airway adjuncts:
 - a. Administer non-invasive positive pressure ventilation **[EMT-O; AEMT-R]** via continuous positive airway pressure (CPAP) or bi-level positive airway pressure (BiPAP) for severe respiratory distress.
 - b. Utilize bag-valve-mask ventilation in children with respiratory failure.
 9. Non-visualized airways **[EMR-O; EMT-R]** and intubation **[INT-O; PARA-R]**: Utilize *only* if bag-valve-mask ventilation fails. The airway should be managed in the least invasive way possible.

Patient Safety Considerations

1. Routine use of lights and sirens is not recommended during transport.
2. Giving positive pressure in the setting of bronchoconstriction, either via a supraglottic airway or intubation, increases the risk of air trapping which can lead to pneumothorax and cardiovascular collapse. These interventions should be reserved for situations of respiratory failure.

Notes and Educational Pearls

Key Considerations

- Inhaled magnesium sulfate should not be administered.
- Heliox should not be administered.
- COPD patients not in respiratory distress should be given oxygen to maintain adequate oxygen saturation above 90%.
- Nebulizer droplets can carry viral particles, so additional PPE should be considered, including placement of a surgical mask over the nebulizer to limit droplet spread.
- In the asthmatic patient, pharmacologic intervention should take priority over CPAP/BiPAP and be given in line with CPAP/BiPAP.

Pertinent Assessment Findings

In the setting of severe bronchoconstriction, wheezing might not be heard. Patients with known asthma who complain of chest pain or shortness of breath should be empirically treated, even if wheezing is absent.

Pulmonary Edema

Aliases

Congestive heart failure, respiratory distress, respiratory failure, acute respiratory distress syndrome, myocardial infarct, pulmonary embolism, COPD, asthma, anaphylaxis

Patient Care Goals

1. Decrease respiratory distress and work of breathing.
2. Maintain adequate oxygenation and perfusion.
3. Provide direct supportive efforts towards decreasing afterload and increasing preload.

Patient Presentation

Inclusion Criteria

1. Respiratory distress with presence of rales
2. Clinical impression consistent with congestive heart failure

Exclusion Criteria

1. Clinical impression consistent with infection (e.g. fever)
2. Clinical impression consistent with asthma or COPD

Patient Management

Assessment

1. History
 - a. Use of diuretics and compliance
 - b. Weight gain
 - c. Leg swelling
 - d. Orthopnea
2. Exam
 - a. Breath sounds—crackles or rales
 - b. Lower extremity edema
 - c. JVD
 - d. Cough and/or productive cough with pink and/or frothy sputum
 - e. Diaphoresis
 - f. Chest discomfort
 - g. Hypotension
 - h. Shock
 - i. Respiratory distress, assess:
 - i. Patient's ability to speak in full sentences
 - ii. Respiratory accessory muscle use

Treatment and Interventions

1. Manage airway as necessary.
2. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
3. Initiate monitoring and perform 12-lead ECG.
4. Establish IV access [*AEMT-R*].

5. Consider Nitroglycerin *[AEMT-R]*
6. Consider non-invasive positive pressure ventilation *[EMT-O; AEMT-R]*, non-visualized airway *[EMR-O; EMT-R]*, or endotracheal intubation *[INT-O; PARA-R]* for severe distress or if not improving with less invasive support.

Patient Safety Considerations

No recommendations

Notes and Educational Pearls

Key Considerations

1. Differential:
 - a. MI
 - b. CHF
 - c. Asthma
 - d. Anaphylaxis
 - e. Aspiration
 - f. COPD
 - g. Pleural effusion
 - h. Pneumonia
 - i. PE
 - j. Pericardial tamponade
 - k. Toxin exposure
2. Non-invasive positive pressure ventilation:
 - a. Contraindications:
 - i. Hypoventilation
 - ii. Altered level of consciousness
 - iii. Airway compromise
 - iv. Aspiration risk
 - v. Pneumothorax
 - vi. Facial trauma or burns
 - vii. Systolic BP less than 90 mmHg
 - viii. Recent oropharyngeal, tracheal, or bronchial surgery
 - b. Benefits:
 - i. Increased oxygenation and perfusion by reducing work of breathing
 - ii. Maintaining inflation of atelectatic alveoli
 - iii. Improving pulmonary compliance
 - iv. Decreases respiratory rate and the work of breathing, HR, and SBP
 - v. Improves delivery of bronchodilators
 - vi. Reduces preload and afterload, improving cardiac output
 - c. Complications:
 - i. Anxiety (most common)
 - ii. Theoretical risk of hypotension and pneumothorax as non-invasive positive pressure ventilation increases intrathoracic pressure which decreases venous return and cardiac output
 - iii. Sinusitis
 - iv. Skin abrasions
 - v. Conjunctivitis—minimized with proper size mask
 - vi. Potential for barotrauma—pneumothorax or pneumomediastinum (rare)

3. Positioning: Allow patient to remain in position of comfort—patients may decompensate if forced to lie down.
4. Common causes of pulmonary edema:
 - a. CHF (most common)
 - b. Medications
 - c. High altitude exposure
 - d. Kidney failure
 - e. Lung damage caused by gases or severe infection
 - f. Major injury
5. Nitrates: These medications provide both subjective and objective improvement, and might decrease intubation rates, incidence of MIs, and mortality. High-dose nitrates can reduce both preload and afterload and potentially increase cardiac output. Because many CHF patients present with very elevated arterial and venous pressure, frequent doses of nitrates may be required to control blood pressure and afterload. High dose nitrate therapy in patients in severe distress such as hypoxia, altered mentation, diaphoresis, or speaking in one-word sentences. An approach is to give two SL NTG (0.8 mg) for SBP greater than 160 mmHg or three SL NTG (1.2 mg) when SBP is greater than 200 mmHg every 5 minutes. A concern with high doses of nitrates is that some patients are very sensitive to even normal doses and may experience marked hypotension. It is therefore critical to monitor blood pressure during high-dose nitrate therapy.
6. Nitrates and phosphodiesterase inhibitors: The use of nitrates should be avoided in any patient who has used a phosphodiesterase inhibitor within the past 48 hours. Examples are: sildenafil (Viagra®, Revatio®), vardenafil (Levitra®, Staxyn®), tadalafil (Cialis®, Adcirca®) which are used for erectile dysfunction and pulmonary hypertension. Also avoid use in patients receiving intravenous epoprostenol (Flolan®) or treprostenil (Remodulin®) which is used for pulmonary hypertension. Administer nitrates with extreme caution, if at all, to patients with an inferior STEMI or suspected STEMI with right ventricular involvement because these patients require adequate RV preload.
7. Nitroglycerin: This drug reduces left ventricular filling pressure primarily via venous dilation. At higher doses the drug variably lowers systemic afterload and increases stroke volume and cardiac output. Although some have advocated early use of ACE inhibitors in patients with acute decompensated heart failure, we do not recommend this approach. There are limited data on the safety and efficacy of initiating new ACE inhibitors or angiotensin receptor blockers therapy in the early phase of therapy of acute decompensated heart failure (i.e. the first 12 to 24 hours).
8. Furosemide (Lasix®): Use is not recommended in the prehospital setting. Pulmonary edema is more commonly a problem of volume distribution than overload, so administration of furosemide provides no immediate benefit for most patients. Misdiagnosis of CHF and subsequent inducement of inappropriate diuresis can lead to increased morbidity and mortality in patients.

Trauma

General Trauma Management

Aliases

None noted

Patient Care Goals

1. Rapid assessment and management of life-threatening injuries
2. Safe movement of patient to prevent worsening injury severity
3. Rapid and safe transport to the appropriate level of trauma care

Patient Presentation

Inclusion Criteria

1. Patients of all ages who have sustained an injury as a result of mechanical trauma, including:
 - a. Blunt injury
 - b. Penetrating injury
 - c. Burns

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Primary survey
 - a. Hemorrhage control
 - i. Assess for and stop severe hemorrhage [see [Extremity Trauma/External Hemorrhage Management guideline](#)]
 - b. Airway
 - i. Assess airway patency by asking the patient to talk to assess stridor and ease of air movement.
 - ii. Look for injuries that may lead to airway obstruction including unstable facial fractures, expanding neck hematoma, blood or vomitus in the airway, facial burns/inhalation injury.
 - iii. Evaluate mental status for ability to protect airway (patients with a GCS less than or equal to 8 are likely to require airway protection).
 - c. Breathing
 - i. Assess respiratory rate and pattern.
 - ii. Assess symmetry of chest wall movement.
 - iii. Listen bilaterally on lateral chest wall for breath sounds.
 - d. Circulation
 - i. Assess blood pressure and heart rate.
 - ii. Look for signs of hemorrhagic shock (these include tachycardia, hypotension, pale, cool clammy skin, capillary refill ***greater than*** 2 seconds).

- e. Disability
 - i. Perform neurologic status assessment [see [Appendix VII](#)].
 - ii. Assess gross motor movement of extremities.
 - iii. Evaluate for clinical signs of traumatic brain injury with herniation including:
 - 1. Unequal pupils.
 - 2. Lateralizing motor signs.
 - 3. Posturing.
- f. Exposure
 - i. Perform rapid evaluation of entire body to identify sites of penetrating wounds or other blunt injuries. Be sure to roll patient and examine the back.
 - ii. Prevent hypothermia.

Treatment and Interventions

1. Hemorrhage control
 - a. Stop severe hemorrhage [see [Extremity Trauma/External Hemorrhage Management guideline](#)].
2. Airway
 - a. Establish patent airway with cervical spine precautions, per the [Airway Management](#) and [Spinal Care](#) guidelines.
 - b. If respiratory efforts are inadequate, assist with bag-mask ventilation and consider airway adjuncts. If patient is unable to maintain airway, consider oral airway (nasal airway should not be used with significant facial injury or possible basilar skull fracture).
 - c. If there is an impending airway obstruction or altered mental status resulting in inability to maintain airway patency, secure definitive airway.
3. Breathing
 - a. If absent or diminished breath sounds in a hypotensive patient, consider tension pneumothorax and perform needle decompression *[INT-R]*.
 - b. For open chest wound, place semi-occlusive dressing *[EMR-O; EMT-R]*.
 - j. Monitor oxygen saturation *[EMR-O; EMT-R]* and, if indicated, administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
4. Circulation
 - a. If pelvis is unstable and patient is hypotensive, place pelvic binder or sheet to stabilize pelvis *[EMR through INT-O; PARA-R]*.
 - b. Establish IV access *[AEMT-R]*.
 - c. Normal saline fluid resuscitation *[AEMT-R]*
 - i. Adults
 1. If SBP greater than 90 mmHg, no IV fluids required.
 2. If SBP less than 90 mmHg or HR greater than 120, consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*) and reassess.
 3. Penetrating trauma: target SBP 90mmHg (or palpable radial pulse).
 4. Head injury: target SBP 110-120 mmHg. Hypotension should be avoided to maintain cerebral perfusion.
 - ii. Pediatrics
 1. If child demonstrates tachycardia for age with signs of poor perfusion (low BP; greater than 2-second capillary refill; altered

- mental status; hypoxia; weak pulses; pallor; or mottled, cool skin), give 20ml/kg crystalloid bolus and reassess.
2. Target normal BP for age [see [Appendix VIII – Abnormal Vital Signs](#)].
 2. Disability
 - a. If clinical signs of traumatic brain injury, see [Head Injury](#) guideline.
 3. Exposure
 - a. Avoid hypothermia:
 - i. Remove wet clothing.
 - ii. Cover patient to prevent further heat loss.
 4. **NOTE:** Patients with major hemorrhage, hemodynamic instability, penetrating torso trauma, or signs of traumatic brain injury often require rapid surgical intervention. Minimize scene time (goal is under 10 minutes) and initiate rapid transport to the highest level of care within the trauma system.
 5. Decisions regarding transport destination should be based on the [Wisconsin Field TraumaTriage Guidelines](#).

Secondary Assessment, Treatment, and Interventions

1. Assessment
 - a. Obtain medical history from patient or family including:
 - i. Allergies
 - ii. Medications
 - iii. Past medical and surgical history
 - iv. Events leading up to the injury
 - b. Secondary survey: Head to toe physical exam
 - i. Head
 1. Palpate head and scalp and face and evaluate for soft tissue injury or bony crepitus.
 2. Assess pupils.
 - ii. Neck
 1. Check for:
 - a. Contusions.
 - b. Abrasions.
 - c. Hematomas.
 - d. JVD.
 - e. Tracheal deviation.
 2. Palpate for crepitus.
 3. Conduct a spinal assessment per the [Spinal Care guideline](#).
 - iii. Chest
 1. Palpate for instability or crepitus.
 2. Listen to breath sounds.
 3. Inspect for penetrating or soft tissue injuries.
 - iv. Abdomen
 1. Palpate for tenderness.
 2. Inspect for penetrating or soft tissue injuries.
 - v. Pelvis
 1. Inspect for penetrating or soft tissue injuries.
 2. Palpate once for instability by applying medial pressure on the iliac crests bilaterally.
 - vi. Back

1. Maintain spinal alignment. Refer to [Spinal Care guideline](#).
2. Inspect for penetrating or soft tissue injuries.
- vii. Neurologic status assessment [see [Appendix VII](#)]:
 1. Perform serial assessment of mental status.
 2. Perform gross exam of motor strength and sensation in all four extremities.
- viii. Extremities
 1. Assess for fracture or deformity.
 2. Assess peripheral pulses/capillary refill.
- c. Additional treatment considerations:
 - i. Maintain spine precautions per the [Spinal Care guideline](#).
 - ii. Splint obvious extremity fractures per the [Extremity Trauma/External Hemorrhage Management guideline](#).
 - iii. Provide pain medication per the [Pain Management guideline](#).

Patient Safety Considerations

1. Manage life-threatening injuries identified on primary survey immediately, with rapid transport to a trauma center. Perform secondary survey enroute.
2. Monitor patient for deterioration over time with serial vital signs and repeat neurologic status assessment [see [Appendix VII](#)].
 - a. Patients with compensated shock may not manifest hypotension until severe blood loss has occurred.
 - b. Patients with traumatic brain injury may deteriorate as intracranial swelling and hemorrhage increase.
3. Anticipate potential for progressive airway compromise in patients with trauma to head and neck.

Notes and Educational Pearls

Key Considerations

- Optimal trauma care requires a structured approach to the patient, emphasizing ABCDE (Airway, Breathing, Circulation, Disability, Exposure).
- Target scene time less than 10 minutes for unstable patients or those likely to need surgical intervention.
- Provider training should include the [State of Wisconsin Trauma Field Triage Guidelines](#).
- Frequent reassessment of the patient is important.
 - If patient develops difficulty with ventilation, reassess breath sounds for development of tension pneumothorax.
 - If extremity hemorrhage is controlled with pressure dressing or tourniquet, reassess for evidence of continued hemorrhage.
 - If mental status declines, reassess ABCs and repeat neurologic status assessment [see [Appendix VII](#)].

Traumatic Arrest: Withholding and Termination of Resuscitative Efforts

Resuscitative efforts should be withheld for trauma patients with the following:

1. Decapitation
2. Hemiorpectomy
3. Signs of rigor mortis or dependent lividity
4. Blunt trauma: apneic, pulseless, no organized cardiac activity on monitor

Note: Adult and Pediatric: Resuscitative efforts may be terminated in patients with traumatic arrest who have no return of spontaneous circulation after 15-30 minutes of resuscitative efforts, including airway management; evaluation and treatment for possible tension pneumothorax; fluid bolus; and minimally interrupted CPR.

Blast Injuries

Aliases

None noted

Patient Care Goals

1. Maintain patient and provider safety by identifying ongoing threats at the scene of an explosion.
2. Identify multi-system injuries which may result from a blast, including possible toxic contamination.
3. Prioritize treatment of multi-system injuries to minimize patient morbidity.

Patient Presentation

Inclusion Criteria

1. Patients exposed to explosive force. Injuries may include any or all of the following:
 - a. Blunt trauma
 - b. Penetrating trauma
 - c. Burns
 - d. Pressure-related injuries (barotrauma)
 - e. Toxic chemical contamination

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Hemorrhage Control
 - a. Assess for and stop severe hemorrhage [see [Extremity Trauma/External Hemorrhage Management guideline](#)].
2. Airway
 - a. Assess airway patency.
 - b. Consider possible thermal or chemical burns to airway.
3. Breathing
 - a. Evaluate adequacy of respiratory effort, oxygenation, quality of lung sounds, and chest wall integrity.
 - b. Consider possible pneumothorax or tension pneumothorax (as a result of penetrating, blunt trauma or barotrauma).
4. Circulation
 - a. Look for evidence of external hemorrhage.
 - b. Assess BP, pulse, skin color/character, and distal capillary refill for signs of shock.
5. Disability
 - a. Assess patient responsiveness (AVPU) and level of consciousness (GCS) [see [Appendix VII](#)].
 - b. Assess pupils.
 - c. Assess gross motor movement and sensation of extremities.
6. Exposure
 - a. Perform rapid evaluation of entire skin surface, including back (log roll), to

identify blunt or penetrating injuries.

Treatment and Interventions

1. Hemorrhage control:
 - a. Control any severe external hemorrhage [see [Extremity Trauma/External Hemorrhage Management guideline](#)].
2. Airway:
 - a. Secure airway, utilizing airway maneuvers, airway adjuncts, non-visualized airways [*EMR-O; EMT-R*], or endotracheal tube [*INT-O; PARA-R*] [see [Airway Management guideline](#)].
 - b. If thermal or chemical burn to airway is suspected, early airway control is vital.
3. Breathing:
 - a. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients. Assist respirations as needed.
 - b. Cover any open chest wounds with semi-occlusive dressing [*EMR-O; EMT-R*].
 - c. If patient has evidence of tension pneumothorax, perform needle decompression [*INT-R*].
4. Circulation:
 - a. Establish IV access with two large bore IVs or IOs.
 - i. Administer NS or LR, per the [General Trauma Management guideline](#).
 - ii. If patient is burned, administer NS or LR per the [Burns guideline](#).
5. Disability:
 - a. If evidence of head injury, treat per the [Head Injury guideline](#).
 - b. Apply spinal precautions per the [Spinal Care guideline](#).
 - c. Monitor GCS during transport to assess for changes.
6. Exposure:
 - a. Keep patient warm to prevent hypothermia.

Patient Safety Considerations

1. Ensuring scene safety is especially important at the scene of an explosion.
 - a. Consider possibility of subsequent explosions, structural safety, possible toxic chemical contamination, the presence of noxious gasses, and other hazards.
 - b. In a possible terrorist event, consider the possibility of secondary explosive devices.
2. Remove patient from the scene as soon as is practical and safe.
3. If the patient has sustained burns (thermal, chemical, or airway), consider transport to specialized burn center.

Notes and Educational Pearls

Key Considerations

- Scene safety is of paramount importance when responding to an explosion or blast injury.
- 1. Patients sustaining blast injury: Be prepared to address complex, multi-system injuries including: blunt and penetrating trauma, shrapnel, barotrauma, burns, and toxic chemical exposure.
- Patients sustaining airway injury, particularly airway burns: Employ early and aggressive airway management.
- IV fluid resuscitation: Minimize in patients without signs of shock.

- Injuries due to barotrauma
 - Tension pneumothorax
 - Hypotension or other signs of shock associated with decreased or absent breath sounds, jugular venous distension, and/or tracheal deviation
 - Tympanic membrane perforation resulting in deafness which may complicate the evaluation of their mental status and their ability to follow commands
- Transport: Primary transport to a trauma or burn center is preferable, whenever possible.

Pertinent Assessment Findings

1. Evidence of multi-system trauma, especially:
 - a. Airway injury or burn
 - b. Barotrauma to lungs
 - c. Toxic chemical contamination

Burns

Aliases

None noted

Patient Care Goals

Minimize tissue damage and patient morbidity from burns.

Patient Presentation

1. Patient may present with:
 - a. Airway—stridor, hoarse voice
 - b. Mouth and nares—redness, blisters, soot, singed hairs
 - c. Breathing—rapid, shallow, wheezes, rales
 - d. Skin—estimate Total Burn Surface Area (TBSA) and depth (partial vs. full thickness)
 - e. Associated trauma—blast, fall, assault

Inclusion Criteria

Patients sustaining thermal burns

Exclusion Criteria

Electrical, chemical, and radiation burns [see [Toxins and Environmental](#) section]

Special Transport Considerations

1. Transport to most appropriate trauma center when there is airway or respiratory involvement, or when significant trauma or blast injury is suspected.
2. Consider air ambulance transportation for long transport times or airway management needs beyond the scope of the responding ground medic.
3. Consider transport directly to burn center if partial or full thickness burns (TBSA) greater than 10%, involvement of hands, feet, genitalia, face, and/or circumferential burns.

Scene Management

1. Assure crew safety:
 - a. Power off
 - b. Electrical lines secure
 - c. Gas off
 - d. No secondary devices
 - e. Hazmat determinations made
 - f. Proper protective attire including breathing apparatus may be required

Patient Management

Assessment

1. Circumstances of event—consider:
 - a. Related trauma in addition to the burns.
 - b. Inhalation exposures such as CO and cyanide (CN).

- c. Pediatric or elder abuse.
2. Follow ABCs of resuscitation per the [General Trauma Management guideline](#).
3. If evidence of possible airway burn, consider aggressive airway management.
4. Consider spinal precautions for those that qualify per the [Spinal Care guideline](#).
5. Estimate TBSA burned and depth of burn.
 - a. Use "Rule of 9's" [see burn related tables in [Appendix VI](#)].

Note: First- degree burns (skin erythema only) are not included in TBSA calculations.
6. Document pain scale.

Treatments and interventions

1. Stop the burning:
 - a. Remove wet clothing (if not stuck to the patient).
 - b. Remove jewelry.
 - c. Leave blisters intact.
2. Minimize burn wound contamination:
 - a. Cover burns with dry dressing or clean sheet.
 - b. Do not apply gels or ointments.
3. Monitor SPO₂ [*EMR-O; EMT-R*], ETCO₂ and ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*]; consider SPCO monitoring, if available.
4. Administer high-flow supplemental oxygen [*EMR-O; EMT-R*] for all burn patients rescued from an enclosed space.
5. Establish IV access [*AEMT-R*], avoid placement through burned skin.
6. Evaluate distal circulation in circumferentially burned extremities.
7. Consider early management of pain and nausea or vomiting.
8. Initiate fluid resuscitation. Use lactated Ringer's [*AEMT-O*] or normal saline [*AEMT-R*]
 - a. If patient in shock:
 - i. Consider other cause, such as trauma or cyanide toxicity.
 - ii. Administer IV fluid [*AEMT-R*] per the [Shock guideline](#).
9. Prevent systemic heat loss and keep the patient warm.

Special Treatment Considerations

1. If blast mechanism, treat per the [Blast Injury guideline](#).
2. Recognize that airway burns can rapidly lead to upper airway obstruction and respiratory failure.
3. Consider the potential for cyanide poisoning in a patient with depressed GCS, respiratory difficulty, and cardiovascular collapse in the setting of an enclosed-space fire. Give the antidote (hydroxocobalamin), if available, in this circumstance.
4. Recognize that carbon monoxide toxicity, particularly in enclosed-space fires, may affect the accuracy of pulse oximetry [see [Carbon Monoxide/Smoke Inhalation guideline](#)].
5. For specific chemical exposures (cyanide, hydrofluoric acid, other acids and alkali) [see [Topical Chemical Burn guideline](#)].
6. Consider decontamination and notification of receiving facility of potentially contaminated patient (e.g. methamphetamine (meth) lab incident).

Notes and Educational Pearls

- Onset of stridor and change in voice are sentinel signs of potentially significant airway

burns, which may rapidly lead to airway obstruction or respiratory failure.

- If the patient is in shock within one hour of burn, it is not from the burn. Evaluate the patient carefully for associated trauma or cyanide toxicity.
- If the patient is not in shock, the fluid rates recommended above will adequately maintain patient's fluid volume.
- Pain management is critical in acute burns.
- ETCO₂ monitoring may be particularly useful to monitor respiratory status in patients receiving significant doses of narcotic pain medication.
- ECG cardiac monitor is important in electrical burns and chemical inhalations.
- TBSA is calculated only based on percent of second and third degree burns. First degree burns are not included in this calculation.

Crush Injury

Aliases

Crush, compartment syndrome

Patient Care Goals

1. Recognizing traumatic crush injury mechanism
2. Minimizing systemic effects of the crush syndrome

Patient Presentation

Inclusion criteria

Traumatic crush mechanism of injury

Exclusion criteria

Non-crush injuries

Patient Management

Assessment

1. Identify any severe hemorrhage.
2. Assess airway, breathing, and circulation.
3. Evaluate for possible concomitant injury (e.g. fractures, solid organ damage, or spinal injury.)
4. Monitor for development of compartment syndrome.

Treatment and Interventions

1. Treat crushed casualties as soon as they are discovered.
2. If severe hemorrhage is present, see [Extremity Trauma/External Hemorrhage Management guideline](#).
3. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
4. Establish intravenous access *[AEMT-R]* with normal saline initial bolus of 10–15 ml/kg (prior to extrication if possible).
5. Consider sodium bicarbonate for significant crush injuries or prolonged entrapment of an extremity *[PARA-R]*.
6. Attach ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]*. Obtain/interpret 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*, if available. Carefully monitor for dysrhythmias or signs of hypokalemia before and immediately after release of pressure and during transport (e.g. peaked T waves, wide QRS, lengthening QT interval, loss of P wave).
7. Consider analgesics for pain control [see [Pain Management guideline](#)].
8. Consider the following post-extrication:
 - a. Continued resuscitation with normal saline (500-1000 ml/hr for adults, 10 cc/kg/hr for children)
 - b. If ECG *[Acquisition EMT-O; Interpretation INT-R]* suggestive of hyperkalemia, if findings of hyperkalemia, administer normal saline IV fluids and consider administration of calcium gluconate (preferred) *[PARA-R]*.
 - c. If not already administered, for significant crush injuries with ECG *[Acquisition*

EMT-O; Interpretation INT-R] suggestive of hyperkalemia, administer sodium.

- d. If ECG [*Acquisition EMT-O; Interpretation INT-R*] suggestive of hyperkalemia, consider albuterol 5 mg via small volume nebulizer.

Patient Safety Considerations

Scene safety for both rescuers and patients is of paramount importance.

Notes and Educational Pearls

1. Causes of mortality in untreated crush syndrome:
 - a. Immediate
 - i. Severe head injury
 - ii. Traumatic asphyxia
 - iii. Torso injury with damage to intrathoracic or intra-abdominal organs
 - b. Early
 - i. Hyperkalemia (potassium is released from injured muscle cells)
 - ii. Hypovolemia, shock
 - c. Late
 - i. Renal failure (from release of toxins from injured muscle cells)
 - ii. Coagulopathy and hemorrhage
 - iii. Sepsis

Key Considerations

- Perform rapid extrication and evacuation to a definitive care facility (trauma center preferred).
- Maintain a high index of suspicion for any patient with a compressive mechanism of injury, as a patient with a crush injury may initially present with very few signs and symptoms.
- Recognize that a fatal medical complication of crush syndrome is hyperkalemia. Suspect hyperkalemia if T- waves become peaked, QRS becomes prolonged (greater than 0.12 seconds), P wave is absent, or QTc is prolonged.
- Avoid lactated Ringer's solution as it contains potassium.
- Continue fluid resuscitation through extrication and transfer to hospital.

Pertinent Assessment Findings

- Evaluation of mental status, GCS
- Evaluation for fractures and potential compartment syndrome development (neurovascular status of injured extremity)
- Examination of spine
- Evidence of additional trauma, potentially masked by with other painful injuries

Extremity Trauma and External Hemorrhage Management

Aliases

None noted

Patient Care Goals

1. Minimize blood loss from extremity hemorrhage.
2. Avoid hemorrhagic shock as a result of extremity hemorrhage.
3. Minimize pain and further injury as a result of potential fractures or dislocations.

Patient Presentation

Inclusion Criteria

1. Traumatic extremity hemorrhage (external hemorrhage)
2. Potential extremity fractures or dislocations

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Evaluate for obvious deformity, shortening, rotation, or instability
2. Assess neurologic status of extremity
 - a. Sensation to light touch
 - b. Distal movement of extremity
3. Assess vascular status of extremity
 - a. Pallor
 - b. Pulse
 - c. Capillary refill
 - d. Degree of bleeding/blood loss with assessment of the color of the blood (venous or arterial) and whether it is pulsatile or not

Treatments and Interventions (also, see protocol diagram below)

1. Manage bleeding
 - a. Apply direct pressure to bleeding site followed by pressure dressing.
 - b. If direct pressure and/or pressure dressing is ineffective or impractical:
 - i. If the bleeding site is amenable to tourniquet placement, apply tourniquet to extremity:
 1. Tourniquet should be placed 2–3 inches proximal to wound, not over a joint, and tightened until bleeding stops and distal pulse is eliminated.
 2. If bleeding continues, place a second tourniquet proximal to the first.
 3. For thigh wounds, consider placement of two tourniquets, side-by-side, and tighten sequentially to eliminate distal pulse.
 - ii. If the bleeding site is not amenable to tourniquet placement (i.e. junctional injury), pack wound tightly with a hemostatic gauze **[ALL**

EMS PRACTICE LEVELS-O and apply direct pressure.

- c. Groin or axillary injury
 - i. Apply direct pressure to wound.
 - ii. If still bleeding, pack wound tightly with hemostatic gauze ***[ALL EMS PRACTICE LEVELS--O]*** and apply direct pressure.
 - iii. Consider using a junctional hemostatic device if available.
2. Manage pain [see [Pain Management guideline](#)]
 - a. Pain management should be strongly considered for patients with suspected fractures.
 - b. If tourniquet placed, an alert patient will likely require pain medication to manage tourniquet pain.
3. Stabilize suspected fractures or dislocations
 - a. Strongly consider pain management before attempting to move a suspected fracture.
 - b. If distal vascular function is compromised, gently attempt to restore normal anatomic position.
 - c. Use splints as appropriate to limit movement of suspected fracture.
 - d. Elevate extremity fractures above heart level whenever possible to limit swelling.
 - e. Apply ice/cool packs to limit swelling in suspected fractures or soft tissue injury. Do not apply ice directly to skin.
 - f. Reassess distal neurovascular status after any manipulation or splinting of fractures or dislocations.

Patient Safety Considerations

1. If tourniquet use:
 - a. Ensure that it is sufficiently tight to occlude the distal pulse, in order to avoid compartment syndrome.
 - b. Ensure that it is well marked and visible and that all subsequent providers are aware of the presence of the tourniquet.
 - c. Do not cover with clothing or dressings.
2. Mark time of tourniquet placement prominently on the patient.
3. If pressure dressing or tourniquet used, frequently re-check to determine if bleeding has restarted. Check for blood soaking through the dressing or continued bleeding distal to the tourniquet. Do not remove tourniquet or dressing in order to assess bleeding.

Notes and Educational Pearls


Key Considerations

- Tourniquet may be placed initially to stop obvious severe hemorrhage, then replaced later with pressure dressing after stabilization of ABCs and packaging of patient. Tourniquet should not be removed if:
 - Transport time short (less than 30 minutes)
 - Amputation or near-amputation
 - Unstable or complex multiple-trauma patient
 - Unstable clinical or tactical situation
- If tourniquet is replaced with pressure dressing, leave loose tourniquet in place so it may be retightened if bleeding resumes.
- Survival is markedly improved when a tourniquet is placed *before* shock ensues.
- Commercial, properly tested tourniquets are preferred over improvised tourniquets.
- If hemostatic gauze is not available, plain gauze tightly packed into a wound has

been shown to be effective.

- Arterial pressure points are not effective in controlling hemorrhage.
- Amputated body parts should be transported with patient for possible re-implantation.
 - Keep the amputated body part cool but dry.
 - Place the amputated part in a plastic bag.
 - Place the bag with the amputated part on ice in a second bag.
 - Do not let the amputated part come into direct contact with the ice.

Prehospital External Hemorrhage Control Protocol



* Use of tourniquet for extremity hemorrhage is strongly recommended if sustained direct pressure is ineffective or impractical; use a commercially-produced, windlass, pneumatic, or ratcheting device, which has been demonstrated to occlude arterial flow, and avoid narrow, elastic, or bungee-type devices; utilize improvised tourniquets only if no commercial device is available; do not release a properly-applied tourniquet until the patient reaches definitive care.

Apply a topical hemostatic agent, in combination with direct pressure, for wounds in anatomic areas where tourniquets cannot be applied and sustained direct pressure alone is ineffective or impractical; only apply topical hemostatic agents in a gauze format that support wound packing; only utilize topical hemostatic agents which have been determined to be effective and safe in a standardized laboratory injury model.

Source: Bulger et al. 2014

Facial and/or Dental Trauma

Aliases

None noted

Patient Care Goals

1. Preservation of a patent airway
2. Preservation of vision
3. Preservation of dentition

Patient Presentation

Inclusion Criteria

Isolated facial injury, including trauma to the eyes, nose, ears, midface, mandible, dentition

Exclusion Criteria

1. General Trauma [see [General Trauma Management guideline](#)]
2. Burn trauma [see [Burns guideline](#)]

Patient Management

Assessment

1. Consider patient medications with focus on blood thinners or anti-platelet agents.
2. Apply ABCs with particular focus on ability to keep airway patent:
 - a. Stable midface
 - b. Stable mandible
 - c. Stable dentition (poorly anchored teeth require vigilance for possible aspiration)
3. Assess bleeding (which may be severe—epistaxis, oral trauma, and facial lacerations).
4. Assess cervical spine pain or tenderness [see [Spinal Care guideline](#)].
5. Assess mental status for possible traumatic brain injury [see [Head Injury guideline](#)].
6. Perform gross vision assessment.
7. Watch for dental avulsions.
8. Collect any avulsed tissue or teeth.
9. Check airway for obstructions: Lost teeth not recovered on scene may be in the airway.
10. Assess for overall trauma.
11. Perform specific re-examination geared toward airway and ability to ventilate adequately.

Treatment and Interventions

1. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
2. Use ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] to help monitor for hypoventilation and apnea.
3. Establish IV access, as needed, for fluid or medication administration [*AEMT-R*].
4. Administer pain medication per the [Pain Management guideline](#).
5. Perform the following for an avulsed tooth:
 - a. Avoid touching the root of the avulsed tooth. Do not wipe off tooth.
 - b. Pick up at crown end. If dirty, rinse off under cold water for 10 seconds.
 - c. Place in milk or saline as the storage medium. Alternatively, an alert and cooperative patient can hold tooth in mouth using own saliva as storage medium.

6. Eye trauma:
 - a. Place eye shield for any significant eye trauma.
 - b. If globe is avulsed, do not put back into socket. Cover with moist saline dressings and then place cup over it.
7. Mandible unstable :
 - a. Expect patient cannot spit or swallow effectively and have suction readily available.
 - b. Transport sitting up (preferred) with emesis basin or suction available. (In the absence of a suspected spinal injury, see [Spinal Care guideline](#).)
8. Epistaxis: squeeze nose (or have patient do so) for 10–15 minutes continuously.
9. Nose or ear avulsion:
 - a. Recover tissue if it does not waste scene time.
 - b. Transport with tissue wrapped in dry sterile gauze in a plastic bag placed on ice.
 - c. Address severe ear and nose lacerations with a protective moist sterile dressing.

Patient Safety Considerations

1. Conduct frequent reassessment of airway.
2. Maintain patency of airway; this is the highest priority: Conduct cervical spine assessment for field clearance [*EMR-O; EMT-R*] (per [Spinal Care guideline](#)) to enable transport sitting up for difficulty with bleeding, swallowing, or handling secretions.

Notes and Educational Pearls

Key Considerations

- Airway may be compromised because of fractures or bleeding
- After nasal fractures, epistaxis may be posterior and may not respond to direct pressure over the nares with bleeding running down posterior pharynx, potentially compromising airway
- Protect avulsed tissue and teeth
 - Avulsed teeth may be successfully re-implanted if done so in a very short period after injury
 - Use sterile dressing for ear and nose cartilage

Pertinent Assessment Findings

1. Unstable facial fractures that can abruptly compromise airway
2. Loose teeth and retro-pharynx bleeding

Head Injury

Aliases

None noted

Patient Care Goals

1. Limit disability and mortality from head injury by:
 - a. Promoting adequate oxygenation (avoid hypoxia and hyperventilation).
 - b. Promoting adequate cerebral perfusion (avoid hypotension).
 - c. Limiting development of increased intracranial pressure.
 - d. Limiting secondary brain injury.

Patient Presentation

Inclusion Criteria

Adult or pediatric patient with blunt or penetrating head injury—LOC or amnesia not required

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Spine stabilization: Maintain cervical stabilization [see [Spinal Care guideline](#)].
2. Assessment: Conduct primary survey per the [General Trauma Management guideline](#).
3. Monitoring:
 - a. Conduct continuous pulse oximetry [*EMR-O; EMT-R*].
 - b. Conduct frequent systolic and diastolic blood pressure measurement.
 - c. Conduct initial neurologic status assessment [see [Appendix VII – Neurologic Status Assessment](#)] and reassessment with any change in mentation.
 - d. Apply continuous waveform ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*], if available, in moderate to severe head injury.
4. Secondary survey pertinent to isolated head injury:
 - a. Head: Gently palpate skull to evaluate for depressed or open skull fracture.
 - b. Eyes:
 - i. Evaluate pupil size and reaction to light to establish baseline.
 - ii. Reassess pupils if decrease in mentation.
 - c. Nose, mouth, and ears: Evaluate for blood or other fluid drainage.
 - d. Face: Evaluate for bony stability.
 - e. Neck: Palpate for cervical spine tenderness or deformity.
 - f. Neurologic:
 - i. Perform neurologic status assessment (GCS or AVPU).
 - ii. Evaluate for focal neurologic deficit: motor and sensory.

Treatment and Interventions

Note: These are not necessarily listed in the order they are to be performed, but are grouped by conceptual areas.

1. Airway:

- a. Administer oxygen **[EMR-O; EMT-R]** as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
- b. If patient unable to maintain airway, consider oral airway (nasal airway should not be used with significant facial injury or possible basilar skull fracture).
- c. Initiate oral endotracheal intubation **[INT-O; PARA-R]**. Non-visualized airway **[EMR-O; EMT-R]** insertion can be used if BVM ventilation is ineffective in maintaining oxygenation, or if airway is continually compromised.
- d. Do not use nasal intubation in patients with head injury.
2. Breathing:
 - a. For patients with a moderate or severe head injury who are unable to maintain their airway: Use continuous waveform capnography, and EtCO₂ measurement **[Interpretation INT-R]** if available, with a target EtCO₂ of 35-40 mmHg.
 - b. Non-visualized airway placement **[EMR-O; EMT-R]** or endotracheal intubation **[INR-O; PARA-R]** should only be performed if BVM ventilation is inadequate to maintain adequate oxygenation with a target EtCO₂ of 35-40 mmHg.
 - c. For patients with a severe head injury with signs of herniation: hyperventilate to a target EtCO₂ of 30–35 mmHg **[Acquisition EMT-O; Interpretation INT-O/PARA-R]** as a short-term option, and only for severe head injury with signs of herniation.
3. Circulation:
 - a. Wound care
 - i. Control bleeding with direct pressure if no suspected open skull injury.
 - ii. Apply moist sterile dressing to any potential open skull wound.
 - iii. Cover an injured eye with moist saline dressing and place cup over it.
 - b. Moderate or severe closed head injury
 - i. Blood pressure: Avoid hypotension and low cerebral perfusion by administering normal saline bolus 20 mg/kg **[AEMT-R]** to achieve the following goals:
 1. Adult (age greater than 10 yo): Maintain SBP greater than or equal to 110 mmHg.
 2. Pediatric: Maintain SBP:
 - a. Less than 1 month: greater than 60 mmHg
 - b. 1–12 months: greater than 70 mmHg
 - c. 1–10 yo: greater than 70 + 2x age in years
 - c. Do not delay transport to initiate IV access.
4. Disability :
 - a. Evaluate for other causes of altered mental status; check blood glucose.
 - b. Conduct spinal assessment and management, per [Spinal Care guideline](#).
 - c. Perform and trend neurologic status assessment (moderate/severe: GCS 3-13, P {pain} or U {unresponsive} on AVPU scale).
 - i. Early signs of deterioration:
 1. Confusion
 2. Agitation
 3. Drowsiness
 4. Vomiting
 5. Severe headache
 - ii. Monitor for signs of herniation.
 - d. Severe head injury: Elevate head of bed 30 degrees.

5. Use transport destination specific to head trauma.
 - a. Preferential transport to highest level of care within regional trauma system:
 - i. GCS 3–13, P (pain) or U (unresponsive) on AVPU scale
 - ii. Penetrating head trauma
 - iii. Open or depressed skull fracture

Patient Safety Considerations

1. Do not hyperventilate patient unless signs of herniation.
2. Assume concomitant cervical spine injury in patients with moderate or severe head injury.
3. **Geriatric Consideration:** Elderly patients with ankylosing spondylitis or severe kyphosis should be padded and immobilized in a position of comfort and may not tolerate a cervical collar.

Notes and Educational Pearls

Key Considerations

- Head injury severity guideline:
 - Mild: GCS 13–15 / AVPU = (A)
 - Moderate: GCS 9–12 / AVPU = (V)
 - Severe: GCS 3–8 / AVPU = (P) or (U)
- Providers should be specifically trained in accurate neurologic status assessment. [See [Appendix VII – Neurologic Status Assessment](#).]
- If endotracheal intubation or invasive airways are used, continuous waveform capnography is required to document proper tube placement and assure proper ventilation rate.
- Signs of herniation:
 - Decreasing mental status
 - Abnormal respiratory pattern
 - Asymmetric or unreactive pupils
 - Decorticate posturing
 - Cushing's response (bradycardia and hypertension)
 - Decerebrate posturing

Pertinent Assessment Findings

1. Neurologic status assessment findings
2. Pupils
3. Trauma findings on physical exam

High Threat Considerations, Active Shooter Scenario

Aliases

None noted

Definitions

- Hot Zone or Direct Threat Zone: an area within the inner perimeter where active threat and active hazards exist
- Warm Zone or Indirect Threat Zone: an area within the inner perimeter where security and safety measures are in place; this zone may have potential hazards, but no active danger exists

Patient Care Goals

1. Assess scene.
2. Mitigate further harm.
3. Accomplish goal with minimal additional injuries.

Patient Presentation

Inclusion Criteria

High threat environment: when greater than normal conditions exist that are likely to cause damage or danger to provider or patient

Exclusion Criteria

No significant threat exists to provider and patient, allowing for the performance of routine care.

Patient Management

Assessment, Treatment, and Interventions

1. Hot Zone or Direct Threat care considerations:
 - a. Defer in depth medical interventions if engaged in ongoing direct threat (e.g. active shooter, unstable building collapse, improvised explosive device, hazardous material threat).
 - b. Employ threat mitigation techniques; they will minimize risk to patients and providers.
 - c. Defer triage to a later phase of care.
 - d. Prioritize extraction based on resources available and the situation at hand.
 - e. Employ minimal interventions; these circumstances warrant it.
 - f. Encourage patients to provide self-first aid or instruct aid from uninjured bystander
 - g. Consider hemorrhage control:
 - i. Tourniquet application is the primary "medical" intervention to be considered in Hot Zone or Direct Threat circumstance.
 - ii. Consider instructing patient to apply direct pressure to the wound if no tourniquet available (or application is not feasible).
 - iii. Consider quickly placing or directing patient to be placed in position to protect airway, if not immediately moving patient.
2. Warm Zone or Indirect Threat care considerations:
 - a. Maintain situational awareness.

- b. Ensure safety of both responders and patients by rendering equipment and environment safe (firearms, vehicle ignition).
- c. Conduct primary survey, per the [General Trauma Management guideline](#), and initiate appropriate life-saving interventions:
 - i. Maintain hemorrhage control:
 - 1. Apply a tourniquet.
 - 2. Employ wound packing if feasible *[EMR-O; EMT-R]*.
 - ii. Maintain airway and support ventilation [see [Airway Management guideline](#)].
- d. Do not delay patient extraction and evacuation for non-life-saving interventions.
- e. Consider establishing a casualty collection point if multiple patients are encountered.
- f. Limit triage in this phase of care to the following categories (unless in a fixed casualty collection point):
 - i. Uninjured and/or capable of self-extraction
 - ii. Deceased/expectant
 - iii. All others

Patient Safety Considerations

- 1. Anticipate unique threats based on situation.
- 2. Consider provider safety in balancing the risks and benefits of patient treatment.

Notes and Educational Pearls

Key Considerations

- Novel risk assessment should be considered. Provider and patient safety will need to be simultaneously considered.
- Integrated response in partnership with other public safety entities may be warranted.
- A little risk may reap significant benefits to patient safety and outcome.
- Maintaining communications and incident management concepts may be crucial to maximizing efficiency and mitigating dangers.

Spinal Care

(Adapted from an evidence-based guideline created using the National Prehospital Evidence-Based Guideline Model Process)

Aliases

None noted

Patient Care Goals

1. Identify patients for whom spinal motion restriction (SMR) *[EMR-O; EMT-R]* is indicated.
2. Minimize secondary injury to spine in patients who have, or may have, an unstable spinal injury.
3. Minimize patient morbidity from the use of immobilization devices.

Patient Presentation

Inclusion criteria

Traumatic mechanism of injury

Exclusion criteria

No recommendations

Patient Management

Assessment

1. Assess the scene to determine the mechanism of injury.
 - a. Mechanism alone should not determine if a patient requires spinal motion restriction; however, mechanisms that have been associated with a higher risk of injury are:
 - i. Motor vehicle crashes (including automobiles, all-terrain vehicles, and snowmobiles).
 - ii. Axial loading injuries to the spine.
 - iii. Falls greater than 10 feet.
2. Assess the patient in the position found for findings associated with spine injury:
 - a. Mental status
 - b. Neurologic deficits
 - c. Spinal pain or tenderness
 - d. Any evidence of intoxication
 - e. Other severe injuries, particularly associated torso injuries

Treatment and Interventions

1. Place patient in cervical collar *[EMR-O; EMT-R]* if there are any of the following:
 - a. Patient complains of midline neck or spine pain
 - b. Any midline neck or spinal tenderness with palpation
 - c. Any abnormal mental status (including extreme agitation)
 - d. Focal or neurologic deficit
 - e. Any evidence of alcohol or drug intoxication
 - f. Another severe or painful distracting injury is present
 - g. Torticollis (in children)
 - h. A communication barrier that prevents accurate assessment

- Note:** If none of the above apply, patient may be managed without a cervical collar.
2. Do not place patients with penetrating injury to the neck in a cervical collar, and do not employ other spinal precautions regardless of whether or not they are exhibiting neurologic symptoms. Doing so can lead to delayed identification of injury or airway compromise, and has been associated with increased mortality.
 3. Employ the following if extrication is required:
 - a. From a vehicle: After placing a cervical collar, if indicated, children in a booster seat and adults should be allowed to self-extricate. For infants and toddlers already strapped in a car seat with a built-in harness, extricate the child while strapped in his or her car seat.
 - b. From other situations: A padded long board may be used for extrication, using the lift and slide (rather than a logroll) technique.
 4. Employ the following if patient is wearing a helmet:
 - a. Football helmet: Remove the face mask followed by manual removal (rather than using automated devices) of the helmet while keeping the neck manually immobilized. Occipital and shoulder padding should be applied, as needed, with the patient in a supine position, in order to maintain neutral cervical spine positioning.
 - b. Other helmet types: Evidence is lacking to provide guidance about other types of helmet removal.
 5. Do not transport patients on rigid long boards, unless the clinical situation warrants long board use. An example of this may be facilitation of immobilization of multiple extremity injuries or an unstable patient where removal of a board will delay transport and/or other treatment priorities. In these situations, long boards should ideally be padded or have a vacuum mattress applied to minimize secondary injury to the patient.
 6. Transport patients to the nearest appropriate facility in accordance with the State of Wisconsin Field Trauma Triage Guidelines.
 7. Immobilize patients with severe kyphosis or ankylosing spondylitis in a position of comfort using towel rolls or sand bags. These patients may not tolerate a cervical collar.

Patient Safety Considerations

- Immobilized patients with nausea or vomiting, or with facial or oral bleeding can potentially suffer airway compromise or aspiration.
- Excessively tight immobilization straps can limit chest excursion and cause hypoventilation.
- Prolonged immobilization on spine board can lead to ischemic pressure injuries to skin.
- Prolonged immobilization on spine board can be very uncomfortable for patient.
- Children are abdominal breathers, so immobilization straps should go across chest and pelvis and not across the abdomen, when possible.
- Children have disproportionately larger heads. When securing pediatric patients to a spine board, the board should have a recess for the head, or the body should be elevated approximately 1–2 cm to accommodate the larger head size and avoid neck flexion when immobilized.
- In an uncooperative patient, avoid interventions that may promote increased spinal movement.
- The preferred position for all patients with spine management is flat and supine. There are three circumstances under which raising the head of the bed to 30

degrees should be considered:

- Respiratory distress
- Suspected severe head trauma
- Promotion of patient compliance

Notes and Educational Pearls

Key Considerations

- Evidence is lacking to support or refute the use of manual stabilization prior to spinal assessment in the setting of a possible traumatic injury, when the patient is alert with spontaneous head and neck movement.
Providers should not manually stabilize these alert and spontaneously moving patients, since patients with pain will self-limit movement, and forcing immobilization in this scenario may unnecessarily increase discomfort and anxiety.
- Certain populations with musculoskeletal instability may be predisposed to cervical spine injury. However, evidence does not support or refute that these patients should be treated differently than those who do not have these conditions. These patients should be treated according to the [Spinal Care guideline](#) like other patients without these conditions.
- Age alone should not be a factor in decision-making for prehospital spine care, yet the patient's ability to reliably be assessed at the extremes of age should be considered. Communication barriers with infants and toddlers or elderly patients with dementia may prevent the provider from accurately assessing the patient.
- Spinal precautions should be considered a treatment or preventive therapy.
- Patients who are likely to benefit from immobilization should undergo this treatment.
- Patients who are not likely to benefit from immobilization, who have a low likelihood of spinal injury, should not be immobilized.
- Ambulatory patients may be safely immobilized on gurney with cervical collar and straps and will not generally require a spine board.
- Reserve long spine board use for the movement of patients whose injuries limit ambulation and who meet criteria for the use of spinal precautions. Remove from the long board as soon as is practical.

Pertinent Assessment Findings

- Mental status
- Normal neurologic examination
- Evidence of intoxication
- Evidence of multiple trauma with other severe injuries

Toxins and Environmental

Poisoning or Overdose Universal Care

Aliases

Toxin, overdose, poison, exposure

Patient Care Goals

1. Remove patient from hazardous material environment and decontaminate to remove continued sources of absorption, ingestion, inhalation, or injection (if properly trained and equipped).
2. Identify intoxicating agent by toxidrome or appropriate environmental testing.
3. Assess risk for organ impairments (heart, brain, kidney).
4. Identify antidote or mitigating agent.
5. Treat signs and symptoms in effort to stabilize patient.

Patient Presentation

Inclusion (Suspect Exposure) Criteria

1. Presentation may vary depending on the concentration and duration of exposure. Signs and symptoms may include, but are not limited to, the following:
 - a. Absorption:
 - i. Nausea
 - ii. Vomiting
 - iii. Diarrhea
 - iv. Altered mental status
 - v. Abdominal pain
 - vi. Rapid heart rate
 - vii. Dyspnea
 - viii. Wheezing
 - ix. Seizures
 - x. Arrhythmias
 - xi. Respiratory depression
 - xii. Sweating
 - xiii. Tearing
 - xiv. Defecation
 - xv. Constricted or dilated pupils
 - xvi. Rash
 - xvii. Burns to the skin
 - b. Ingestion:
 - i. Nausea
 - ii. Vomiting
 - iii. Diarrhea
 - iv. Altered mental status
 - v. Abdominal pain
 - vi. Rapid or slow heart rate
 - vii. Dyspnea
 - viii. Seizures

- ix. Arrhythmias
- x. Respiratory depression
- xi. Chemical burns around or inside the mouth
- xii. Abnormal breath odors
- c. Inhalation:
 - i. Nausea
 - ii. Vomiting
 - iii. Diarrhea
 - iv. Altered mental status
 - v. Abnormal skin color
 - vi. Dyspnea
 - vii. Seizures
 - viii. Burns to the respiratory tract
 - ix. Stridor
 - x. Sooty sputum
 - xi. Known exposure to toxic or irritating gas
 - xii. Respiratory depression
 - xiii. Sweating
 - xiv. Tearing
 - xv. Constricted or dilated pupils
 - xvi. Dizziness
- d. Injection:
 - i. Local pain
 - ii. Puncture wounds
 - iii. Reddening skin
 - iv. Local edema
 - v. Numbness
 - vi. Tingling
 - vii. Nausea
 - viii. Vomiting
 - ix. Diarrhea
 - x. Altered mental status
 - xi. Abdominal pain
 - xii. Seizures
 - xiii. Muscle twitching
 - xiv. Hypoperfusion
 - xv. Respiratory depression
 - xvi. Metallic or rubbery taste
- 1. **Toxidromes** (constellations of signs and symptoms that add in the identification of certain classes of medications and their toxic manifestations). These toxidrome constellations may be masked or obscured in poly pharmacy events:
 - a. Anticholinergic
 - i. Red as a beet (flushed skin)
 - ii. Dry as a bone (dry skin)
 - iii. Mad as a hatter (altered mental status)
 - iv. Blind as a bat (mydriasis)
 - v. Hot as a pistol (hyperthermia)
 - vi. Full as a flask (urinary retention)

- vii. "Tachy" like a pink flamingo (tachycardia and hypertension)
- b. Cholinergic (DUMBELS)

DUMBELS is a mnemonic used to describe the signs and symptoms of acetylcholinesterase inhibitor agent poisoning. All patient age groups are included where the signs and symptoms exhibited are consistent with the toxidrome of DUMBELS.

 - i. **D**iarrrhea
 - ii. **U**rination
 - iii. **M**iosis or **M**uscle weakness
 - iv. **B**ronchospasm, **B**ronchorrhea, **B**radycardia (the killer Bs)
 - v. **E**mesis
 - vi. **L**acrimation
 - vii. **S**alivation or **S**weating
- c. Opioids
 - i. Respiratory depression
 - ii. Miosis (pinpoint pupils)
 - iii. Altered mental status
 - iv. Decreased bowel sounds
- d. Sedative Hypnotic
 - i. Central nervous system depression
 - ii. Ataxia (unstable gait or balance)
 - iii. Slurred speech
 - iv. Normal or depressed vital signs (pulse, respirations, blood pressure)
- e. Stimulants (Sympathomimetic)
 - i. Tachycardia, tachydysrhythmias
 - ii. Hypertension
 - iii. Diaphoresis
 - iv. Delusions or paranoia
 - v. Seizures
 - vi. Hyperthermia
 - vii. Mydriasis (dilated pupils)
- f. Serotonin Syndrome (presentation with at least three of the following)
 - i. Agitation
 - ii. Ataxia
 - iii. Diaphoresis
 - iv. Diarrhea
 - v. Hyperreflexia
 - vi. Mental status changes
 - vii. Myoclonus
 - viii. Shivering
 - ix. Tremor
 - x. Hyperthermia
 - xi. Tachycardia

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Make sure the scene is safe. Use environmental Carbon Monoxide (CO) detector on "first in" bag if possible *[ALL EMS PRACTICE LEVELS-O]*
2. Consider body substance isolation (BSI) or appropriate PPE.
3. Assess ABCD and, if indicated, expose patient for assessment, and then re-cover to assure retention of body heat.
4. Check vital signs including temperature.
5. Attach ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]* and examine rhythm strip for arrhythmias (consider 12-lead ECG) *[Acquisition EMT-O; Interpretation INT-R]*.
6. Check blood glucose level *[EMR-O; EMT-R]*.
7. Monitor pulse oximetry *[EMR-O; EMT-R]* and ETCO₂ *[Acquisition EMT-O; Interpretation INT-O/PARA-R]* for respiratory decompensation.
8. Perform carboxyhemoglobin device assessment *[ALL EMS PRACTICE LEVELS-O]*, if available.
9. Identify specific medication taken, when indicated (including immediate release vs sustained release), time of ingestion, dose, and quantity. When appropriate, bring all medications (prescribed and not prescribed) in the environment.
10. Obtain an accurate ingestion history (as patient may become unconscious before arrival at ED):
 - a. Time of ingestion
 - b. Route of exposure
 - c. Quantity of medication or toxin taken (safely collect all possible medications or agents)
 - d. Alcohol or other intoxicant taken
11. Consider the threat to yourself and the destination facility if bringing in exposure agent.
12. Obtain pertinent cardiovascular history and other prescribed medications.
13. Check for needle marks, paraphernalia, bites, bottles, or evidence of agent involved in exposure, self-inflicted injury, or trauma.
14. Check for weapons and drugs (law enforcement should have checked, but you may decide to re-check).
15. Obtain pertinent patient history.
16. Perform physical examination.

Treatment and Interventions

1. Assure a patent airway.
2. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
3. Initiate IV access for infusion treatment medication and/or lactated Ringer's or normal saline if indicated *[AEMT-R]*.
4. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*) if evidence of hypoperfusion.
5. Administer appropriate antidote or mitigating medication (refer to specific agent guideline if not listed below).
 - a. Acetaminophen overdose:
 - i. Consider activated charcoal without sorbitol PO only if within the first

- hour of ingestion and prolonged transport to definitive care.
 - ii. Do not administer oral agents if there is a risk of rapidly decreasing mental status.
- b. Aspirin overdose:
 - i. Consider activated charcoal without sorbitol (1 gm/kg) PO.
 - 1. Administer charcoal early (highly recommended), as aspirin is erratically absorbed.
 - 2. Do not administer oral agents including activated charcoal if altered mental status or risk of rapid decreasing mental status from polypharmacy.
 - ii. Avoid manual ventilation in salicylate poisonings. Let the patient breathe on their own, even if tachypnea, until there is evidence of decompensation or dropping oxygen saturation. Acid or base disturbances and outcomes worsen when the patient is manually ventilated.
- c. Benzodiazepine overdose:
 - i. Provide respiratory support.
 - ii. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
 - iii. Consider vasopressors after adequate fluid resuscitation (1–2 liters of crystalloid) for the hypotensive patient.
- d. Caustic substances ingestion (e.g. acids and alkali):
 - i. Evaluate for airway compromise secondary to spasm or direct injury associated with oropharyngeal burns.
 - ii. In the few minutes immediately after ingestion, consider administration of water or milk if available.
 - 1. Adults: maximum 240 mL (8 ounces)
 - 2. Pediatrics: maximum 120 mL (4 ounces) to minimize risk of vomiting
 - 3. Do not attempt dilution in patients with respiratory distress, altered mental status, severe abdominal pain, nausea or vomiting, or patients who are unable to swallow or protect their airway.
 - 4. Do not force fluids in anyone who refuses to drink.
- e. Dystonia (symptomatic), extrapyramidal signs or symptoms, or mild allergic reactions
 - i. Consider administration of diphenhydramine *[PARA-O]*.
- f. Monoamine oxidase inhibitor overdose—symptomatic; e.g. MAOI; isocarboxazid (Marplan®), phenelzine (Nardil®), selegiline (Emsam®), tranylcypromine (Parnate®)
 - i. Consider administration of midazolam (benzodiazepine of choice) for temperature control.
- g. Opiate overdose: Treat per the [Opioid Poisoning/Overdose guideline](#).
- h. Oral ingestion unknown poisoning:
 - i. Do not administer oral agents if there is a risk of rapidly decreasing mental status or for petroleum-based ingestions.
 - ii. Consider administration of activated charcoal without sorbitol, particularly if it is within the first 1 hour after ingestion (including acetaminophen), or prolonged transport to definitive care.
 - iii. Administer activated charcoal to patients who have ingested

medications with extended release or delayed absorption.

- i. Selective serotonin reuptake inhibitors (SSRIs)
 - i. Consider early airway management.
 - ii. Treat arrhythmias following ACLS guidelines.
 - iii. Aggressively control hyperthermia with cooling measures.
 - iv. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
 - v. Consider vasopressors after adequate fluid resuscitation (1–2 liters of crystalloid) for the hypotensive patient [see [Shock guideline](#)].
 - vi. Consider midazolam (benzodiazepine of choice) for agitation.
 - vii. Treat seizures per [Seizures guideline](#).
- j. Tricyclic Antidepressant (TCA) Overdose:
 - i. Consider early airway management.
 - ii. Consider sodium bicarbonate if widened QRS (100 msec or greater).
 - iii. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
 - iv. Consider vasopressors after adequate fluid resuscitation (1–2 liters of crystalloid) for the hypotensive patient *[INT-R]* [see [Shock guideline](#)].
 - v. Consider midazolam *[INT-O; PARA-R]* (benzodiazepine of choice) for agitation.
 - vi. Treat seizures per [Seizures guideline](#).

Patient Safety Considerations

- Consider scene or environmental safety for patient and provider: Consider environmental carbon monoxide monitor use.
- Monitor patient airway, breathing, pulse oximetry, ETCO₂ for adequate ventilation as they may change over time.
- Repeat vital signs often.
- Monitor level of consciousness.
- Monitor ECG with special attention to rate, rhythm, QRS and QT duration.
- Maintain or normalize patient temperature.
- Engage the regional poison center as early as reasonably possible to aid in appropriate therapy and to track patient outcomes to improve knowledge of toxic effects. The national 24-hour toll-free telephone number to poison control centers is (800) 222- 1222, and it is a resource for free, confidential expert advice from anywhere in the United States.

Notes and Educational Pearls

Key Considerations

1. Each toxin or overdose has unique characteristics which must be considered in individual protocol.
2. Activated charcoal (which does not bind to all medications or agents) is still a useful adjunct in the serious agent, enterohepatic, or extended release agent poisoning as long as the patient does not have the potential for rapid alteration of mental status, or airway or aspiration risk. Precautions should be taken to avoid or reduce the risk of aspiration.
3. Ipecac is no longer recommended for any poisoning or toxic ingestion. The manufacturer has stopped production of this medication.
4. Flumazenil is not indicated in a suspected benzodiazepine overdose as you can precipitate refractory or intractable seizures if the patient is a benzodiazepine dependent patient.

Pertinent Assessment Findings

Frequent reassessment is essential as patient deterioration can be rapid and catastrophic.

Acetylcholinesterase Inhibitors (Carbamates, Nerve Agents, Organophosphates) Exposure

Aliases

Acetylcholinesterase inhibitor, ATNAA®, carbamate, Duodote®, insecticide, nerve agent, organophosphate, pesticide, weapons of mass destruction, WMD

Patient Care Goals

Rapid recognition of the signs and symptoms of confirmed or suspected acetylcholinesterase inhibitor (AChEI) agents such as carbamates, nerve agents, or organophosphates exposure followed by expeditious and repeated administration of atropine, the primary antidote

Note: Carbamates and organophosphates are commonly active agents in over-the-counter insecticides. Accidental carbamate exposure rarely requires treatment

Patient Presentation

Inclusion Criteria

1. **DUMBELS** is a mnemonic used to describe the signs and symptoms of acetylcholinesterase inhibitor agent poisoning. All patient age groups are included where the signs and symptoms exhibited are consistent with the toxidrome of DUMBELS.
 - a. **D**iarrhea
 - b. **U**rination
 - c. **M**iosis or **M**uscle weakness
 - d. **B**ronchospasm, **B**ronchorrhea, **B**radycardia (the killer B's)
 - e. **E**mesis
 - f. **L**acrimation
 - g. **S**alivation or **S**weating

Exclusion Criteria

No recommendations

Patient Management

1. Don the appropriate PPE; ensure rescuers are properly trained and equipped.
2. Remove the patient's clothing and wash the skin with soap and water. Be aware that:
 - a. Acetylcholinesterase inhibitor agents can be absorbed through the skin.
 - b. Contaminated clothing can provide a source of continued exposure to the toxin.
3. Rapidly assess the patient's respiratory status, mental status, and pupillary status.
4. Administer the antidote **[ALL EMS PRACTICE LEVELS-O]** immediately for confirmed or suspected acetylcholinesterase inhibitor agent exposure.
5. Administer oxygen **[EMR-O; EMT-R]** as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
6. Provide airway management.
7. Establish intravenous access **[AEMT-R]** (if possible).
8. Apply a ECG cardiac monitor (if available).

9. Recognize that the heart rate may be normal, bradycardic, or tachycardic.
10. Characterize clinical improvement based upon the drying of secretions and easing of respiratory effort rather than heart rate or pupillary response.
11. Reassess patient status continuously; this is critical.

Assessment Considerations

1. Acetylcholinesterase inhibitor agents are highly toxic chemical agents and can rapidly be fatal.
2. Patients with low-dose chronic exposures may have a more delayed presentation of symptoms.
3. Antidotes (atropine and pralidoxime) are effective if administered before circulation fails.
4. The patient may develop:
 - a. Miosis (pinpoint pupils).
 - b. Bronchospasm.
 - c. Bradycardia.
 - d. Vomiting.
 - e. Excessive secretions in the form of:
 - i. Tearing.
 - ii. Salivation.
 - iii. Rhinorrhea.
 - iv. Diarrhea.
 - v. Urination.
 - vi. Bronchorrhea.
5. Penetration of an acetylcholinesterase inhibitor agent into the central nervous system (CNS) will cause:
 - a. Headache.
 - b. Confusion.
 - c. Generalized muscle weakness.
 - d. Seizures.
 - e. Lethargy or unresponsiveness.
6. Estimated level of exposure based upon signs and symptoms
 - a. Mild
 - i. Miosis alone (while this is a primary sign in vapor exposure, it may not be present in all exposures)
 - ii. Miosis and severe rhinorrhea
 - b. Mild to moderate (in addition to symptoms of mild exposure)
 - i. Localized swelling
 - ii. Muscle fasciculation
 - iii. Nausea and vomiting
 - iv. Weakness
 - v. Shortness of breath
 - c. Severe (in addition to symptoms of mild to moderate exposure)
 - i. Unconsciousness
 - ii. Convulsions
 - iii. Apnea or severe respiratory distress requiring assisted ventilation
 - iv. Flaccid paralysis
7. Onset of symptoms can be immediate with an exposure to a large amount of the acetylcholinesterase inhibitor.
 - a. There is usually an asymptomatic interval of minutes after liquid exposure

- before these symptoms occur.
- b. Effects from vapor exposure occur almost immediately.
- 8. Signs and symptoms with large acetylcholinesterase inhibitor agent exposures (regardless of route)
 - a. Sudden loss of consciousness
 - b. Seizures
 - c. Copious secretions
 - d. Apnea
 - e. Death
- 9. Obtain an accurate ingestion history (as patient may become unconscious before arrival at ED):
 - a. Time of ingestion or exposure
 - b. Route of exposure
 - c. Quantity of medication or toxin taken (safely collect all possible medications or agents)
 - d. Alcohol or other intoxicant taken
 - e. Pertinent cardiovascular history or other prescribed medications for underlying disease
- 10. The patient can manifest any or all of the signs and symptoms of the toxidrome based on the route of exposure, agent involved, and concentration of the agent:
 - a. Vapor exposures will have a direct effect on the eyes and pupils causing miosis.
 - b. Patients with isolated skin exposures will have normally reactive pupils.
 - c. Certain acetylcholinesterase inhibitor agents can place the patient at risk for both a vapor and skin exposure.

Treatment and Interventions (see dosing tables below)

- 1. Medications:
 - a. Atropine
 - i. Atropine is the primary antidote for organophosphate, carbamate, or nerve agent exposures, and repeated doses should be administered liberally to patients who exhibit signs and symptoms of exposure or toxicity.
 - ii. Atropine may be provided in multi-dose vials, pre-filled syringes, or auto-injectors.
 - iii. Commercially available atropine auto-injectors **[ALL EMS PRACTICE LEVELS-O]** include:
 - 1. Atro-Pen® 1 mg of atropine (dark red container)
 - 2. Atro-Pen® 2 mg of atropine (green container)
 - 3. Pediatric Atro-Pen® 0.25 mg of atropine (yellow container)
 - 4. Pediatric Atro-Pen® 0.5 mg of atropine (blue container)
 - b. Pralidoxime chloride (2-PAM) **[ALL EMS PRACTICE LEVELS-O]**
 - i. Pralidoxime chloride is a secondary treatment and should be given concurrently in an effort to reactivate the acetylcholinesterase.
 - ii. Pralidoxime chloride may be provided in a single dose vial, pre-filled syringes, or auto-injectors.
 - iii. Auto-injectors contain 600 mg of pralidoxime chloride.
 - iv. In order to be beneficial to the victim, a dose of pralidoxime chloride should be administered shortly after the nerve agent or organophosphate poisoning as it has minimal clinical effect if administration is delayed.

- c. Benzodiazepines
 - i. Benzodiazepines are administered as an anticonvulsant for those patients who exhibit seizure activity [see [Seizures guideline](#) for doses and routes of administration].
 - ii. Lorazepam, diazepam, and midazolam are the most frequently used benzodiazepines in the prehospital setting.
 - iii. In the scenario of an acetylcholinesterase inhibitor agent exposure, the administration of diazepam or midazolam is preferable due to their more rapid onset of action.
 - iv. Benzodiazepines may be provided in multi-dose or single-dose vials, pre-filled syringes, or auto-injectors.
 - v. CANA[®] (Convulsive Antidote Nerve Agent) is a commercially available auto-injector that contains 10 mg of diazepam **[ALL EMS PRACTICE LEVELS-O]**.
- d. Mark I[®] Kits **[ALL EMS PRACTICE LEVELS-O]**
 - i. Mark I[®] Kits is a commercially available kit of nerve agent or organophosphate antidote auto-injectors. These are being phased out and replaced with Duodote by the CDC.
 - ii. A Mark I[®] kit consists of one auto-injector containing 2 milligrams of atropine and a second auto-injector containing 600 milligrams of pralidoxime chloride.
- e. Duodote[®] **[ALL EMS PRACTICE LEVELS-O]**
 - i. Duodote[®] is a commercially available auto-injector of nerve agent/organophosphate antidote.
 - ii. Duodote[®] is one auto-injector that contains 2.1 milligrams of atropine and 600 milligrams of pralidoxime chloride.
- f. ATNAA[®] (Antidote Treatment Nerve Agent Auto-injector) **[ALL EMS PRACTICE LEVELS-O]**
 - i. ATNAA[®] is an auto-injector of nerve agent or organophosphate antidote that is typically in military supplies.
 - ii. ATNAA[®] is one auto-injector that contains 2.1 milligrams of atropine and 600 milligrams of pralidoxime chloride.
 - iii. ATNAA[®] may be seen in civilian supplies assets when Duodote[®] is unavailable or in short supply.
- g. CHEMPACK **[ALL EMS PRACTICE LEVELS-O]**
 - i. CHEMPACK is a federally owned cache of nerve agent antidotes that is managed by the Centers for Disease Control and Prevention (CDC) and offered to states that voluntarily agree to maintain custody and security of CHEMPACK assets.
 - ii. These are forward-deployed at sites determined by states that are part of the program such as hospitals and EMS centers.
 - iii. Deployment of CHEMPACKs are reserved for events where the nerve agent or organophosphate exposure will deplete the local or regional supply of antidotes.
 - iv. There are two types of CHEMPACK containers:
 - 1. EMS Containers: CHEMPACK assets for EMS contain a large portion of auto-injectors for rapid administration of antidotes by EMS providers of all levels of licensure/certification. They contain enough antidote to treat roughly 454 patients.

2. Hospital Containers: CHEMPACK assets contain a large portion of multidose vials and powders for reconstitution. They contain enough antidote to treat roughly 1000 patients.
2. Medication Administration:
 - a. Atropine in extremely large, and potentially multiple, doses is the antidote for an acetylcholinesterase inhibitor agent poisoning.
 - b. Atropine should be administered immediately followed by repeated doses until the patient's secretions resolve (see table below).
 - c. Pralidoxime chloride (2-PAM) is a secondary treatment and, when possible, should be administered concurrently with atropine.
 - d. The stock of atropine and pralidoxime chloride available to EMS providers is usually not sufficient to fully treat the victim of an acetylcholinesterase inhibitor agent exposure; however, EMS providers should initiate the administration of atropine and, if available, pralidoxime chloride.
 - e. Seizures should be treated with benzodiazepines. There is some emerging evidence that, for midazolam, the intranasal route of administration may be preferable to the intramuscular route. However, intramuscular absorption may be more clinically efficacious than the intranasal route in the presence of significant rhinorrhea.
 - f. The patient should be emergently transported to the closest appropriate medical facility as directed by on-line medical control.
3. Recommended Doses (see dosing tables below)

The medication dosing tables that are provided below are based upon the severity of the clinical signs and symptoms exhibited by the patient. There are several imperative factors to note:

 - a. For organophosphate or severe acetylcholinesterase inhibitor agent exposure, the required dose of atropine necessary to dry secretions and improve the respiratory status is likely to exceed 20 mg. Atropine should be administered rapidly and repeatedly until the patient's clinical symptoms diminish. Atropine must be given until the acetylcholinesterase inhibitor agent has been metabolized. It may require up to 2000 mg of atropine over several days to weeks.
 - b. Since the antidotes in the Mark I[®] kit are in two separate vials, the atropine auto- injector in the kit can be administered to the patient in the event that Atro-Pen[®] or generic atropine auto-injectors are not available and/or intravenous atropine is not an immediate option.
 - c. Due to the fact that Duodote[®] auto-injectors contain pralidoxime chloride, they should not be used for additional dosing of atropine beyond the recommended administered dose of pralidoxime chloride.
 - d. All of the medications below can be administered intravenously in the same doses cited for the intramuscular route. However, due to the rapidity of onset of signs, symptoms, and potential death from acetylcholinesterase inhibitor agents, intramuscular administration is highly recommended to eliminate the inherent delay associated with establishing intravenous access.
 - e. The antidotes can be administered via the intraosseous route. However, due to the rapidity of onset of signs, symptoms, and potential death from acetylcholinesterase inhibitor agents, intramuscular administration remains preferable due to the inherent delay associated with establishing intraosseous access and the limited use of this route of administration for other

medications.

**Mild Acetylcholinesterase Inhibitor Agent
Exposure**

Patient (Weight)	Atropine Dose IM or via Auto- Injector
Infant: 0–2 yo	0.05 mg/kg IM or via auto-injector (<i>e.g. 0.25 and/or 0.5 mg auto-injector(s)</i>)
Child: 3–7 yo (13–25 kg)	1 mg IM or via auto-injector (<i>e.g. one 1 mg or two 0.5 mg auto-injectors</i>)
Child: 8–14 yo (26–50 kg)	2 mg IM or via auto-injector (<i>e.g. one 2 mg or two 1 mg auto-injectors</i>)
Adolescent/ Adult	2 mg IM or via auto-injector
Pregnant Women	2 mg IM or via auto-injector
Geriatric/ Frail	1 mg IM or via auto-injector

Mild to Moderate Acetylcholinesterase Inhibitor Agent Exposure

Patient (Weight)	Atropine Dose IM or via Auto- Injector	Pralidoxime Chloride Dose IM or via 600 mg Auto-Injector
Infant: 0–2 years	0.05 mg/kg IM or via auto-injector (<i>e.g. 0.25 mg and/or 0.5 mg auto-injector</i>)	15 mg/kg IM
Child: 3–7 yo (13–25 kg)	1 mg IM or via auto-injector (<i>e.g. one 1 mg auto-injector or two 0.5 mg auto-injectors</i>)	15 mg/kg IM OR One auto-injector (600 mg)
Child: 8–14 yo (26–50 kg)	2 mg IM or via auto-injector (<i>e.g. one 2 mg auto-injector or two 1 mg auto-injectors</i>)	15 mg/kg IM OR One auto-injector (600 mg)
Adolescent/ Adult	2–4 mg IM or via auto-injector	600 mg IM OR One auto-injector (600 mg)
Pregnant Women	2–4 mg IM or via auto-injector	600 mg IM OR One auto-injector (600 mg)

Geriatric/ Frail	2 mg IM or via auto-injector	10 mg/kg IM OR One auto-injector (600 mg)
-----------------------------	------------------------------	--

Severe Acetylcholinesterase Inhibitor Agent Exposure

Patient (Weight)	Atropine Dose IM or via Auto-Injector	Pralidoxime Chloride Dose IM or via 600 mg Auto-Injector
Infant: 0–2 yo	0.1 mg/kg IM or via auto-injector (<i>e.g. 0.25 mg and/or 0.5 mg auto-injector</i>)	45 mg/kg IM
Child: 3–7 yo (13–25 kg)	0.1 mg/kg IM OR 2 mg via auto-injector (<i>e.g. one 2 mg auto-injector or four 0.5 mg auto-injectors</i>)	45 mg/kg IM OR One auto-injector (600mg)
Child: 8–14 yo (26–50 kg)	4 mg IM or via auto-injector (<i>e.g. two 2 mg auto-injectors or four 1 mg auto-injectors</i>)	45 mg/kg IM OR Two auto-injectors (1200 mg)
Adolescent: 14 yo or older	6 mg IM or via auto-injector (<i>e.g. three 2 mg auto-injectors</i>)	Three auto-injectors (1800 mg)
Adult	6 mg IM or via auto-injector (<i>e.g. three 2 mg auto-injectors</i>)	Three auto-injectors (1800 mg)
Pregnant Women	6 mg IM or via auto-injector (<i>e.g. three 2 mg auto-injectors</i>)	Three auto-injectors (1800 mg)
Geriatric/Frail	2-4 mg IM or via auto-injector (<i>e.g. one to two 2 mg auto-injectors</i>)	25 mg/kg IM OR two to three auto-injectors (1200 mg–1800 mg)

Guidance for the Treatment of Seizures Secondary to Acetylcholinesterase Inhibitor Agent Exposure

Patient	Diazepam	Midazolam
Infant (0–2 yo)	0.2–0.5 mg/kg IM Repeat every 2-5 minutes	0.2 mg/kg IM Repeat prn in 10 minutes
	0.2–0.5 mg/kg IV every 15-30 minutes May repeat twice as needed	May repeat dose once
	Total maximum dose: 5 mg	Total maximum dose: 0.4 mg/kg
Child (3–13 yo)	0.2–0.5 mg/kg IM Repeat every 2–5 minutes	0.2 mg/kg IM Not to exceed 10 mg Repeat prn in 10 minutes
	0.2–0.5 mg/kg IV every 15-30 minutes May repeat dose twice if needed	May repeat dose once
	Total maximum dose: 5 mg if less than 5 years	Total maximum dose: 0.4 mg/kg Not to exceed 20 mg
	Total maximum dose: 10 mg if age 5 years or older 1 CANA [®] auto-injector	
Adolescent (14 yo or older)	2–3 CANA [®] auto-injectors	0.2 mg/kg IM Total maximum dose of 10 mg Repeat prn in 10 minutes
	5–10 mg IV every 15 minutes	May repeat dose once
	Total maximum dose: 30 mg	Total maximum dose: 20 mg
Adult	2–3 CANA [®] auto-injectors	10 mg IM Repeat prn in 10 minutes
	5–10 mg IV every 15 minutes	May repeat dose once
	Total maximum dose: 30 mg	Total maximum dose: 20 mg
Pregnant Women	2–3 CANA [®] auto-injectors	10 mg IM Repeat prn in 10 minutes
	5–10 mg IV every 15 minutes	May repeat dose once
	Total maximum dose: 30 mg	Total maximum dose: 20 mg
Geriatric	2–3 CANA [®] auto-injectors	10 mg IM Repeat prn in 10 minutes
	5–10 mg IV every 15 minutes	May repeat dose once
	Total maximum dose: 30 mg	Total maximum dose: 20 mg

Tables adapted from: U.S. Department of Health and Human Services, ASPR, National Library of Medicine, *Chemical Hazards Emergency Medical Management: Nerve Agents-*

Prehospital Management, www.chemm.nlm.nih.gov

Patient Safety Considerations

- Continuous and ongoing patient reassessment is critical.
- Clinical response to treatment is demonstrated by the drying of secretion and the easing of respiratory effort.
- Initiation of and ongoing treatment should not be based upon heart rate or pupillary response.
- Precautions for pralidoxime chloride administration:

Although Duodote® and ATNAA® contain atropine (the primary antidote for an acetylcholinesterase inhibitor agent poisoning), the inclusion of pralidoxime chloride in the auto-injector can present challenges if additional doses of atropine are warranted by the patient condition and other formulations of atropine are unavailable:

 - Pediatrics: An overdose of pralidoxime chloride may cause profound neuromuscular weakness and subsequent respiratory depression.
 - Adults: Especially for the geriatric victim, excessive doses of pralidoxime chloride may cause severe systolic and diastolic hypertension, neuromuscular weakness, headache, tachycardia, and visual impairment.
 - Geriatrics: In a victim who may have underlying medical conditions, particularly impaired kidney function or hypertension, the EMS provider should consider administering the lower recommended adult dose of intravenous pralidoxime chloride.
- Considerations during the use of auto-injectors
 - If an auto-injector is administered, a dose calculation prior to administration is not necessary.
 - For atropine, additional auto-injectors should be administered until secretions diminish.
 - Mark I® kits, Duodote® and ATNAA® have not been approved for pediatric use by the Food and Drug Administration (FDA), but they can be considered for the initial treatment of children of any age with severe symptoms of an Acetylcholinesterase inhibitor agent poisoning, especially if other formulations of atropine are unavailable.
 - Pediatric Atro-Pen® auto-injectors are commercially available in a 0.25 mg auto-injector (yellow) and a 0.5 mg auto-injector (red). Atro-Pen® auto-injectors are commercially available in a 1 mg auto-injector (blue) and a 2 mg auto-injector (green).
 - A pralidoxime chloride 600 mg auto-injector may be administered to an infant that weighs greater than 12 kg.

Notes and Educational Pearls

Key Considerations

- Clinical effects of acetylcholinesterase inhibitor agents
 - The clinical effects are caused by the inhibition of the enzyme acetylcholinesterase which allows excess acetylcholine to accumulate in the nervous system.
 - The excess accumulated acetylcholine causes hyperactivity in muscles, glands, and nerves.

- Organophosphates (certain insecticides)
 - Organophosphates can be legally purchased by the general public.
 - Organophosphates (e.g. pesticides) penetrate tissues and bind to the patient's body fat producing a prolonged period of illness and ongoing toxicity even during aggressive treatment.
- Nerve agents
 - Traditionally classified as weapons of mass destruction (WMD)
 - Not readily accessible to the general public
 - Extremely toxic and rapidly fatal with any route of exposure
 - GA (tabun), GB (sarin), GD (soman), GF, and VX are types of nerve agents and are WMDs
 - Nerve agents can persist in the environment and remain chemically toxic for a prolonged period of time.

Pertinent Assessment Findings

The signs and symptoms exhibited with the toxidrome of DUMBELS [see [Patient Presentation – Inclusion Criteria](#) above]

Radiation Exposure

Aliases

None noted

Patient Care Goals

1. Prioritize identification and treatment of immediately life-threatening medical conditions and traumatic injuries above any radiation-associated injury.
2. Identify and appropriately treat acute radiation injury.
3. Reduce risk for contamination of personnel while caring for patients potentially or known to be contaminated with radioactive material.

Patient Presentation

Inclusion Criteria

1. Patients who have been acutely exposed to ionizing radiation from accidental environmental release of a radioactive source
2. Patients who have been acutely exposed to ionizing radiation from a non-accidental environmental release of a radioactive source
3. Patients who have been contaminated with material emitting ionizing radiation

Exclusion Criteria

1. Patients exposed to normal doses of ionizing radiation from medical imaging studies
2. Patients exposed to normal doses of ionizing radiation from therapeutic medical procedures

Patient Management

Assessment

1. Identify and treat life-threatening injuries and medical problems; this takes priority over decontamination.
2. Don PPE capable of preventing: skin exposure to liquids and solids (gown and gloves); mucous membrane exposure to liquids and particles (face mask and eye protection); and inhalational exposure to particles (N95 face mask or respirator).
3. Do not eat or drink any food or beverages while caring for patients with radiation injuries until screening is completed for contamination, and appropriate decontamination performed, if needed.
4. Use caution to avoid dispersing contaminated materials.
5. Provide appropriate condition-specific care for any immediately life-threatening injuries or medical problems.

Treatment and Interventions

1. If patient experiences nausea, vomiting, and/or diarrhea:
 - a. Provide care, per [Nausea-Vomiting guideline](#).
 - b. Document the time gastrointestinal symptoms started.
2. If seizure occurs:
 - a. Consider a primary medical cause or exposure to possible chemical agents unless indicators for a large whole body radiation dose (greater than 20Gy), such as

- rapid onset of vomiting, are present.
- b. Treat per [Seizures guideline](#).

Patient Safety Considerations

Treat life-threatening medical problems and traumatic injuries prior to assessing for and treating radiation injuries or performing decontamination.

Notes and Educational Pearls

Key Considerations

- Irradiated patients pose no threat to medical providers.
- Contaminated patients pose very little threat to medical providers who use appropriate PPE including N95 masks or respirators, gloves, gowns, and face and eye protection.
- Sources of radiation
 - Legal
 - Industrial plants
 - Health care facilities that provide radiologic services
 - Nuclear power plants
 - Mobile engineering sources (e.g. construction sites that are installing cement)
 - Illegal
 - Weapons of mass destruction
 - "Dirty bomb" design to contaminate widespread areas
- Physiology of radiation poisoning
 - Contamination: Poisoning from direct exposure to a radioactive source, contaminated debris, liquids, or clothing where radiation continues to be emitted from particles on surface
 - Exposure: Poisoning from radioactivity, in the form of ionizing rays, penetrating through the bodily tissues of the patient
- Common types of radioactivity that cause poisoning
 - Gamma rays
 - Highest frequency of ionizing rays
 - Penetrates the skin deeply
 - Causes the most severe radiation toxicity
 - Beta rays: Can penetrate up to 1 cm of the skin's thickness
 - Alpha rays
 - Lowest frequency of ionizing rays
 - Short range of absorption
 - Dangerous only if ingested or inhaled
 - Radioactive daughters
 - Products of decay of the original radioactive substance
 - Can produce gamma and beta rays (e.g. uranium decays into a series of radon daughters)
- In general, trauma patients who have been exposed to or contaminated by radiation should be triaged and treated on the basis of the severity of their conventional injuries.
- A patient who is contaminated with radioactive material (e.g. flecks of radioactive material embedded in their clothing and skin) generally poses a minimal exposure risk to medical personnel.

- EMS providers may be asked to assist public health agencies in the distribution and administration of potassium iodide in a mass casualty incident involving radiation release or exposure.

Pertinent Assessment Findings

- Treatment of life-threatening injuries or medical conditions takes priority over assessment for contamination or initiation of decontamination.
- Time to nausea and vomiting is a reliable indicator of the received dose of ionizing radiation. The more rapid the onset of vomiting, the higher the whole-body dose of radiation.
- Tissue burns are a late finding (weeks following exposure) of ionizing radiation injury. If burns are present acutely, they are from a thermal or chemical mechanism.
- Seizures may suggest acute radiation syndrome if accompanied by early vomiting. If other clinical indicators do not suggest a whole-body dose of greater than 20Gy, consider other causes of seizure.
- Delayed symptoms (days to weeks after exposure or contamination)
 - Skin burns with direct contact with radioactive source
 - Skin burns or erythema from ionizing rays
 - Fever
 - Bone marrow suppression presenting as:
 - Immunosuppression
 - Petechiae
 - Spontaneous internal and external bleeding

Topical Chemical Burn

Aliases

Chemical Burn

Patient Care Goals

1. Rapid recognition of a topical chemical burn
2. Initiation of emergent and appropriate intervention and patient transport

Patient Presentation

Inclusion Criteria

1. Patients of all ages who have sustained exposure to a chemical that can cause a topical chemical burn may develop immediate or in some cases a delayed clinical presentation.
2. Agents that are known to cause chemical burns include alkali, acids, mustard agent, and lewisite.

Exclusion criteria

None recommended

Patient Management

1. Don the appropriate PPE; ensure rescuers are properly trained and equipped.
2. Remove the patient's clothing, if necessary.
3. Place contaminated clothing in double bags.
4. Transport patient by EMS providers if deemed necessary and manpower resources permit. The EMS providers should be those who did not participate in the decontamination process, and transport should occur in an emergency response vehicle that has not been exposed to the chemical.
5. Gather information regarding the chemical while on scene, including materials safety data sheet if available.
6. Communicate all data regarding the chemical to the receiving facility.

Assessment

1. Be aware that clinical effects and severity of a topical chemical burn is dependent upon:
 - a. Class of agent (alkali injury or acid injury).
 - b. Concentration of the chemical the (higher the concentration, the greater the risk of injury).
 - c. pH of the chemical:
 - i. Alkali-increased risk with pH greater than or equal to 11
 - ii. Acid-increased risk with pH less than or equal to 3
 - d. Onset of burn
 - i. Immediate
 - ii. Delayed (e.g. hydrofluoric acid)
2. Calculate the estimated total body surface area that is involved.
3. Prevent further contamination.
4. Provide special attention to assessment of ocular or oropharyngeal exposure; evaluate for airway compromise secondary to spasm or direct injury associated

with oropharyngeal burns.

5. Recognize that some acid and alkali agents may manifest systemic effects.

Treatment and Interventions

1. If dry chemical contamination, carefully brush off solid chemical prior to flushing the site as the irrigating solution may activate a chemical reaction.
2. If wet chemical contamination, flush the patient's skin (and eyes, if involved) with copious amounts of water or normal saline.
3. Provide adequate analgesia per the [Pain Management guideline](#).
4. Consider the use of topical anesthetic eye drops (e.g. tetracaine) for chemical burns of the eye.
5. For eye exposure, administer continuous flushing of irrigation fluid to eye **[ALL LEVELS-O]**. Morgan lens may facilitate administration **[PARA-O]**.
6. Administer early airway intervention for airway compromise or spasm associated with oropharyngeal burns.
7. Take measures to minimize hypothermia.
8. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline **[AEMT-R]** or lactated Ringer's **[AEMT-O]**).

Hydrofluoric Acid

Hydrofluoric acid (HF) is a highly corrosive substance that is primarily used for automotive cleaning products, rust removal, porcelain cleaners, etching glass, cleaning cement or brick, or as a pickling agent to remove impurities from various forms of steel. Hydrofluoric acid readily penetrates intact skin and there may be underlying tissue injury. It is unlikely that low concentration HF will cause an immediate acid-like burn, however there may be delayed onset of pain to the exposed area. Higher concentration HF may cause immediate pain as well as more of a burn appearance that can range from mild erythema to an obvious burn. An oral or large dermal exposure can result in significant systemic hypocalcemia with possible QT prolongation and cardiovascular collapse.

1. For all patients in whom a hydrofluoric acid exposure is confirmed or suspected:
 - a. Vigorously irrigate all affected areas with water or normal saline for a minimum of 15 minutes.
 - b. Apply ECG cardiac monitor **[Acquisition EMT-O; Interpretation INT-R]** for oral or large dermal exposures significant HF exposures.
 - c. Apply calcium preparation:
 - i. Calcium prevents tissue damage from hydrofluoric acid.
 - ii. Topical calcium preparations:
 1. Commercially manufactured calcium gluconate gel
 2. If commercially manufactured calcium gluconate gel is not available, a topical calcium gluconate gel preparation can be made by combining 150 mL (5 ounces) of a sterile water-soluble gel (e.g. Surgilube® or KY® jelly) with one of the following **[PARA-R]** :
 - a. 35 mL of calcium gluconate 10% solution
 - b. 10 g of calcium gluconate tablets (e.g. Tums®)
 - c. 3.5 g calcium gluconate powder or
 3. If calcium gluconate is not available, 10 mL of calcium chloride 10% solution in 150 mL in sterile water soluble gel (e.g.

- Surgilube® or KY® jelly)
4. Apply generous amounts of the calcium gluconate gel to the exposed skin sites to neutralize the pain of the hydrofluoric acid.
 - a. Leave the gel in place for at least 20 minutes then reassess.
 - b. This can be repeated as needed.
 5. Although generally low yield, there may be benefit to intravenous pain medication along with the topical calcium gluconate gel for pain control.
 6. If fingers are involved, apply the calcium gel to the hand: Squirt additional calcium gel into a surgical glove, and then insert the affected hand into the glove.
 7. For patients who have ingested hydrofluoric acid or who have a large dermal exposure, consider intravenous calcium gluconate as symptomatic hypocalcemia can precipitate rapidly as manifest by muscle spasms, seizures, hypotension ventricular arrhythmias and QT prolongation.

Patient Safety Considerations

1. Don PPE.
2. Take measures to prevent the patient from further contamination through decontamination.
3. Take measures to protect the EMS provider and others from contamination.
4. Do not attempt to neutralize an acid with an alkali or an alkali with an acid as an exothermic reaction will occur and cause serious thermal injury to the patient.
5. Consider expeditious transport or transfer to a designated burn center for burns that involve a significant percentage of total body surface area or burns that involve the eyes, face, hands, feet or genitals.

Notes and Educational Pearls

Key Considerations

- IV fluid resuscitation should be guided by patient age and clinical status.
- Since the severity of topical chemical burns is largely dependent upon the type, concentration, and pH of the chemical involved as well as the body site and surface area involved, it is imperative to obtain as much information as possible while on scene about the chemical substance by which the patient was exposed. The information gathering process will often include:
 - Transport of the “sealed” container of the chemical to the receiving facility.
 - Transport of the original or a copy of the Material Safety Data Sheet (MSDS) of the substance to the receiving facility.
 - Contacting the reference agency to identify the chemical agent and assist in management (e.g. CHEMTREC®).
- Inhalation of HF should be considered in any dermal exposure involving the face and neck or if clothing is soaked in the product.
- Decontamination is critical for both acid and alkali agents to reduce injury. Removal of chemicals with a low pH (acids) is more easily accomplished than chemicals with a high pH (alkalis) because alkalis tend to penetrate and bind to deeper tissues.

- Some chemicals will also manifest local and systemic signs, symptoms, and bodily damage.

Pertinent Assessment Findings

- An estimate of the total body surface area that is involved
- Patient response to therapeutic interventions
- Patient response to fluid resuscitation
- Patient response to analgesia

Stimulant Poisoning or Overdose

Aliases

Stimulant, cocaine, methamphetamine, amphetamines, PCP, phencyclidine, bath salts

Patient Care Goals

1. Identify intoxicating agent.
2. Protect organs at risk for injury such as heart, brain, liver, kidney.
3. Determine if there is an antidote.
4. Treat the symptoms which may include severe tachycardia and hypertension, agitation, hallucinations, chest pain, seizure, and arrhythmia.

Patient Presentation

Inclusion Criteria

1. Tachycardia or tachydysrhythmias
2. Hypertension
3. Diaphoresis
4. Delusions or paranoia
5. Seizures
6. Hyperthermia
7. Mydriasis (dilated pupils)
8. Stimulant or hallucinogenic (with stimulant properties) agents:
 - a. Cocaine
 - b. Amphetamine or methamphetamine
 - c. Phencyclidine (PCP) (hallucinogen)
 - d. Bupropion
 - e. Synthetic stimulant drugs of abuse (some having mixed properties)
 - f. Ecstasy
 - g. Methamphetamine
 - h. Synthetic cathinones (bath salts)
 - i. Spice
 - j. K2
 - k. Synthetic THC
 - l. Khat

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Begin with the ABCDs:
 - a. Airway is patent
 - b. Breathing is oxygenating
 - c. Circulation is perfusing
 - d. Mental status is stable
2. Treat any compromise of these parameters.
3. Ask about chest pain and difficulty breathing.

4. Check vital signs including temperature for hyperthermia.
5. Apply a ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*].
6. Check blood glucose level [*EMR-O; EMT-R*].
7. Monitor ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] for respiratory decompensation.
8. Check a 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*] when possible.
9. Check for trauma, self-inflicted injury.
10. Check for weapons and drugs (law enforcement should have already performed this check, but you may decide to repeat the inspection).

Treatment and Interventions

1. Establish IV access for any fluids and meds [*AEMT-R*].
2. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]) [see [Shock](#) and [Hyperthermia/Heat Exposure](#) guidelines].
3. Treat chest pain as ACS and follow STEMI protocol if there is ECG is consistent with STEMI.
4. Consider treating shortness of breath as atypical ACS.
 - a. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
5. Consider soft physical management devices [*EMR-O; EMT-R*] especially if law enforcement has been involved in getting patient to cooperate [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].
6. Consider medications to reduce agitation and other significant sympathomimetic findings for the safety of the patients and providers. This may improve behavior and compliance [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].
7. Consider prophylactic use of anti-emetic ondansetron [*INT-O*].
8. If hyperthermia suspected, begin external cooling.

Patient Safety Considerations

- Apply the least amount of physical management devices that are necessary to protect the patient and the providers [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].
- Assess for potential weapons or additional drugs; this is very important since these items can pose a threat not just to the patient but also to the EMS crew.

Notes and Educational Pearls

Key Considerations

- Recognition and treatment of hyperthermia (including sedatives to decrease heat production from muscular activity) is essential as many deaths are attributable to hyperthermia.
- If law enforcement has placed the patient in handcuffs, this patient needs ongoing physical security for safe transport. Have law enforcement in back of ambulance for the handcuffed patient or make sure proper physical management devices are in place before law enforcement leaves and ambulance departs from scene.
- If patient has signs and symptoms of ACS, strive to give nitroglycerin [*AEMT-R*].
 - Vasospasm is often the problem in this case as opposed to a fixed coronary artery lesion.
 - Consider administration of benzodiazepines as if to treat anxiety.

- Maintaining IV access [*AEMT-R*], ECG cardiac monitor, and SPO₂/ETCO₂ monitors are key to being able to catch and intervene decompensations in a timely manner.
- If agitated, consider restraining the patient to facilitate patient assessment and lessen likelihood of vascular access or monitor displacements.
- Cocaine has sodium channel blocking effects and can cause significant cardiac conduction abnormalities with a widened QRS. Treatment is with sodium bicarbonate similar to a tricyclic antidepressant. Check a 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*] to assess for these complications.

Pertinent Assessment Findings

- History is as important as the physical examination.
- If the patient is on psychiatric medication, but has failed to be compliant, this fact alone puts the patient at higher risk for excited delirium.
- If the patient is found naked, this may elevate the suspicion for stimulant use or abuse and increase the risk for excited delirium. Neuroleptic malignant syndrome, serotonin syndrome and excited delirium can present in with similar signs and symptoms.
- If polypharmacy is suspected, hypertension and tachycardia are expected hemodynamic findings secondary to increased dopamine release. Stimulus reduction from benzodiazepines, anti-psychotics, and ketamine will improve patient's vital signs and behavior.
- Be prepared for the potential of cardiovascular collapse as well as respiratory arrest.
- If a vasopressor is needed, epinephrine or norepinephrine is recommended over dopamine.

Cyanide Exposure

Aliases

Cyanide, hydrogen cyanide, blood agent

Patient Care Goals

1. Remove patient from toxic environment.
2. Assure adequate ventilation, oxygenation and correction of hypoperfusion.

Patient Presentation

Cyanide is a colorless, "bitter almond smell" (genetically only 40% of population can smell) gas or white crystal which binds to the ferric ion in cells, blocking the enzyme cytochrome oxidase, thus preventing the use of oxygen by the cell's mitochondria, leading to cellular hypoxia.

Inclusion Criteria

1. Depending on its form, cyanide can enter the body through inhalation, ingestion, or absorption through the skin. Cyanide should be suspected in occupational or other smoke exposures (e.g. firefighting), industrial accidents, natural catastrophes, suicide and murder attempts, chemical warfare and terrorism (whenever there are multiple casualties of an unclear etiology). Non-specific and early signs of cyanide exposure (inhalation, ingestion, or absorption) include the following signs and symptoms: anxiety, vertigo, weakness, headache, tachypnea, nausea, dyspnea, vomiting, and tachycardia.
2. High concentrations of cyanide will produce:
 - a. Markedly altered level of consciousness, including rapid collapse.
 - b. Seizures.
 - c. Respiratory depression or respiratory arrest.
 - d. Cardiac dysrhythmias (other than sinus tachycardia).
3. The rapidity of onset is related to the severity of exposure (inhalation or ingestion) and may be dramatic with immediate effects that include early hypertension with subsequent hypotension, sudden cardiovascular collapse, seizure or coma, and rapid death.

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Remove patient from toxic environment when rescuers are properly trained and equipped.
2. Assess ABCDs and, if indicated, expose the patient, and then re-cover the patient to assure retention of body heat.
3. Assess vital signs including temperature and pulse oximetry *[EMR-O; EMT-R]* (which may not correlate with tissue oxygenation in cyanide/smoke exposure).
4. Attach a ECG cardiac monitor and obtain a 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
5. Check blood glucose level *[EMR-O; EMT-R]*.

6. Monitor pulse oximetry [*EMR-O; EMT-R*] and ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*].
7. Monitor patient for signs of hypoxia (pulse oximetry *less than 94%*) [*EMR-O; EMT-R*] and respiratory decompensation regardless of pulse oximetry reading.
8. Identify the specific agent of exposure, time of ingestion or inhalation, and quantity or timing of exposure.
9. Obtain patient history including cardiovascular history and prescribed medication.
10. Obtain other pertinent patient history.
11. Perform physical exam.

Treatment and Interventions

There is no widely available, rapid, confirmatory cyanide blood test. Many hospitals will not be able to rapidly assess cyanide levels. Therefore, treatment decisions must be made on the basis of clinical history and signs and symptoms of cyanide intoxication. For the patient with an appropriate history and manifesting one or more significant cyanide exposure signs or symptoms:

1. Treat with 100% oxygen [*EMR-O; EMT-R*] via non-rebreather mask or bag valve mask.
2. Collect a pre-treatment blood sample in the appropriate tube for lactate and cyanide levels [*AEMT-O*] *if possible*.
3. Administer one of the following medication regimes:
 - a. Hydroxocobalamin (the preferred agent) [*PARA-O*]
 - OR**
 - b. Sodium thiosulfate [*PARA-O*].
4. Treat seizures per [Seizures guideline](#).

Patient Safety Considerations

- In the event of multiple casualties, be sure to wear appropriate PPE during rescue evacuation from the toxic environment.
- If the patient ingests cyanide, it will react with the acids in the stomach generating hydrogen cyanide gas. Be sure to maximize air circulation in closed spaces (ambulance) as the patient's gastric contents may contain hydrogen cyanide gases when released with vomiting or belching.
- Do not use nitrites in conjunction with suspected carbon monoxide poisoning as it worsens the hemoglobin oxygen carrying capacity even more than carbon monoxide (CO).
- Hydroxocobalamin is the only agent safe for treatment of cyanide poisoning in pregnant patient.

Notes and Educational Pearls

Key Considerations

- Pulse oximetry accurately reflects serum levels of oxygen but does not accurately reflect tissue oxygen levels; therefore, it should not be relied upon in possible cyanide and/or carbon monoxide toxicity.
- After hydroxocobalamin has been administered, pulse oximetry levels are no longer accurate.
- If the patient ingests cyanide, it will react with the acids in the stomach generating hydrogen cyanide gas. Be sure to maximize air circulation in closed spaces (ambulance) as the patient's gastric contents may contain hydrogen cyanide gases

- when released with vomiting or belching.
- Amyl nitrite and sodium nitrite are no longer being used and no longer available in commercial kits.

Pertinent Assessment Findings

Early and repeated assessment is essential

Beta Blocker Poisoning or Overdose

Aliases

Anti-hypertensive

Patient Care Goals

1. Reduce GI absorption of oral agents with some form of binding agent (activated charcoal) especially for extended release.
2. Provide early airway protection; this is required as patients may have rapid mental status deterioration.
3. Assure adequate ventilation, oxygenation, and correction of hypoperfusion.

Patient Presentation

Beta blocker or beta adrenergic antagonist medication to reduce the effects of epinephrine or adrenaline

Inclusion Criteria

1. Patients may present with:
 - a. Bradycardia
 - b. Hypotension
 - c. Altered mental status
 - d. Weakness
 - e. Shortness of breath
 - f. Possible seizures
2. Beta blocker agents examples:
 - a. Acebutolol hydrochloride (Sectral®)
 - b. Atenolol (Tenormin®)
 - c. Betaxolol hydrochloride (Kerlone®)
 - d. Bisoprolol fumarate (Zebeta®)
 - e. Carteolol hydrochloride (Cartrol®)
 - f. Esmolol hydrochloride (Brevibloc®)
 - g. Metoprolol (Lopressor®, Toprol XL®)
 - h. Nadolol (Corgard®)
 - i. Nebivolol (Bystolic®)
 - j. Penbutolol sulfate (LevatoI®)
 - k. Pindolol (Visken®)
 - l. Propranolol (Inderal®, InnoPran®)
 - m. Timolol maleate (Blocadren®)
 - n. Sotalol hydrochloride (Betapace®)
3. Alpha/beta-adrenergic blocking agents examples:
 - a. Carvedilol (Coreg®)
 - b. Labetalol hydrochloride (Trandate®, Normodyne®)

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Assess ABCDs and, if indicated, expose and then cover to assure retention of body heat.
2. Check vital signs, including temperature.
3. Apply a ECG cardiac monitor, examine rhythm strip for arrhythmias, and consider obtaining a 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*].
4. Check blood glucose level [*EMR-O; EMT-R*].
5. Monitor pulse oximetry [*EMR-O; EMT-R*] and ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] for respiratory decompensation.
6. Identify specific medication taken (noting immediate release vs. sustained release formulations), time of ingestion, and quantity.
7. Obtain a pertinent cardiovascular history or other prescribed medications for underlying disease.
8. Obtain pertinent patient history.
9. Administer patient physical.

Treatment and Interventions

1. Consider activated charcoal without sorbitol [*EMT-O*].
 - a. If risk of rapid decreasing mental status, do not administer oral agent without adequately protecting the airway.
2. Check blood glucose level [*EMR-O; EMT-R*] on all patients, but especially on pediatric patients as beta blockers can cause hypoglycemia in pediatric population.
3. Consider atropine sulfate for symptomatic bradycardia [*INT-R*].
4. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]).
5. For symptomatic patients with cardiac effects (i.e. hypotension, bradycardia) consider Glucagon [*INT-O*].
6. Consider vasopressors after adequate fluid resuscitation (1–2 liters of crystalloid) for the hypotensive patient [see [Shock guideline](#) for pediatric vs. adult dosing].
7. Consider transcutaneous pacing if refractory to initial pharmacologic interventions [*INT-R*].
8. Treat seizures per [Seizures guideline](#).
9. If widened QRS (100 msec or greater), consider sodium bicarbonate [*PARA-R*]. This can be repeated as needed to narrow QRS.

Patient Safety Considerations

- Transcutaneous pacing may not always capture or correct hypotension when capture is successful.
- Aspiration of activated charcoal can produce a patient where airway management is nearly impossible. Do not administer activated charcoal to any patients that may have a worsening mental status.

Notes and Educational Pearls

Key Considerations

1. **Pediatric Considerations:**
 - a. Pediatric patient may develop hypoglycemia from beta blocker overdose therefore it is important to perform glucose evaluation.
 - b. A single pill can kill a toddler. It is very important that a careful assessment of medications the toddler could have access to be done by EMS and all suspect medications should be brought into the ED.

2. All Patient Considerations

- a. Glucagon has a side effect of increased vomiting. Ondansetron prophylaxis should be considered.
- b. Atropine may have little or no effect (likely to be more helpful in mild overdoses). The hypotension and bradycardia may be mutually exclusive and the blood pressure may not respond to correction of bradycardia.
- c. Propranolol crosses the blood brain barrier and can cause altered mental status, seizure, and widened QRS similar to TCA toxicity.

Pertinent Assessment Findings

- Certain beta blockers, such as acebutolol and propranolol, may increase QRS duration.
- Certain beta blockers, such as acebutolol and pindolol, may produce tachycardia and hypertension.
- Sotalol can produce increase in QTc interval and ventricular dysrhythmia.
- Frequent reassessment is essential as patient deterioration can be rapid and catastrophic.

Bites and Envenomation

Aliases

Stings

Patient Care Goals

Bites, stings, and envenomation can come from a variety of insects, marine and terrestrial animals. There is a spectrum of toxins or envenomation with very limited EMS interventions.

1. Assure adequate ventilation, oxygenation and correction of hypoperfusion.
2. Maintain pain control which also includes limited external interventions to reduce pain.

Patient Presentation

Inclusion Criteria

1. Bites, stings, and envenomation can come from a variety of marine and terrestrial animals and insects causing local or systemic effects.
2. Patients may present with toxin specific reactions which may include:
 - a. Site pain
 - b. Swelling
 - c. Muscle pain (hallmark of black widow spider bites)
 - d. Erythema
 - e. Discoloration
 - f. Bleeding
 - g. Nausea
 - h. Abdominal pain
 - i. Hypotension
 - j. Tachycardia
 - k. Tachypnea
 - l. Muscle incoordination
 - m. Confusion
 - n. Anaphylaxis or allergic reactions
3. There is a spectrum of toxins or envenomation and limited EMS interventions that will have any mitigating effect on the patient in the field.
 - a. The critical intervention is to get the patient to a hospital that has access to the antivenin if applicable.

Exclusion Criteria

None

Patient Management

Assessment

1. Assess ABCDs and, if indicated, expose and then cover to assure retention of body heat.
2. Monitor vital signs which include temperature.
3. Apply a ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]*, examine rhythm strip for arrhythmias, and consider obtaining a 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]*.
4. Check blood glucose Level *[EMR-O; EMT-R]*.

5. Monitor pulse oximetry [*EMR-O; EMT-R*] and ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] for respiratory decompensation.
6. Obtain pertinent patient history.
7. Conduct patient physical with special consideration to area of envenomation or bite.

Treatment and Interventions

1. Consider an IV Normal saline fluid bolus (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]) 20 mL/kg up to 2 liters.
2. Consider vasopressors after adequate fluid resuscitation for the hypotensive patient [for adult vs. pediatric dosing see [Shock guideline](#)].
3. Treat seizures per [Seizures guideline](#).
4. Administer specific therapy for select bites, stings, or envenomation:
 - a. Crotalidae (vipers such as the timber rattlesnake and the eastern massasauga rattlesnake) are found in Wisconsin. Antivenin for this type of snake is commercially available; consider transport to hospital that has access to antivenin, if feasible.
5. Provide adequate analgesia per the [Pain Management guideline](#).

Patient Safety Considerations

- Do *not*:
 - Apply tourniquets, tight Ace® or crepe bandage, or constricting bands above or below the site of the envenomation.
 - Perform incision and/or suction wound to remove toxin.
 - Apply cold packs or immerse the affected extremity in ice water (cryotherapy).
 - Try to capture the offending marine or terrestrial animal or insect
- If the offending organism has been killed, be aware that many dead insect, marine, or fanged animals can continue to bite or sting with venom and should be safely placed in a hard sided and closed container for future identification.
- Patient may still have an imbedded stinger, tooth, nematocyst, or barb which may continue to deliver toxin if left imbedded. Consider safe removal without squeezing the toxin delivery apparatus.

Notes and Educational Pearls

Key Considerations

Vinegar has potential to increase pain associated with jellyfish sting as it can increase nematocysts discharge in certain species. Providers must be familiar with endemic species and how to best address exposure.

Pertinent Assessment Findings

1. Assess for signs and symptoms of local and systematic impact of the suspected toxin.
2. Be aware that patient may still have an imbedded stinger, tooth, nematocysts, or barb which may continue to deliver toxin if left imbedded.

Calcium Channel Blocker Poisoning or Overdose

Aliases

Anti-hypertensive

Patient Care Goals

1. Reduce GI absorption of oral agents with some form of binding agent (activated charcoal) especially for extended release.
2. Provide early airway protection; this is required as patients may have rapid mental status deterioration.
3. Assure adequate ventilation, oxygenation and correction of hypoperfusion.

Patient Presentation

Calcium channel blockers interrupt the movement of calcium across cell membranes. Calcium channel blockers are used to manage hypertension, certain rate-related arrhythmias, prevent cerebral vasospasm, and angina pectoris. Patients may present with:

1. Bradycardia
2. Hypotension
3. Decreased AV Nodal conduction
4. Cardiogenic shock
5. Hyperglycemia

Inclusion Criteria

1. Patients who have may have taken or been administered calcium channel blockers
 - a. Calcium channel blocker examples:
 - i. Amlodipine (Norvasc®)
 - ii. Diltiazem (Cardizem®, Tiazac®)
 - iii. Felodipine
 - iv. Isradipine
 - v. Nicardipine
 - vi. Nifedipine (Adalat CC®, Afeditab CR®, Procardia®)
 - vii. Nisoldipine (Sular®)
 - viii. Verapamil (Calan®, Verelan®)

Exclusion criteria

No recommendations

Patient Management

Assessment

1. Assess ABCDs and, if indicated, expose and then cover to assure retention of body heat.
2. Monitor vital signs, including temperature.
3. Apply a ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*], and consider obtaining a 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*].
4. Check blood glucose Level [*EMR-O; EMT-R*].
5. Monitor pulse oximetry [*EMR-O; EMT-R*] and ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] for respiratory decompensation.
6. Identify specific medication taken (noting immediate release vs. sustained

- release formulations), time of ingestion, and quantity.
7. Obtain pertinent cardiovascular history or other prescribed medications for underlying disease.
 8. Patient pertinent history.
 9. Physical exam.

Treatment and Interventions

1. Consider activated charcoal without sorbitol. If risk of rapid decreasing mental status, do not administer oral agent without adequately protecting the airway.
2. Consider atropine sulfate for symptomatic bradycardia *[INT-R]*.
3. Consider calcium gluconate (preferred) *[PARA-R]*.
4. Consider IV fluid bolus (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*) 20 mL/kg up to 2 liters *[AEMT-R]*.
5. Consider vasopressors after adequate fluid resuscitation for the hypotensive patient *[INT-R]* [see [Shock guideline](#) for adult vs. pediatric dosing] if atropine, calcium, and vasopressors have failed in the symptomatic bradycardia patient
6. Consider transcutaneous pacing *[INT-R]* if refractory to initial pharmacologic interventions.
7. If seizure, consider midazolam *[INT-O; PARA-R]* (benzodiazepine of choice).

Patient Safety Considerations

Transcutaneous pacing may not always capture nor correct hypotension when capture is successful.

Notes and Educational Pearls

Key Considerations

- While most calcium channel blockers cause bradycardia, dihydropyridine class calcium channel blockers (e.g. nifedipine, amlodipine) can cause a reflex tachycardia (torsade de pointes) early in the ingestion. The patient can become bradycardic as the intoxication worsens.
- Administering calcium chloride or calcium gluconate to a patient on cardiac glycosides (e.g. digoxin) is an appropriate intervention. It is now accepted that the traditional avoidance of this intervention (due to its supposed potential to precipitate toxicity and associate fatal arrhythmias) is a historical belief and is not supported.
- Glucagon has a side effect of increased vomiting. Ondansetron prophylaxis should be considered.
- A single pill can kill a toddler. It is very important that a careful assessment of medications the toddler could have access to is done by EMS, and suspect medications brought into the ED.
- Calcium channel blockers can cause many types of rhythms that can range from sinus bradycardia to complete heart block.
- Hyperglycemia is the result of the blocking of L-type calcium channels in the pancreas. This can help differentiate these ingestions from beta blockers. There may also be a relationship between the severity of the ingestion and the extent of the hyperglycemia.
- Atropine may have little or no effect (likely to be more helpful in mild overdoses).
- Hypotension and bradycardia may be mutually exclusive and the blood pressure

may not respond to correction of bradycardia.

Pertinent Assessment Findings

- Conduct close monitoring of ECG and document changes and dysrhythmias
- Serial frequent assessments are essential as these patients often have rapid deterioration with profound hypotension.

Carbon Monoxide or Smoke Inhalation

Aliases

CO

Patient Care Goals

1. Remove patient from toxic environment.
2. Assure adequate ventilation, oxygenation and correction of hypoperfusion.
3. Consider use of environmental carbon monoxide (CO) monitors on “first in” bags to assist in detection of occult CO toxicity.

Patient Presentation

Carbon monoxide is a colorless, odorless gas which has a high affinity for binding to red cell hemoglobin, thus preventing the binding of oxygen to the hemoglobin, leading to hypoxia (pulse oximetry less than 94%). A significant reduction in oxygen delivery to tissues and organs occurs with carbon monoxide poisoning. Carbon monoxide is also a cellular toxin which can result in delayed or persistent neurologic sequelae in significant exposures. With any form of combustion (fire or smoke [e.g. propane, kerosene, or charcoal stoves or heaters], combustion engines [e.g. generators, lawn mowers, motor vehicles, home heating systems]), carbon monoxide will be generated. People in a fire may also be exposed to cyanide from the combustion of some synthetic materials. Cyanide toxicity may need to be considered in the hemodynamically unstable patient removed from a fire.

Inclusion Criteria

1. Patients exposed to carbon monoxide may present with a spectrum of symptoms:
 - a. Mild intoxication:
 - i. Nausea
 - ii. Fatigue
 - iii. Headache
 - iv. Vertigo
 - v. Lightheadedness
 - b. Moderate to severe:
 - i. Altered mental status
 - ii. Tachypnea
 - iii. Tachycardia
 - iv. Convulsion
 - v. Cardiopulmonary arrest

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Remove patient from toxic environment by rescuers who are properly trained and equipped.
2. Assess ABCDs and, if indicated, expose patient and re-cover to assure retention of body heat.
3. Monitor vital signs including pulse oximetry [*EMR-O*; *EMT-R*], temperature, and ETCO₂

- [Acquisition EMT-O; Interpretation INT-O/PARA-R]* if available.
4. Apply an ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]* and obtain a 12-lead ECG *[Acquisition EMT-O; Interpretation INT-R]* if available.
 5. Check blood glucose level *[EMR-O; EMT-R]*
 6. Monitor pulse oximetry *[EMR-O; EMT-R]* and ETCO₂ for respiratory decompensation.
 7. Obtain pertinent patient history.
 8. Conduct patient physical examination.

Treatment and Interventions

1. Administer 100% oxygen *[EMR-O; EMT-R]* via non-rebreather mask or bag valve mask or advanced airway as indicated.
2. Treat seizures per [Seizures guideline](#).
3. Consider transporting patients with severe carbon monoxide poisoning directly to a facility with hyperbaric oxygen capabilities, if feasible and patient does not meet criteria for other specialty care (e.g. trauma or burn).

Patient Safety Considerations

- Consider affixing a carbon monoxide detector *[ALL EMS PRACTICE LEVELS-O]* to an equipment bag that is routinely taken into scene (if it signals alarm, don appropriate respiratory protection and exit scene) to assist with detection of occult CO toxicity.
- Remove patient and response personnel from potentially hazardous environment as soon as possible.
- Prohibit the patient, the patient's family, and other appropriate bystanders from entering the environment (e.g. building, car) where the carbon monoxide exposure occurred until the source of the poisoning has been eliminated.
- Do not look for cherry red skin coloration as an indication of carbon monoxide poisoning, as this is an unusual finding.
- Transport all patients with probable or suspected CO poisoning to the nearest appropriate hospital based on their presenting signs and symptoms. Do not depend on CO oximeter devices, as they may yield inaccurate results for patients with CO poisoning.

Notes and Educational Pearls

Key Considerations

- Pulse oximetry is inaccurate due to the carbon monoxide binding with hemoglobin.
- Pregnant patients are more likely to be treated successfully with hyperbaric oxygen as maternal carboxyhemoglobin levels do not accurately reflect fetal carboxyhemoglobin levels.
- Consider cyanide toxicity if carbon monoxide poisoning is from a fire.
- A patient light wavelength analysis device to detect carboxyhemoglobin is useful to indicate if there is a carbon monoxide exposure in a non-arrested patient. Do not anticipate an immediate change in readings with oxygen administration.

Pertinent Assessment Findings

- Early and repeat assessment of patient's mental status and motor function are extremely useful in determining response to therapy and the need for hyperbaric therapy.
- Identification and documentation of possible etiology of poisoning
- Time of symptom onset and time of initiation of exposure-specific treatment
- Response to therapy

Opioid Poisoning or Overdose

Aliases

Carfentanil, Dilaudid®, drug abuse, EVZIO®, fentanyl, heroin, hydrocodone, hydromorphone, methadone, morphine, naloxone, Narcan®, opiate, opioid, overdose, oxycodone, Oxycontin®, Percocet®, Percodan®, Suboxone, U-47700, Vicodin®

Patient Care Goals

1. Rapid recognition and intervention of a clinically significant opioid poisoning or overdose.
2. Prevention of respiratory and/or cardiac arrest.

Patient Presentation

Inclusion Criteria

Patients exhibiting miosis (pinpoint pupils), decreased mental status, and respiratory depression of all age groups with known or suspected opioid use or abuse.

Exclusion Criteria:

Patients with altered mental status exclusively from other causes (e.g. head injury, or hypoglycemia).

Patient Management

1. Don the appropriate PPE.
2. Initiate therapeutic interventions to support the patient's airway, breathing, and circulation prior to the administration of naloxone **[EMR-O; EMT-R]**.
3. Identify specific medication taken (including immediate release versus sustained release) time of ingestion, and quantity, if possible.
4. Obtain and document pertinent cardiovascular history or other prescribed medications for underlying disease.
5. Be aware that unsecured hypodermic needles may be on scene if the intravenous route may have been used by the patient, and that there is a higher risk of needle sticks during the management of this patient population which may also have an increased incidence of blood-borne pathogens.
6. Consider Naloxone, an opioid antagonist, for administration to patients with respiratory depression in a confirmed or suspected opioid overdose **[EMR-O; EMT-R]**.
7. Be aware that Naloxone administration:
 - Via the intravenous route provides more predictable bioavailability and flexibility in dosing and titration.
 - Via the intranasal or intramuscular routes or as a nebulized solution provides additional options of medication delivery.
8. If naloxone was administered to the patient prior to the arrival of EMS, obtain the dose and route through which it was administered and, if possible, bring the devices containing the dispensed naloxone with the patient along with all other medications on scene.


Assessment

1. Assess the patient's airway, breathing, circulation, and mental status.

2. Support the patient's airway by positioning, oxygen administration, and ventilator assistance with a bag valve mask if necessary.
3. Assess the patient for other etiologies of altered mental status including hypoxia (pulse oximetry less than 94%) [EMR-O; EMT-R], hypoglycemia [EMR-O; EMT-R], hypotension, and traumatic head injury.
4. Remove any adhesive patches from the skin (legally prescribed opioids manufactured for transdermal absorption), if found.

Treatments and Interventions

Figure 6. Opioid-Associated Emergency for Healthcare Providers Algorithm.



© 2020 American Heart Association

2020 American Heart Association. Available at https://cpr.heart.org/-/media/cpr-files/cpr-guidelines-files/highlights/hghlghts_2020_ecc_guidelines_english.pdf. Accessed 11/29/2020

1. Check responsiveness, call for help, get naloxone [EMR-O; EMT-R] and AED.
2. Assess breathing.
3. Prevent deterioration.
4. Conduct ongoing assessment.
5. Assess pulse.
6. Support ventilation; give naloxone [EMR-O; EMT-R].
7. Start CPR, use AED or ECG, refer to resuscitation algorithm, and consider naloxone [EMR-O; EMT-R].

Patient Safety Considerations

1. Clinical duration of naloxone
 - a. The clinical opioid reversal effect of naloxone is limited and may end within an hour whereas opioids often have a duration of 4 hours or longer.
 - b. Monitor the patient for recurrent respiratory depression and decreased mental status.

2. Opioid withdrawal
 - a. Be aware that patients with altered mental status secondary to an opioid overdose may become agitated or violent following naloxone administration due to opioid withdrawal. Therefore the goal is to use the lowest dose possible to avoid precipitating withdrawal.
 - b. Be prepared for this potential scenario and take the appropriate measures in advance to ensure and maintain scene safety.
3. EMS providers should be prepared to initiate airway management before, during, and after naloxone administration and to provide appropriate airway support until the patient has adequate respiratory effort.

Notes and Educational Pearls

Key Considerations

- The essential feature of opioid overdose requiring EMS intervention is respiratory depression or apnea.
- Some opioids have additional toxic effects (e.g. methadone can produce QT prolongation, and tramadol can produce seizures).
- Overuse and abuse of prescribed and illegal opioids has led to an increase in accidental and intentional opioid overdoses.
- DEA and Opioids:
 - Legally prescribed opioids are controlled under the Drug Enforcement Administration (DEA).
 - Opioids have a high potential for abuse, but have an accepted medical use in patient treatment and can be prescribed by a physician.
 - Common legally prescribed opioids include codeine, fentanyl, hydrocodone, morphine, hydromorphone, methadone, morphine, oxycodone, and oxymorphone.
 - Opioid derivatives, such as heroin, are illegal in the United States.
- Opioid combinations:
 - Some opioids are manufactured as a combination of analgesics with acetaminophen, acetylsalicylic acid (aspirin), or other substances.
 - In the scenario of an overdose, there is a potential for multiple drug toxicities.
 - Examples of opioid combination analgesics:
 - Vicodin® is a combination of acetaminophen and hydrocodone
 - Percocet® is a combination of acetaminophen and oxycodone
 - Percodan® is a combination of aspirin and oxycodone
 - Suboxone® is a combination of buprenorphine and naloxone
- High-potency opioids:
 - Fentanyl is 50–100 times more potent than morphine. It is legally manufactured in an injectable and oral liquid, tablet, and transdermal (worn as a patch) forms, however much of the fentanyl adulterating the heroin supply are illegal fentanyl analogs such as acetyl fentanyl.
 - Carfentanil is 10,000 times more potent than morphine.
 - It is legally manufactured in a liquid form, however, a powder or tablet is the most common form of this drug in illegal production.
 - In the concentration in which it is legally manufactured (3 mg/mL), an intramuscular dose of 2 mL of carfentanil will sedate an elephant.
 - Synthetic opioids (e.g. W-18, are 10,000 times more potent than morphine).

Many synthetic opioids are not detectable by routine toxicology screening assays.

- The IN route provides no risk of needle stick to the provider.
- Patients with opioid overdose from fentanyl or fentanyl analogs may rapidly exhibit chest wall rigidity and require positive end expiratory pressure (PEEP), in addition to multiple and/or larger doses of naloxone, to achieve adequate ventilation.
- PPE that provides additional cutaneous, respiratory, or ocular protection may be considered when providing care in jurisdictions experiencing an increased incidence of overdose from high potency opioids.

Pertinent Assessment Findings

- The primary clinical indication for the use of opioid medications is analgesia.
- In the opioid overdose scenario, signs and symptoms include:
 - Miosis (pinpoint pupils).
 - Respiratory depression.
 - Decreased mental status.
- Additional assessment precautions:
 - The risk of respiratory arrest with subsequent cardiac arrest from an opioid overdose as well as hypoxia (pulse oximetry *less than* 94%), hypercarbia, and aspiration may be increased when other substances such as alcohol, benzodiazepines, or other medications have also been taken by the patient.
 - **Pediatric Considerations:** The signs and symptoms of an opioid overdose may also be seen in newborns who have been delivered from a mother with recent or chronic opioid use. Neonates who have been administered naloxone for respiratory depression due to presumed intrauterine opioid exposure may be narcotic dependent and should be monitored closely for seizures.

Airway Respiratory Irritants

Aliases

Respiratory irritant, airway injury, respiratory injury, chemical respiratory injury, toxic inhalation

Patient Care Goals

Rapid recognition of the signs and symptoms of confirmed or suspected airway respiratory irritants.

Patient Presentation

Inclusion Criteria

1. Inhalation of a variety of gases, mists, fumes, aerosols, or dusts may cause irritation or injury to the airways, pharynx, lung, asphyxiation, or other systemic effects.
2. Inhaled airway or respiratory irritant agents will interact with the mucus membranes, upper and lower airways based on solubility, concentration, particle size, and duration of exposure.
3. The less soluble and smaller the particle size of the agent, the deeper it will travel into the airway and respiratory systems before reacting with adjoining tissues, thus causing a greater delay in symptom onset.

Exclusion Criteria

No recommendations

Signs and Symptoms

As the type, severity, and rapidity of signs and symptoms onset depends on agent, water solubility, concentration, particle size, and duration of exposure, the below signs and symptoms are often overlapping and escalating in severity.

1. Many airway and respiratory irritant agents have "warning properties" such as identifiable or unpleasant smells or irritation to eyes or airways.
2. Some agents do not have clear warning properties and will often have delayed onset of any sign or symptom:
 - a. Unusual odor or smell
 - b. Tearing or itchy eyes
 - c. Burning sensation and burns to the nose, pharynx and respiratory tract
 - d. Sneezing
 - e. General excitation
 - f. Cough
 - g. Chest tightness
 - h. Nausea
 - i. Shortness of breath or dyspnea
 - j. Wheezing
 - k. Stridor
 - l. Dyspnea on exertion
 - m. Dizziness
 - n. Change in voice
 - o. Airway obstruction includes laryngospasm and laryngeal edema
 - p. Pulmonary edema (non-cardiogenic)
 - q. Seizures

- r. Cardiopulmonary arrest
- 3. High water solubility or highly irritating (oral, nasal, and pharynx; particle size greater than 10 micrometers)
 - a. Acrolein
 - b. Ammonia
 - c. Chloramine
 - d. Ethylene oxide
 - e. Formaldehyde
 - f. Hydrogen chloride
 - g. Methyl bromide
 - h. Sodium azide
 - i. Sulfur dioxide
- 4. Intermediate water solubility (bronchus and bronchiole; particle size 5 to 10 micrometers)
 - a. Chlorine
- 5. Low water solubility and less irritating (alveolar, particle size less than 5 micrometers)
 - a. Cadmium fume
 - b. Fluorine
 - c. Hydrogen sulfide (rotten egg odor; olfactory fatigue)
 - d. Mercury fume
 - e. Mustard gas (also delayed blistering skin manifestations)
 - f. Nickel carbonyl
 - g. Ozone
 - h. Phosgene
- 6. Asphyxia agents (two categories)
 - a. Oxygen deprivation below 19.5% oxygen atmosphere ("simple asphyxiants")

Any gas that reduces oxygen fraction or displaces oxygen from the inspired air

 - i. Argon
 - ii. Carbon dioxide
 - iii. Ethane
 - iv. Helium
 - v. Methane
 - vi. Natural gas (e.g. heptane, propane)
 - vii. Nitrogen
 - viii. Nitrogen dioxide (delayed symptom onset)
 - b. Chemical interfering with oxygen delivery or utilization ("chemical asphyxiants")
 - i. Carbon monoxide [see [Carbon Monoxide/Smoke Exposure guideline](#)]
 - ii. Cyanide [see [Cyanide Exposure guideline](#)]
 - iii. Hydrogen sulfide
- 7. Inhalants of abuse
 - a. These agents or substances are a diverse class of substances that include volatile solvents, aerosols, and gases.
 - b. These chemicals are intentionally inhaled to produce a state that resembles alcohol intoxication with initial excitation, drowsiness, lightheadedness, and agitation.
 - c. The abusers of these inhaled agents are often called huffers, sniffers, baggers, or snorters.
 - i. These individuals often present after inhaling an aerosol or gas with a loss of consciousness, and with the presence of the aerosol can or

- residue or paint around or in the mouth, nose, and oral pharynx.
- d. Common household products that are used as inhalants of abuse:
 - i. Volatile solvents
 1. Paint remover
 2. Degreasers
 3. Dry-cleaning fluids
 4. Gasoline
 5. Lighter fluid
 6. Correction fluid
 7. Felt tip markers
 8. Glue
 - ii. Cosmetic or paint spray
 1. Deodorant spray
 2. Vegetable oil spray
 3. Fabric protector spray
 4. Spray paint
 - iii. Propellants, asphyxiants, nitrous oxide
 1. Propane gas
 2. Balloon tanks (helium)
 3. Computer keyboard cleaner
 4. Ether
 5. Halothane
 6. Chloroform
 7. Butane
 8. Propane
 9. Whipped cream dispensers
 8. Riot control agents [see [Riot Control Agent guideline](#)]
 9. A prototype agent is identified with each region of the effected airway respiratory tract for *mild to moderate exposures*, as severe concentrated exposures of many of these agents overlap in signs and symptoms. The deeper the symptoms are in the respiratory tract and the slower the rate of symptom onset, the less water soluble the airway respiratory irritant.
 - a. Nasal and oral pharynx irritation: highly water-soluble agents (ammonia)
 - b. Bronchial irritation (chlorine)
 - c. Acute pulmonary edema or deep alveolar injury: poorly water soluble (phosgene)
 - d. Direct neurotoxin (hydrogen sulfide)
 - e. Asphyxia agent with additional symptoms (nitrogen dioxide, Silo Filler's disease)
 - f. Inhalants of abuse (volatile solvents, cosmetics, paints, propellants, asphyxiants, nitrous oxide)
 - g. Riot control agents [see [Riot Control Agents guideline](#)]
 - h. Anticholinesterase inhibitors [see [Acetylcholinesterase Inhibitors guideline](#)]
 10. Ammonia
 - a. Immediate detection of unique sharp smell
 - b. Nasal pharyngeal burning or irritation sensation
 - c. Ocular tearing and irritation
 - d. Sneezing
 - e. Altered mental status: sleepy to agitated
 - f. Cough
 - g. Shortness of breath

- h. Chest tightness
- i. Bronchospasm wheezing
- j. Change in voice
- k. Upper airway obstruction includes laryngospasm and laryngeal edema
- l. Corneal burns or ulcers
- m. Skin burns
- n. Pharyngeal, tracheal, bronchial burns
- o. Dyspnea or tachypnea
- p. High concentrations and or protracted exposure may develop non-cardiac pulmonary edema
- q. Esophageal burns
- 11. Chlorine
 - a. All the above (Ammonia)
 - b. Increased likelihood of the following
 - i. Bronchiole burns
 - ii. Bronchospasm wheezing
 - iii. Non-cardiac pulmonary edema develops within 6 to 24 hours of higher exposures
- 12. Phosgene
 - a. Often have **none** of the above symptoms for first half hour to several hours, then are much milder until more severe lower respiratory tract symptoms develop
 - i. Only warning is report of "fresh mowed hay" odor
 - ii. Mild airway irritation or drying
 - iii. Mild eye irritation
 - iv. Fatigue
 - v. Chest tightness
 - vi. Dyspnea or tachypnea
 - vii. Significant delay up to 24 hours for
 - 1. Exertional dyspnea
 - 2. Bronchospasm wheezing
 - 3. Hypoxia
 - 4. Severe non-cardiac pulmonary edema
 - 5. Cardiopulmonary arrest
- 13. Hydrogen sulfide is a direct neurotoxin and is rapidly absorbed through lungs, generating systemic effects.
 - a. Distinctive rotten egg smell which rapidly causes olfactory fatigue or loss of sense of smell
 - b. Cough
 - c. Shortness of breath
 - d. Rapid alternations in cognition or consciousness
 - e. Bronchiole and lung hemorrhage or hemoptysis
 - f. Non-cardiac pulmonary edema
 - g. Hydrogen sulfide is known as the "knock down" gas because of near immediate and sudden loss of consciousness with high concentrations
 - h. Asphyxia
 - i. Death
- 14. Nitrogen dioxide (also called Silo Filler's disease)
 - a. Heavier than air, displaces oxygen from low lying areas and closed spaces

- causing direct asphyxia
- b. Low concentrations may cause
 - i. Ocular irritation
 - ii. Cough
 - iii. Dyspnea or tachypnea
 - iv. Fatigue
- c. High concentrations:
 - i. Altered mental status including agitation
 - ii. Cyanosis
 - iii. Vomiting
 - iv. Dizziness
 - v. Loss of consciousness
 - vi. Cardiopulmonary arrest
- 15. Inhalants of abuse (e.g. felt tip markers, spray paint)
 - a. Physical presences of paint or residue on individual from the inhaled agent
 - b. Slurred speech
 - c. Altered mental status (excitation, drowsiness to unconsciousness)
 - d. Loss of consciousness
 - e. Cardiac dysrhythmias
 - f. Cardiopulmonary arrest

Patient Management

1. Don appropriate PPE; respiratory protection is critical.
2. Remove patient from the toxic environment; rescuers should be properly trained and equipped.
 - a. Remove the patient's clothing that may retain gases or decontaminate if liquid or solid contamination.
 - b. Flush effected or burned areas.
3. Rapidly assess the patient's respiratory status, mental status, and oxygenation [*EMR-O; EMT-R*].
4. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
5. Establish intravenous access [*AEMT-R*] (if possible).
6. Apply a ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*] (if available).
7. Conduct continuous and ongoing patient reassessment; this is critical.

Assessment

1. Make sure the scene is safe as many gases are heavier than air and will build up in low lying areas. This is especially true of hydrogen sulfide and it's "knock down" effect of the initial unprotected responder. Be aware that subsequent casualties can occur when unprotected rescuers attempt to save the first downed responder.
2. Consider BSI or appropriate PPE.
3. Remove patient from toxic environment by rescuers who are properly trained and equipped.
4. Decontaminate; rescuers should be properly trained and equipped.
5. Assess ABCD and if indicated, expose the patient and then cover the patient to assure retention of body heat.

6. Monitor vital signs, including temperature.
7. Place ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*] and examine rhythm strip for arrhythmia potentials (consider 12-lead ECG).
8. Check blood glucose level.
9. Monitor pulse oximetry [*EMR-O; EMT-R*] and ETCO₂ [*Acquisition EMT-O; Interpretation INT-O/PARA-R*] for respiratory decompensation.
10. Perform carboxyhemoglobin and cyanide device assessment, if available.
11. Identify specific suspected agent if possible.
12. Obtain pertinent cardiovascular history or history of other prescribed medications for underlying disease.
13. Obtain pertinent patient history.
14. Conduct patient physical examination.

Treatment and Interventions

1. Assure a patent airway.
 - a. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
 - b. Maintain the airway and assess for airway burns, stridor, or airway edema and if indicated, perform intubation early [*INT-O; PARA-R*] (recommendation to avoid non-visualized airways); cricothyroidotomy [*PARA-O*] may be required in rarer severe cases.
 - c. Apply non-invasive ventilation techniques.
 - i. Use continuous positive airway pressure (CPAP) or bi-level positive airway pressure (BiPAP) [*EMT-O; AEMT-R*] or High Flow Nasal Cannula (HFNC) [*PARA-O*] for severe respiratory distress or impending respiratory failure.
 - ii. Use bag-valve-mask (BVM) ventilation in the setting of hypoventilation, respiratory failure or arrest.
2. Administer albuterol [*EMR-O; EMT-R*] to all patients in respiratory distress with signs of bronchospasm, either by basic life support BLS or ALS providers. This medication should be repeated at this dose with unlimited frequency for ongoing distress.
3. Administer ipratropium nebulized up to 3 doses [*EMT-O*], in conjunction with albuterol.
4. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]).
5. If the patient is experiencing significant pain, administer IV/IO analgesics:
 - a. Fentanyl [*INT-O*] or
 - b. Morphine [*INT-O*]
6. Administer early eye irrigation.
7. Treat topical chemical burns [see appropriate [Toxins and Environmental](#) section guideline(s)].
8. In severe respiratory irritation, in particular hydrogen sulfide, with altered mental status and no improvement with removal from the toxic environment, administer oxygen as appropriate with a target of achieving greater than 93% saturation [*EMR-O; EMT-R*]; consider consultation for transfer to a hyperbaric oxygen therapy.

Patient Safety Considerations

1. Generally, in speaking to patients with exposure to highly soluble airway or respiratory irritants, you will find that they have self-extricated due to the warning properties such as the smell, rapidity of onset of irritation, and other symptoms,
2. The less soluble agents may generate only an odor (e.g. mowed hay smell for Phosgene) symptom and will have delayed serious symptoms such as acute pulmonary edema, hypoxia, and shortness of breath with minimal exertion.

Notes and Educational Pearls

Key Considerations

- Airway respiratory irritants can exacerbate underlying reactive airway diseases (e.g. asthma, COPD) and precipitate or exacerbate bronchospasm, respiratory distress, and hypoxia.
- As patients may be off gassing (particularly hydrogen sulfide and hydrogen cyanide) in the back of the transport vehicle, it is recommended to have adequate ventilation of the patient compartment.
- Removal from the toxic environment, oxygen, general supportive therapy, bronchodilators, respiratory support, and time are core elements of care as there are no specific antidotes for any of these inhaled agents (with the exception of heavy metals that may be chelated by physicians after agent identification).
- Hydrogen sulfide causes the cells responsible for the sense of smell to be stunned into inaction. Therefore, with a very short exposure they will shut down and the exposed victim will not perceive the smell, yet the victim will continue to absorb the gas as it is still present.
- Inhaled agents have become popular as a means of committing suicide. If there is some form of suicide signage—hoses, or buckets of substances—visible as you arrive at the vehicle or residence, immediately retreat to a well ventilated area and don SCBA before opening the vehicle or making entry. These gases may be highly concentrated and potentially lethal to EMS responders.
- Household bathroom, kitchen, and oven cleaners, when mixed, can generate a variety of airway respiratory irritants (ammonia, chloramine, and chlorine gas releases are particularly common). A very common exposure is to chloramine, a gas liberated when bleach (hypochlorite) and ammonia are combined. Chloramine then hydrolyzes in the distal airways and alveoli to ammonia and hypochlorous acid.
- Sudden sniffing death can result from a single use of inhalant of abuse.
 - Some inhalants can cause the heart to beat rapidly and erratically and cause cardiac arrest.
 - This syndrome most often is associated with abuse of butane, propane and effects of the chemicals in the aerosols.

Pertinent Assessment Findings

- Patient may describe a specific odor (chlorine swimming pool smell, ammonia smell, fresh mowed hay smell [phosgene]) which may be helpful but should not be relied upon as the human nose is a poor discriminator of scent.
- Respiratory distress (retractions, wheezing, stridor)
- Decreased oxygen saturation
- Skin color
- Neurologic status assessment
- Reduction in work of breathing after treatment
- Improved oxygenation after breathing

Riot Control Agents

Aliases

CN (Mace®), CS, OC (pepper spray), tear gas, harassing agents, incapacitating agents, chemical crowd control agents, lacrimators

Patient Care Goals

1. Address side effects of exposed individuals.
2. Decontaminate affected individuals.
3. Minimize effect to provider.

Patient Presentation

Inclusion Criteria

1. Exposure to identifiable agents that are not intended to cause significant injury or fatality

Exclusion Criteria

1. Exposure to chlorine, phosgene, ammonia or other agents that are intended to cause significant injury or fatality
2. Exposure to an unknown agent

Patient Management

Assessment

1. Assess scene safety: Evaluate for hazards to EMS personnel, patient, and bystanders.
 - a. Determine riot control agent being used.
 - b. Don appropriate PPE.
 - c. Determine number of patients.
2. Note symptoms exhibited by the exposed individual.
3. Examine as appropriate to complaints.

Treatment and Interventions

1. Move affected individuals from contaminated environment into fresh air if possible.
2. Remove contaminated clothing as able.
3. Have patient remove contact lenses if appropriate.
4. Note that irrigation with water or saline may facilitate resolution of symptoms and is recommended for decontamination of dermal and ocular exposure.
5. If patient is in respiratory distress, go to [Respiratory](#) section.
6. If patient is wheezing, go to [Bronchospasm guideline](#).
7. For persistent pain of the eye or skin, go to [Topical Chemical Burn guideline](#).
8. Re-evaluate exposed individuals who are persistently symptomatic warrant further evaluation and treatment per local standards.

Patient Safety Considerations

- Toxicity is related to duration of exposure and concentration of agent used (exposure in non-ventilated space).
- Patients with pre-existing pulmonary conditions (e.g. asthma, COPD) may be prone to more severe respiratory effects.
- Traumatic injury may result when exposed individuals are in proximity to the device used to disperse the riot control agent (e.g. hose or stream under pressure, riot

control agent projectile, grenade).

Notes and Educational Pearls

Key Considerations

- CN, CS, and OC are the most commonly encountered riot control agents.
- CN, CS and OC have a high safety ratio. All three have a high median lethal concentration (LCt50) and a low median effective concentration (ECt50).
- Toxicity is related to time of exposure and concentration of agent used (exposure in non-ventilated space).
- Symptoms that may be experienced after exposure:
 - Eyes: tearing, pain, conjunctivitis, blurred vision
 - Nose, mouth, throat: rhinorrhea, burning or pain, trouble swallowing, drooling
 - Lungs: chest tightness, coughing, choking sensation, wheezing, dyspnea
 - Skin: burning, redness, dermatitis
 - GI: nausea and vomiting are rare and may be posttussive
- Symptoms begin within seconds of exposure, are self-limited and are best treated by removing patient from ongoing exposure. Symptoms frequently decrease over time (15–45 minutes) after exposure ends.

Pertinent Assessment Findings

- Riot control agent used
- Symptoms of exposed
- Lung sounds
- Evidence of other traumatic injuries

Hyperthermia and Heat Exposure

Aliases

Hyperthermia, heat cramps, heat exhaustion, heat syncope, heat edema, heat stroke

Definitions

1. *Heat cramps* are minor muscle cramps usually in the legs and abdominal wall. Patient temperature is normal.
2. *Heat exhaustion* has both salt and water depletion usually of a gradual onset. As it progresses tachycardia, hypotension, elevated temperature, and very painful cramps occur. Symptoms of headache, nausea and vomiting occur. Heat exhaustion can progress to heat stroke.
3. *Heat stroke* occurs when the cooling mechanism of the body (sweating) ceases due to temperature overload and/or electrolyte imbalances. Patient temperature is usually **greater than** 104°F. When no thermometer is available, it is distinguished from heat exhaustion by altered level of consciousness.
4. *Heat syncope* is a transient loss of consciousness with spontaneous return to normal mentation attributable to heat exposure.
5. *Heat edema* is dependent extremity swelling caused by interstitial fluid pooling.

Patient Care Goals

1. Administer cooling and rehydration.
2. Mitigate high risk for decompensation.
3. Mitigate high risk for agitation and uncooperative behavior.

Patient Presentation

Inclusion Criteria

1. Heat cramps
2. Heat exhaustion
3. Heat stroke
4. Heat syncope
5. Heat edema
6. Stimulant drug abuse
7. Excited delirium [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)]

Exclusion Criteria

1. Fever from infectious or inflammatory conditions
2. Malignant hyperthermia
3. Serotonin syndrome
4. Neuroleptic malignant syndrome

Patient Management

Assessment

1. Patient Assessment:
 - a. Age
 - b. Oral intake
 - c. Medications
 - d. Alcohol
 - e. Illicit drugs
 - f. Overdose
 - g. Withdrawal risk
2. Environmental Assessment:
 - a. Ambient temperature and humidity
 - b. Exertion level
 - c. Length of time at risk
 - d. Attire (clothing worn)
 - e. Confined space
 - i. **Pediatric Considerations:** Children left in cars who show signs of altered mental status and elevated body temperature should be presumed to have hyperthermia.
3. Associated Symptoms:
 - a. Cramps
 - b. Headache
 - c. Orthostatic symptoms
 - d. Nausea
 - e. Weakness
 - f. Mental status changes, including
 - i. Confusion.
 - ii. Coma.
 - iii. Seizures.
 - iv. Psychosis.
4. Vital signs:
 - a. Temperature: usually 104°F or greater (if thermometer available)
 - b. Skin:
 - i. Flushed and hot
 - ii. Dry or sweaty
 - iii. Signs of first or second degree burns from sun exposure
 - c. Other signs of poor perfusion or shock

Treatment and Interventions

1. Move victim to a cool area and shield from the sun or any external heat source.
2. Remove as much clothing as is practical and loosen any restrictive garments.
3. If alert and oriented, give small sips of cool liquids.
4. If altered mental status, check blood glucose level *[EMR-O; EMT-R]*.
5. Manage airway as indicated.
6. Place on ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]* and record ongoing vital signs and level of consciousness.
7. If temperature is greater than 104°F (40°C) or if altered mental status is present, begin active cooling:

- a. Immerse the patient in an ice bath; this provides the most rapid cooling mechanism but may not be available to EMS. If shivering occurs during cooling, administer Midazolam (preferred benzodiazepine) *[INT-O; PARA-R]*.
 - b. Mist the exposed skin continually with tepid water while fanning the victim (most effective).
 - c. Note that using truncal ice packs are acceptable, but are less effective than evaporation.
 - d. DO NOT apply wet clothes or wet clothing, as they may trap heat and prevent evaporative cooling.
8. Continue cooling efforts until the patient's temperature is less than 102.2°F (39°C) and the patient demonstrates improvement in mental status.
 9. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
 10. Monitor for arrhythmia and cardiovascular collapse [see [Cardiovascular](#) section guidelines].
 11. Treat seizures per the [Seizures guideline](#).
 12. Transport all patients suffering from life threatening heat illness (including heat stroke) to the hospital.

Patient Safety Considerations

Consider use of physical securing devices [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)] to protect vascular access sites.

Notes and Educational Pearls

Key Considerations

- Patients at risk for heat emergencies include neonates, infants, geriatric patients, and patients with mental illness.
- Contributory risk factors may come from:
 - Prescription and over-the-counter herbal supplements.
 - Cold medications.
 - Heart medications.
 - Diuretics.
 - Psychiatric medications.
 - Drug abuse.
 - Accidental or intentional drug overdose.
- Heat exposure can occur either due to increased environmental temperatures or prolonged exercise or a combination of both.

Note: Environments with temperature *greater than* 90°F and humidity *greater than* 60% present the most risk.
- Heat stroke is associated with cardiac arrhythmias independent of drug ingestion or overdose. Heat stroke has also been associated with cerebral edema.
- Do not forget to look for other causes of altered mental status such as low blood glucose level, or, in the proper circumstances (e.g. endurance exercise events), consider exercise associated hyponatremia (EAH), especially in the patient with altered mental status, normal blood glucose, and normal temperature.
- *Controversy:*
 - Shivering may occur while treating heat stroke.
 - It is uncertain how harmful shivering is to heat stroke patients.

- Cooling should be continued until the above temperature and mental status goals are met.
- Treat shivering as described above in 7a, under Treatment and Interventions.
- Research does not demonstrate the value of one benzodiazepine over another in shivering patients.
- Hyperthermia that is ***not*** the result of environmental factors has a differential that includes the following:
 - Fever and delirium
 - Hyperthyroid storm
 - Delirium tremens (DTs)
 - CNS lesion or tumor
 - Adverse drug event: neuroleptic malignant syndrome, malignant hyperthermia
 - Mental status changes without hyperthermia in the correct circumstances could be exercise associated hyponatremia
- There is no evidence supporting EMS utilizing orthostatic vital signs.

Pertinent Assessment Findings

1. Warning signs: fever, altered mental status
2. Blood glucose level for AMS

Hypothermia and Cold Exposure

Aliases

Hypothermia, frost bite, cold induced injuries

Patient Care Goals

1. Maintain hemodynamic stability.
2. Prevent further heat loss.
3. Rewarm the patient in a safe manner.
4. Manage hypothermia induced cardiac arrest appropriately.
5. Prevent loss of limbs.

Patient Presentation

1. Patients may suffer from hypothermia due to exposure to a cold environment (increased heat loss) or may suffer from a primary illness or injury that, in combination with cold exposure (heat loss in combination with decreased heat production), leads to hypothermia.
2. Patients may suffer systemic effects from cold (hypothermia) or localized effects (e.g. frostbite).
3. Patients with mild hypothermia will have normal mental status, shivering, and may have normal vital signs while patients with moderate to severe hypothermia will manifest mental status changes, eventual loss of shivering, and progressive bradycardia, hypotension, and decreased respiratory status.
4. Patients with frostbite will develop numbness involving the affected body part along with a "clumsy" feeling and areas of blanched skin. Later findings include a "woody" sensation, decreased or loss of sensation, bruising or blister formation, or a white and waxy appearance to affected tissue.

Inclusion Criteria

Patients suffering systemic or localized cold injuries

Exclusion Criteria

1. Patients without cold exposure
- OR**
2. Patients with cold exposure but no symptoms referable to hypothermia or frostbite

Patient Management

Assessment

1. Begin with the primary survey, looking for evidence of circulatory collapse and ensuring effective respirations.
 - a. The patient suffering from moderate or severe hypothermia may have severe alterations in vital signs including weak and extremely slow pulses, profound hypotension and decreased respirations.
 - b. The rescuer may need to evaluate the hypothermic patient for longer than the normothermic patient (up to 60 seconds).
2. Obtain standard SAMPLE-type patient history.
3. Obtain additional history, including:

- a. Any associated injury or illness.
- b. Duration of cold exposure.
- c. Ambient temperature.
- d. Any treatments initiated before EMS arrival.
4. Categorize the severity of hypothermia based on core body temperature readings and clinical evaluation.
 - a. Body temperature: Perform, if possible and reliable, core body temperature measurements and categorize patients into one of the four following levels of hypothermia:
 - Mild: normal body temperature 35–32.1°C (95–89.8°F)
 - Moderate: 32°–28°C (89.7°–82.5°F)
 - Severe: 28°–24°C (82.4°–75.2°F)
 - Profound: less than 24°C (75.2°F)
 - b. Balance the above temperature-based categorization against the following clinical findings, as these are equally important:
 - Mild: vital signs not depressed normal mental status, shivering is preserved; body maintains ability to control temperature
 - Moderate to severe: progressive bradycardia, hypotension, and decreased respirations, alterations in mental status with eventual coma, shivering will be lost in moderate hypothermia (generally between 31–30° C), and general slowing of bodily functions; the body loses ability to thermo-regulate

Treatment and Interventions

1. Maintain patient and rescuer safety—the patient has fallen victim to cold injury and rescuers have likely had to enter the same environment. Maintain rescuer safety by preventing cold injury to rescuers.
2. Manage airway per the [Airway Management guideline](#).
3. Initiate interventions according to severity of patient condition (a. Mild, b. Moderate or Severe, or b. Frostbite):
 - a. **Mild hypothermia:**
 - i. Remove the patient from the environment and prevent further heat loss by removing wet clothes and drying skin; insulate from the ground; shelter the patient from wind and wet conditions; and insulate the patient with dry clothing or a hypothermia wrap or blanket. Cover the patient with a vapor barrier and, if available, move the patient to a warm environment.
 - ii. Assess patient need for oxygen.
 - Hypothermic patients have decreased oxygen needs and may not require supplemental oxygen.
 - If oxygen is deemed necessary, it should be warmed, to a maximum temperature between 40–42°C (104–108°F) and humidified if possible.
 - iii. Fuel shivering through caloric replacement. Vigorous shivering can substantially increase heat production.
 - Provide beverages or foods containing glucose if feasible and patient is awake and able to manage airway independently.
 - iv. Consider field-rewarming methods such as placement of large heat

packs or heat blankets (chemical or electric if feasible) to the anterior chest or wrapped around the patient's thorax if large enough. Forced air warming blankets (e.g. Bair Hugger®) can be an effective field rewarming method if available.

- v. Monitor frequently—if temperature or level of consciousness decreases, refer to [Severe Hypothermia](#).
- vi. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]).
 - Indications for IV access and IV fluids in the mildly hypothermic patient are similar to those of the non-hypothermic patient.
 - IV fluids, if administered, should be warmed, ideally to 42°C.
 - Normal saline bolus therapy 20 mg/kg is preferable to continuous drip.
 - The recommended fluid for volume replacement in the hypothermic patient is normal saline.
- vii. If alterations in mental status, consider measuring blood glucose [*EMR-O*; *EMT-R*] and treat as indicated (treat per [Hypoglycemia](#) or [Hyperglycemia](#) guidelines) and assess for other causes of alterations of mentation.
- viii. Transport to a hospital capable of rewarming the patient.

b. Moderate or severe hypothermia:

- i. Perform ABCs—pulse checks for patients suffering hypothermia should be performed for 60 second—and obtain core temperature if possible for patients exhibiting signs or symptoms of moderate or severe hypothermia.
 - Core temperatures are best measured by esophageal probe [*EMR-O*; *PARAM-R*], if one is available, the patient's airway is secured, and the provider has been trained in its insertion and use.
 - If esophageal temperature monitoring is not available or appropriate, use an epitympanic thermometer designed for field conditions with an isolating ear cap.
 - Rectal temperatures may also be used, but only once the patient is in a warm environment. Rectal temperatures are not reliable or suitable for taking temperatures in the field and should only be done in a warm environment (such as a heated ambulance).
- ii. Manage airway as needed.
 - Care must be taken not to hyperventilate the patient as hypocarbia may reduce the threshold for ventricular fibrillation in the cold patient.
 - Indications and contraindications for advanced airway devices are similar in the hypothermic patient as in the normothermic patient.
- iii. Prevent further heat loss by removing the patient from the environment and removing wet clothes and drying skin, insulate from the ground, shelter the patient from wind and wet conditions, and insulate the patient with dry clothing or a hypothermia wrap or blanket. Cover the patient with a vapor barrier and, if available, move the patient to a warm

environment.

- iv. Initiate field-rewarming methods such as placement of large heat packs or heat blankets (chemical or electric if feasible) to the anterior chest or wrapped around the patient's thorax if large enough.
 - Never apply chemical or electrical heat sources directly to the skin.
 - Use a barrier between the skin and heat source to prevent burns.
 - Use forced air warming blankets (e.g. Bair Hugger®), if available; they can be an effective field rewarming method.
- v. Handle the patient gently.
 - Attempt to keep the patient in the horizontal position, especially limiting motion of the extremities to avoid increasing return of cold blood to the heart.
 - Once in a warm environment, clothing should be cut off (rather than removed by manipulating the extremities).
 - Move the patient only when necessary such as to remove the patient from the elements.
- vi. Apply ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*] or AED if available.
- vii. Establish IV and provide warmed NS bolus 20 mg/kg [*AEMT-R*] – Repeat as necessary.
- viii. If alterations in mental status, consider measuring blood glucose [*EMR-O; EMT-R*] and treat as indicated (treat per [Hypoglycemia](#) or [Hyperglycemia](#) guidelines) and assess for other causes of alterations of mentation.
- ix. Transport as soon as possible to a hospital capable of resuscitation. If cardiac arrest develops, consider transport to a center capable of extracorporeal circulation (ECMO) or cardiopulmonary bypass (if feasible).
- x. Warm the patient compartment of the ambulance to 24°C (75.2°F) during transport.

c. Frostbite:

- i. Avoid rewarming of extremities until definitive treatment is possible, if ambulation or travel is necessary for evacuation or safety. Additive injury occurs when the area of frostbite is rewarmed and then inadvertently refrozen. Only initiate rewarming if refreezing is absolutely preventable.
 - If rewarming is feasible and refreezing can be prevented, use circulating warm water (37–39°C or 98.6–102°F) to rewarm effected body part, and thaw injury completely. If warm water is not available, rewarm frostbitten parts by contact with non-affected body surfaces. Do not rub or cause physical trauma.
 - After rewarming, cover injured parts with loose, sterile dressing. If blisters are causing significant pain, and the provider is so trained, these may be aspirated, however, should not be de-roofed. Do not allow injury to refreeze. Treat per the [Pain Management guideline](#).

Patient Safety Considerations

1. Given the additive effects of additional cold stress, the patient should be removed from the cold environment as soon as operationally feasible.
2. In patients suffering from moderate to severe hypothermia, it is critical to not allow these patients to stand or exercise as this may cause circulatory collapse.
3. Devices that self-generate heat (e.g. heat packs) and are being utilized during the rewarming process should be wrapped in a barrier to avoid direct contact with the skin and to prevent burns. Available evidence suggests that heat packs with peak temperatures above 45°C (113°F) are most likely to cause burns. In patients who are unresponsive, or unable to recognize a developing injury, please check the area in which the heating pad is placed regularly to ensure no tissue damage occurs.

Notes and Educational Pearls

Key Considerations

Considerations in cardiac arrest associated with hypothermia.

- The following are contraindications for initiation of resuscitation in the hypothermic patient:
 - Obvious fatal injuries (such as decapitation)
 - The patient exhibits signs of being frozen (such as ice formation in the airway)
 - Chest wall rigidity such that compressions are impossible
 - Danger to rescuers or rescuer exhaustion
 - Avalanche victims buried for 35 minutes or longer with airway obstruction by ice or snow
- Recognize that fixed and dilated pupils, apparent rigor mortis, and dependent lividity may not be contraindication for resuscitation in the severely hypothermic patient.
- The mainstay of therapy in severe hypothermia and cardiac arrest should be effective chest compressions and attempts at rewarming.
 - Note:** Chest compressions should be provided at the same rate as in normothermic patients.
- The temperature at which defibrillation should first be attempted in the severely hypothermic cardiac arrest victim and the number of defibrillation attempts is unclear. There are different approaches regarding resuscitation of the hypothermic arrest patient.
 - Per the American Heart Association (AHA), if the patient has a shockable rhythm (VF/VT), defibrillation should be attempted. It is reasonable to continue defibrillation attempts per AHA protocols concurrently with rewarming strategies.
 - If defibrillation is unsuccessful and the patient's core temperature is greater than 30°C (86°F), follow guidelines for normothermic patients.
 - If available monitors reveal asystole, CPR alone is the mainstay of therapy.
 - If monitoring reveals an organized rhythm (other than VF or VT) and no pulses are detected, do not start CPR, but continue to monitor.
 - While this may represent pulseless electrical activity (PEA), this may also represent situations in which the patient's pulses are not detectable but remain effective due to decreased metabolic needs.
 - In the case of PEA, the rhythm will deteriorate rapidly to asystole, in which case, CPR should be initiated.
 - Given the potential to cause VF with chest compressions, the Alaska guidance offers that it is better to maintain effective cardiac activity than to start CPR and cause VF.

- Manage the airway per standard care in cardiac arrest victims [see [Cardiac Arrest guideline](#)].
 - In the absence of advanced airways, ventilate the patient at the same rate as a normothermic patient.
 - If the patient has an advanced airway, ventilate at half the rate recommended for a normothermic patient to prevent hyperventilation. If ETCO₂ is available, ventilate to maintain normal ETCO₂ levels.
 - There is little evidence to guide use of medications in severe hypothermia with cardiac arrest, however 2010 AHA updates to advanced cardiac life support recommend use of vasopressors according to standard ACLS protocols.
 - Above 30°C (86°F), intervals between medication provision should be doubled until the patient reaches 35°C (95°F), at which time, normal medication intervals may be adopted.
- Upon ROSC, treat per [Adult Post-ROSC guideline](#).
- Patients with severe hypothermia and arrest may benefit from resuscitation even after prolonged downtime, and survival with intact neurologic function has been observed even after prolonged resuscitation.

Note: Patients should not be considered deceased until rewarming has been attempted.
- If a hypothermic patient clearly suffered cardiac arrest and subsequently became hypothermic afterward with prolonged down time between arrest and rescue, there is no rationale for initiating resuscitation and warming the patient.

Pertinent Assessment Findings

1. Identification of associated traumatic injuries (when present)
2. Identification of localized freezing injuries
3. Patient core temperature (when available)

Drowning

Aliases

Near-drowning, non-fatal drowning, fatal drowning, submersion, immersion

Patient Care Goals

1. Rapid assessment and management of life-threatening injuries
2. Rescue from the water-based environment
3. Transport all patients suffering from drowning for hospital evaluation

Patient Presentation

Inclusion Criteria

Patients suffering from drowning or drowning events independent of presence or absence of symptoms

Exclusion Criteria

Patients without history of drowning

Patient Management

Assessment

1. Follow [Universal Care guideline](#).
2. Obtain patient history: It should include circumstances leading to the submersion, details of mechanism of injury, time under water, and water temperature (if available).
Note: Consider possible c-spine injury when taking history, mechanism of injury, and conducting exam. If evaluation suggests injury to the cervical spine, manage c-spine [*EMR-O; EMT-R*].
3. Conduct primary survey: It should include aggressive airway management and restoration of adequate oxygenation and ventilation. Unlike the CAB strategy used in standard cardiac arrest, patients suffering cardiac arrest from drowning require an ABC approach with prompt airway management and supplemental breathing.
4. Assess for other associated injury such as injury to the head or dive-related emergency.

Treatment and Interventions

1. Ensure scene safety for patient and rescuers. Remove patient from water as soon as possible by rescuers who are properly trained and equipped.
 - a. Practice the safest water rescue technique possible, given circumstances on scene.
 - b. Evacuate to land or a water craft as soon as possible.
 - c. If there is a delay to accessing shore or a rescue boat, initiate in-water basic life support consisting of ventilation only.
2. Manage airway per the [Airway Management guideline](#).
3. Follow [Cardiac Arrest guideline](#) as indicated with consideration of ABC strategy for drowning victims in cardiac arrest.
 - a. Initiate 5 rescue breaths followed by 30 chest compressions.
 - b. After the initial 5 breaths, use a 2 breaths to 30 compression ratio.
4. If mechanism or history suggest cervical spine injury, manage c-spine per the [Spinal Care guideline](#).

5. Monitor vital signs including oxygen saturations *[EMR-O; EMT-R]*.
6. Administer oxygen *[EMR-O; EMT-R]* as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
7. Consider positive pressure ventilation in patients with signs or symptoms of respiratory difficulty.
8. Consider hypothermia, treat per [Hypothermia/Cold Exposure guideline](#).
9. If the victim was involved in underwater diving and uncertainty exists regarding the most appropriate therapy, consider contacting on-line medical control and discussing need for hyperbaric treatment. Include discussion regarding:
 - a. Submersion time.
 - b. Greatest depth achieved.
 - c. Ascent rate.
 - d. Gas mix.
10. Establish IV access *[AEMT-R]*.
11. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline *[AEMT-R]* or lactated Ringer's *[AEMT-O]*).
12. Perform advanced airway management as indicated. Consider non-invasive positive pressure ventilation *[EMT-O; AEMT-R]* in awake patients with respiratory distress.
13. Apply ECG cardiac monitor *[Acquisition EMT-O; Interpretation INT-R]*.

Patient Safety Considerations

1. Avoidance of hyperoxygenation of the drowning victim
2. Consideration of rescuer safety

Notes and Educational Pearls

Key Considerations

- The World Health Organization definition of drowning is "the process of experiencing respiratory impairment from submersion or immersion in liquid."
- Drowning is further defined in the following categories:
 - Non-fatal drowning: patients rescued from drowning
 - Fatal drowning: any death, acutely or subacutely, resultant from drowning
- Submersion refers to situations in which the patient's airway is underwater.
- Immersion refers to situations in which the patient's body is in water but the patient's airway remains out of the water.
- **Pediatric Considerations:**
- Drowning is a common cause of death in children.
- Risk factors for drowning include male gender, age less than 14 yo, alcohol use, lack of supervision, and risky behavior.
- Rescue efforts should be coordinated between all responding agencies to ensure patient is rapidly accessed and removed from the water.
- Initiation of in-water ventilations may increase survival. In-water chest compressions are futile.
- The European Resuscitation Council recommends 5 initial breaths be provided to the drowning victim.
- The initial ventilations may be more difficult to achieve as water in the airways may impede alveolar expansion.
- After the initial 5 breaths and 30 compressions, the standard ratio of 2 breaths to 30

compressions may be resumed.

- Active efforts to expel water from the airway (by abdominal thrusts or other means) should be avoided as they delay resuscitative efforts and increase the potential for vomiting and aspiration.
- Long-standing teaching has suggested that rescuers should always assume c-spine injury in victims of drowning.
 - The 2010 American Heart Association update on special circumstances in cardiac arrest notes that routine c-spine precautions in all victims of drowning is likely unnecessary unless the mechanism or injury, history, or physical exam suggests a cervical spine injury.
 - Mechanisms of injury highly suggestive of cervical spine injury include diving, water skiing, surfing or watercraft accidents.
- Uncertainty exists regarding survival in cold water drowning, however, recent literature suggests the following:
 - If water temperature is less than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest:
 - Survival is possible for submersion time less than 90 minutes and resuscitative efforts should be initiated.
 - Survival is not likely for submersion time greater than 90 minutes and providers may consider not initiating resuscitation or termination of resuscitation on scene.
 - If water temperature is greater than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest:
 - Survival is possible for submersion time less than 30 minutes and resuscitative efforts should be initiated.
 - Survival is not likely for submersion time greater than 30 minutes and providers may consider not initiating resuscitation or termination of resuscitation on scene.
- Patients may develop subacute respiratory difficulty after drowning and therefore all victims of drowning should be transported for observation.

Dive (SCUBA) Injury or Accidents

Aliases

Barotrauma, bends, squeeze

Patient Care Goals

1. Rapid assessment and management of life-threatening injuries
2. Rescue from the water-based environment
3. Transport of patients suffering from self-contained underwater breathing apparatus (SCUBA) diving injury or illness for hospital evaluation and consideration of repressurization or hyperbaric oxygen therapy (HBOT)

Patient Presentation

Inclusion Criteria

Patients with history of recent (within 48 hours) SCUBA diving activity who are exhibiting potential signs and/or symptoms of dive related illness or injury, regardless of dive table compliance

Note: SCUBA-related complications may occur anywhere, particularly when divers travel by air within 24-hours of diving.

Exclusion Criteria

Patients without history of recent (within 48 hours) SCUBA diving exposure

Patient Management

Assessment

1. Follow [Universal Care guideline](#).
2. Obtain patient history: It should include circumstances leading to the complaint; details of mechanism of injury; time under water; depth of dive; compliance with dive tables and decompression stops, gas mixture used, and water temperature (if available).
3. Be alert for signs of barotrauma (pulmonary barotrauma; arterial gas embolism; pneumothorax; ear, sinus, or dental barotrauma etc.); and/or decompression sickness (joint pain, mental status change, other neurologic symptoms including paralysis) or nitrogen narcosis (confusion, intoxication).
4. Assess for other associated injury such as injury to the head or spine (if mechanism and symptoms suggest); marine envenomation; hypothermia; or other injury.

Treatment and Interventions

1. Consider instructions for drowning or near-drowning, if a SCUBA accident includes this context [see [Drowning guideline](#)].
2. Manage airway as indicated.
3. If air embolism suspected, place in left lateral recumbent position (patient lying with the left side down, knees drawn upward, and flat).
 - a. Trendelenburg position is sometimes recommended to help trap the air in the dependent right ventricle, and may be useful if a central venous catheter is being used to withdraw the air, but this position may increase cerebral edema.

4. Monitor vital signs including oxygen saturations [*EMR-O; EMT-R*] and cardiac rhythm (if possible).
 - a. Administer oxygen [*EMR-O; EMT-R*] as appropriate for dyspnea or distress with a target of achieving greater than 93% saturation for most acutely ill patients.
 - b. Use positive pressure ventilation carefully in patients for whom pulmonary barotrauma is a consideration [see [Airway Management guideline](#)].
5. Place patients with symptoms suggesting decompression illness on supplemental oxygen, regardless of saturations, to enhance washout of inert gases.
6. Assess for hypothermia, treat per [Hypothermia/Cold Exposure guideline](#).
7. Consider contacting on-line medical control and discussing need for hyperbaric treatment and primary transport to facility with HBOT capability. Include discussion regarding factors such as submersion time, greatest depth achieved, ascent rate, and gas mix.
8. Establish IV access [*AEMT-R*].
9. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated Ringer's [*AEMT-O*]).

Patient Safety Considerations

1. If the patient is still in the water, seek safest and most rapid means of removal (within your scope of training) while minimizing risk of further injury.
2. Seek assistance early for special rescue, extrication, and transportation needs.
3. Check for multiple patients (e.g. group dive table calculation error(s) or contaminated dive gases).

Notes and Educational Pearls

Key Considerations

- Rescue efforts should be coordinated between all responding agencies to ensure that the patient is rapidly accessed and safely removed from the water if the diver is unable to do so themselves.
- If air medical transport is necessary, the patient should be transported with the cabin pressurized to lowest possible altitude. If an unpressurized aircraft is used (e.g. most helicopter [HEMS] services), patient should be flown at the lowest safe altitude possible.
- Decompression illness may have a variety of presentations depending on system affected (e.g. skin, joint(s), pulmonary, neurologic),
- SCUBA accidents or incidents can result in a variety of issues, including barotrauma, air embolism and decompression illness.

Pertinent Assessment Findings

- Vital signs findings
- Neurologic status assessment findings
- Respiratory assessment findings (e.g. oxygen saturation, respiratory rate)
- Subcutaneous emphysema findings

Conducted Electrical Weapon Injury (e.g. TASER®)

Aliases

Tased

Patient Care Goals

1. Manage the condition that triggered the application of the conducted electrical weapon; focus special attention on patients meeting criterion for excited delirium [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].
2. Make sure patient is appropriately secured or restrained with assistance of law enforcement to protect the patient and staff [see [Agitated or Violent Patient/Behavioral Emergency guideline](#)].
3. Perform comprehensive trauma and medical assessment as patients who have received conducted electrical weapon may have already been involved in physical confrontation.
4. Assess distance of discharge: If discharged from a distance, locate and remove two single barbed darts (13mm length) from the patient.
Note: Do not remove barbed dart from sensitive areas (head, neck, hands, feet or genitals).

Patient Presentation

Inclusion Criteria

1. Patient received either the direct contact discharge or the distance two barbed dart discharge of the conducted electrical weapon
2. Patient may have sustained fall or physical confrontation trauma
3. Patient may be under the influence of toxic substances and or may have underlying medical or psychiatric disorder

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Secure or restrain patient with assistance of law enforcement.
2. Perform primary and secondary assessment including 3-lead ECG and pulse oximeter. Consider using 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*].
3. Evaluate patient for evidence of excited delirium manifested by varied combination of agitation, reduced pain sensitivity, elevated temperature, persistent struggling, or hallucinosis.

Treatment and Interventions

1. Make sure patient is appropriately secured with assistance of law enforcement to protect the patient and staff. Consider psychologic management medications if patient is struggling against physical devices and may harm themselves or others.
2. Remove barbed darts except for sensitive areas (head, neck, hands, feet, or genitals).
3. Treat medical and traumatic injury.

Patient Safety Considerations

- Before removal of the barbed dart, make sure the cartridge has been removed from the conducted electrical weapon.
- Patient should not be restrained in the prone, face down, or hog-tied position as respiratory compromise is a significant risk.
- The patient may have underlying pathology before being tased (refer to appropriate guidelines for managing the underlying medical or traumatic pathology).
- Perform a comprehensive assessment with special attention looking for signs and symptoms that may indicate agitated delirium.
- Transport the patient to the hospital if they have concerning signs or symptoms.
- EMS providers who respond for a conducted electrical weapon patient should not perform a “medical clearance” for law enforcement.

Notes and Educational Pearls

Key Considerations

- Conducted electrical weapon can be discharged in three fashions:
 - By direct application of weapon without the use of the darts
 - By a single dart combined with direct application of weapon
 - By two darts fired from a distance up to 35 feet
- The device delivers 19 pulses per second with an average current per pulse of 2.1 milliamps which—in combination with toxins or drugs, patient’s underlying diseases, excessive physical exertion, and trauma—may precipitate arrhythmias. Consider ECG monitoring and 12-lead ECG assessment.
- Drive Stun is a direct weapon two-point contact which is designed to generate pain and not incapacitate the subject. Only local muscle groups are stimulated with the Drive Stun technique.

Pertinent Assessment Findings

1. Thoroughly assess the tased patient for trauma as the patient may have fallen from standing or higher.
2. Ascertain if more than one TASER® cartridge was used (by one or more officers, in effort to identify total number of possible darts and contacts).

Electrical Injuries

Aliases

Electrical burns, electrocution

Patient Care Goals

1. Prevent additional harm to patient.
2. Identify life threatening issues such as dysrhythmias and cardiac arrest.
3. Identify characteristics of electrical source to communicate to receiving facility (voltage, amperage, alternating current [AC] versus direct current [DC]).
4. Understand that deep tissue injury can be far greater than external appearance.
5. Have high index of suspicion for associated trauma due to patient being thrown.
6. Determine most appropriate disposition for the patient as many will require burn center care and some may require trauma center care.

Patient Presentation

Inclusion Criteria

Exposure to electrical current (AC or DC).

Exclusion Criteria

None

Patient Management

Assessment

1. Verify scene is secure. The electrical source must be disabled prior to assessment by rescuers who are properly trained and equipped.
2. Conduct primary survey with specific focus on dysrhythmias or cardiac arrest. Apply a ECG cardiac monitor [*Acquisition EMT-O; Interpretation INT-R*].
3. Identify all sites of burn injury. If the patient became part of the circuit, there will be an additional site near the contact with ground; electrical burns are often full thickness and involve significant deep tissue damage.
4. Assess for potential associated trauma and note if the patient was thrown from contact point. If patient has altered mental status, assume trauma was involved and treat accordingly.
5. Assess for potential compartment syndrome from significant extremity tissue damage.
6. Determine characteristics of source, if possible: AC or DC, voltage, amperage, and also time of injury.

Treatment and Interventions

1. Identify dysrhythmias or cardiac arrest—even patients who appear dead (particularly dilated pupils) may have good outcomes with prompt intervention [see appropriate guideline for additional information and patient assessment/treatment].
2. Immobilize if associated trauma suspected [see [Trauma](#) section guidelines].
3. Apply dry dressing to any wounds.
4. Remove constricting clothing and jewelry since additional swelling is possible.
5. Consider isotonic IV/IO fluid bolus 20 ml/kg (normal saline [*AEMT-R*] or lactated

Ringer's *[AEMT-O]*).

6. Remember that external appearance will underestimate the degree of tissue injury.
7. Treat pain per [Pain Management guideline](#); electrical injuries may be associated with significant pain.
8. Take electrical injury patients to a burn center whenever possible, since these injuries can involve considerable tissue damage.
9. Prioritize treatment of trauma when there is significant associated trauma and if local trauma resources and burn resources are not in the same facility.

Patient Safety Considerations

1. Verify no additional threat to patient
2. Shut off electrical power
3. Move patient to shelter if electrical storm activity still in area

Notes and Educational Pearls

Key Considerations

- Electrical current causes injury through three main mechanisms:
 - Direct tissue damage, altering cell membrane resting potential, and eliciting tetany in skeletal and/or cardiac muscles
 - Conversion of electrical energy into thermal energy, causing massive tissue destruction and coagulative necrosis
 - Mechanical injury with direct trauma resulting from falls or violent muscle contraction
- Anticipate atrial and/or ventricular dysrhythmias as well as cardiac arrest.
- The mortality related to electrical injuries is impacted by the following:
 - Route current takes through the body (current traversing the heart has higher mortality)
 - Type of current—AC vs. DC
 - AC is more likely to cause cardiac dysrhythmias while DC is more likely to cause deep tissue burns; however, either type of current can cause any injury.
 - DC typically causes one muscle contraction while AC can cause repeated contractions.
 - Both types of current can cause involuntary muscle contractions that do not allow the victim to let go of the electrical source
 - AC is more likely to cause ventricular fibrillation while DC is more likely to cause asystole
 - The amount of current impacts mortality more than the voltage

Current level (Milliamperes)	Probable Effect on Human Body of 120 V, 60 Hz AC for 1 second
1 mA	Perception level; slight tingling sensation; still dangerous if wet conditions
5mA	Slight shock felt; not painful but disturbing; average individual can let go; however, strong involuntary reactions to shocks in this range may lead to injuries
6mA–16mA	Painful shock; begin to lose muscular control; commonly referred to as the freezing current or "let-go" range
17mA–99mA	Extreme pain, respiratory arrest, severe muscular contractions; individual cannot let go; death is possible
100mA–2000mA	Ventricular fibrillation (uneven, uncoordinated pumping of the heart); muscular contraction and nerve damage begins to occur; death is likely
> 2,000mA	Cardiac arrest, internal organ damage, and severe burns; death is probable

Source: https://www.osha.gov/SLTC/etools/construction/electrical_incidents/eleccurrent.html

Pertinent Assessment Findings

1. Identification of potential trauma concomitant with electrical injury
2. Presence of cardiac dysrhythmias

Lightning or Lightning Strike Injury

Aliases

Lightning burn

Patient Care Goals

1. Identify patient(s) as lightning strike victim(s).
2. Move to safe area.
3. Initiate immediate resuscitation of cardiac arrest victim(s), within limits of mass casualty care, also known as "reverse triage."
4. Perform ECG cardiac monitoring during transport.
5. Treat associated traumatic injuries.

Patient Presentation

1. Environment: Lightning strikes may happen in a variety of environmental conditions. Most commonly they occur in outdoor or wilderness circumstances:
Golf courses, exposed mountains or ledges and farms or fields all present conditions that increase risk of lightning strike when hazardous meteorological conditions exist.
2. Injury: Lacking bystander observations or history, it is not always immediately apparent that the patient has been the victim of a lightning strike.
Subtle findings such as injury patterns might suggest lightning injury.

Inclusion Criteria

Patients of all ages who have been the victim of lightning strike injury

Exclusion Criteria

No recommendations

Patient Management

Assessment

1. Respiratory
 - a. Apnea
 - b. Agonal respirations
 - c. Respiratory paralysis
2. Cardiovascular
 - a. Dysrhythmias
 - b. Transient hypertension
3. Neurologic
 - a. Seizures
 - b. Confusion
 - c. Paralysis
 - d. Paraplegia
 - e. Vertigo or dizziness
 - f. Parasthesias
 - g. Amnesia
 - h. Memory deficits

- i. Anxiety
- j. Fixed or dilated pupils possible (autonomic dysfunction)
- 4. Skin
 - a. Ferning or fern-like superficial skin burn ("Lichtenberg figures")
 - b. Vascular instability may result in cool, mottled extremities
 - c. Numerous first and/or second degree burns
 - d. Third degree burns (less common)
- 5. Cardiopulmonary, as injury is a result of DC current:
 - a. Patient may be in full cardiopulmonary arrest
 - b. Patient may have only respiratory arrest
- 6. Neurologic
 - a. May have stroke-like findings as a result of neurologic insult
- 7. Other
 - a. May have secondary traumatic injury as a result of overpressurization, blast or missile injury
 - b. Fixed/dilated pupils may be a sign of neurologic insult, rather than a sign of death or impending death. Do not use as a solitary, independent sign of death for the purpose of discontinuing resuscitation in this patient population.

Treatment and Interventions

1. Assure patent airway. If in respiratory arrest only, manage airway as appropriate.
2. If in cardiopulmonary arrest, treat per [Cardiac Arrest guideline](#)
3. Consider IV initiation [*AEMT-R*]. Avoid initiation through burned skin.
4. Monitor ECG. Be alert for potential arrhythmias. Consider 12-lead ECG [*Acquisition EMT-O; Interpretation INT-R*], when available.
5. Consider early pain management for burns or associated traumatic injury [see [Pain Management guideline](#)].

Patient Safety Considerations

1. Recognize that repeat strike is a risk. Patient and rescuer safety are paramount.
2. Recognize that victims do not carry or discharge a current, so the patient is safe to touch and treat.

Notes and Educational Pearls

Key Considerations

- Lightning strike cardiopulmonary arrest patients have a high rate of successful resuscitation, if initiated early, in contrast to general cardiac arrest statistics.
- There may be multiple victims.
- If multiple victims, cardiac arrest patients whose injury was witnessed or thought to be recent should be treated first and aggressively (reverse from traditional triage practices).
 - Patients suffering cardiac arrest from lightning strike initially suffer a combined cardiac and respiratory arrest.
 - Return of spontaneous circulation may precede resolution of respiratory arrest.
 - Patients may be successfully resuscitated if provided proper cardiac and respiratory support, highlighting the value of "reverse triage."
- It may not be immediately apparent that the patient is a lightning strike victim.
- Injury pattern and secondary physical exam findings may be key in identifying patient as a victim of lightning strike.

- Lightning strike is a result of very high voltage, very short duration DC current exposure.

Pertinent Assessment Findings

1. Presence of thermal or non-thermal burns
2. Evidence of trauma
3. Evidence of focal neurologic deficits

APPENDICES

I. Universal Documentation Guideline

Aliases

NEMSIS, Documentation

Patient Care Goals

1. Support continuity of patient care and continuous performance improvement (CPI) of patient care through meeting minimum documentation standards for all EMS events where a patient was encountered.
2. This guideline defines minimum standards and inclusions used and referenced throughout this document under the "Quality Improvement" section of each guideline.
3. The National EMS Information System (NEMSIS) submission requirements, state and local EMS systems, and EMS billing reimbursement services will have more extensive minimum requirements that exceed this guideline.
4. Use this guideline as a starting point for systems looking to more formally define documentation requirements.

Patient Presentation

Inclusion Criteria

All EMS events where a patient was encountered and one or more clinical guideline was used to determine patient treatment and/or disposition

Exclusion Criteria

None

Toolkit for Key Categories of Data

Elements Incident

Demographics

1. Incident Demographics include the type of incident, location, time, dispatch information, response resources and patient and incident disposition of the EMS event.
 - a. This information will always apply and be available, even if the responding unit never arrives on scene (is cancelled) or never makes patient contact.
 - b. Incident demographics are important for filtering incident types and outcomes when doing CPI reviews, providing aggregate descriptive data, and billing for reimbursement.
2. Minimum Incident Demographic Fields include:
 - a. Incident Times
 - i. eTimes.03: Unit Notified by Dispatch Date/Time (*NEMSIS mandatory*)
 - ii. eTimes.05: Unit En Route Date/Time (*Unit responding*)
 - iii. eTimes.06: Unit Arrived on Scene Date/Time (*If arrived*)
 - iv. eTimes.07: Arrived at Patient Date/Time (*If patient contact made*)
 - v. eTimes.09: Unit Left Scene Date/Time (*Unit Transporting Time, if applicable*)

- vi. eTimes.11: Patient Arrived at Destination Date/Time (*If applicable*)
 - vii. eTimes.13: Unit Back in Service Date/Time (*NEMSIS mandatory*)
 - b. eResponse.05: Type of Service Requested (*e.g. 911 vs interfacility*)
 - c. eResponse.07: Primary Role of the Unit (*e.g. Transport or non-transport*)
 - d. eDispatch.01: Complaint Reported by Dispatch (*Dispatch reason from EMD*)
 - e. Crew Responding:
 - i. eCrew.01: Crew Member ID (*Crew name or license # depending on software*)
 - ii. eCrew.02: Crew Member Level (*License level for this call*)
 - iii. eCrew.03: Crew Member Response Role (*e.g. Primary or secondary care giver*)
 - f. eScene.09: Incident Location Type
 - i. Used for multiple purposes, including CARES (Cardiac Arrest Registry to Enhance Survival)
 - g. Response Modes (*e.g. lights and sirens*)
 - i. eResponse.23: Response Mode to Scene
 - ii. eResponse.24: Additional Response Mode Descriptors
 - h. Delays:
 - i. eResponse.09: Type of Response Delay
 - ii. eResponse.10: Type of Scene Delay

Patient Demographics and Medical History

Patient demographics in this section include the minimum information required for CPI review and do not include protected health information (PHI) or patient identifiable information. Local systems may require additional PHI to support EMS reimbursement and link local level CPI reviews to specific incidents or outcome data.

1. Minimum Patient Demographic and History Fields include:
 - a. ePatient.13 - Gender
 - b. ePatient.15 - Age
 - c. ePatient.16 - Age Units
 - d. eHistory.06 - Medication Allergies
 - e. eHistory.07 - Environmental/Food Allergies
 - f. eHistory.08 - Medical/Surgical History
 - g. eHistory.12 - Current Medications
 - h. eHistory.17 - Alcohol/Drug Use Indicators
 - i. eHistory.01 - Barriers to Patient Care
 - j. eExam.01 - Estimated Body Weight in Kilograms
 - k. eExam.02 - Length-Based Tape Measure

Patient Complaints and Symptoms

1. Patient and situational history for this EMS event generally addresses issues leading up to EMS being requested and include patient complaints; SAMPLE history; signs or symptoms; barriers and confounders; onset times; and trauma and cardiac arrest historical information.
2. Patient Complaints, Signs and Symptoms, and Key Related Times:
 - a. eSituation.02 - Possible Injury
 - b. Patient Complaint Group
 - i. eSituation.03 - Complaint Type
 - ii. eSituation.04 - Complaint
 - iii. eSituation.05 - Duration of Complaint
 - iv. eSituation.06 - Time Units of Duration of Complaint

- c. eSituation.07 - Chief Complaint Anatomic Location
- d. eSituation.08 - Chief Complaint Organ System
- e. Signs and Symptoms
 - i. eSituation.01 - Date/Time of Symptom Onset
 - ii. eSituation.09 - Primary Symptom [Single Choice]
 - iii. eSituation.10 - Other Associated Symptoms [Choose All that Apply]
- f. eSituation.18 - Date/Time Last Known Well (Stroke/CVA)

Situational History for this EMS Event

3. SAMPLE History

Note: Although many assessment guidelines refer to this history mnemonic, many electronic patient care report (ePCR) systems do not collect this information in a tool organized specifically in this group, but rather throughout the EMS record in the appropriate areas to the topics.

a. Symptoms

- i. eSituation.09 - Primary Symptom
- AND**
- ii. eSituation.10 - Other Associated Symptoms

b. Allergies

- i. eHistory.06 - Medication Allergies
- AND**
- ii. eHistory.07 - Environmental/Food Allergies

a. Medications

- i. eHistory.12 - Current Medications

b. Past medical and surgical history

- i. eHistory.08 - Medical/Surgical History

c. Last Oral Intake

- i. eHistory.19 - Last Oral Intake (*if software configured to collect*)
and/or
- ii. eNarrative.01 - Patient Care Report Narrative

d. Events leading to activation of EMS

- i. eSituation.17 - Patient Activity and/or
- ii. eNarrative.01 - Patient Care Report Narrative

4. Barriers and Situational Confounders

- a. eHistory.01 - Barriers to Patient Care
- b. eHistory.17 - Alcohol/Drug Use Indicators

5. Stroke

- a. eSituation.18 - Date/Time Last Known Well (Stroke/CVA)

6. Trauma History and Situation

- a. eSituation.02 - Possible Injury (*Yes/No - based on mechanism, not listing an actual injury*)
- b. eInjury.01 - Cause of Injury
 - i. Known to providers as *Mechanism of Injury*; values are from ICD-10
 - ii. Intent is included where possible in ICD-10, but is no longer a separate field as it was in NEMSIS v2
- c. eInjury.03 - Trauma Center Criteria (*Combined steps 1 and 2 of CDC's "Guidelines for Field Triage of Injured Patients"*)
- d. eInjury.04 - Vehicular, Pedestrian, or Other Injury Risk Factor (*Combined steps 3 and 4 of CDC's "Guidelines for Field Triage of Injured Patients"*)

- e. eInjury.07 - Use of Occupant Safety Equipment
 - f. Destination Pre-Arrival Alerts (e.g. trauma alerts)
 - i. eDisposition.24 - Destination Team Pre-Arrival Alert or Activation
 - ii. eDisposition.25 - Date/Time of Destination Pre-Arrival Alert or Activation
7. Cardiac Arrest History and Situation
- Note:** The following fields meet the needs of Utstein Criteria reports and many of the fields in CARES. CARES has additional custom fields that may be available from your software vendor.
- a. eArrest.01 - Cardiac Arrest [Yes/No]
 - b. eArrest.02 - Cardiac Arrest Etiology
 - c. eArrest.03 - Resuscitation Attempted By EMS
 - d. eArrest.04 - Arrest Witnessed By
 - e. eArrest.05 - CPR Care Provided Prior to EMS Arrival
 - f. eArrest.06 - Who Provided CPR Prior to EMS Arrival
 - g. eArrest.07 - AED Use Prior to EMS Arrival
 - h. eArrest.08 - Who Used AED Prior to EMS Arrival
 - i. eArrest.09 - Type of CPR Provided
 - j. eArrest.11 - First Monitored Arrest Rhythm of the Patient
 - k. eArrest.12 - Any Return of Spontaneous Circulation
 - l. eArrest.14 - Date/Time of Cardiac Arrest
 - m. eArrest.15 - Date/Time Resuscitation Discontinued
 - n. eArrest.16 - Reason CPR/Resuscitation Discontinued
 - o. eArrest.17 - Cardiac Rhythm on Arrival at Destination
 - p. eArrest.18 - End of EMS Cardiac Arrest Event
 - q. eScene.02 - Other EMS or Public Safety Agencies at Scene
 - r. eScene.03 - Other EMS or Public Safety Agency ID Number
 - s. eScene.04 - Type of Other Service at Scene

Provider Impressions and Incident or Patient Disposition

- 1. Provider Impressions (Provider Field Working Diagnosis)
 - a. eSituation.11 - Provider's Primary Impression [Single Choice]
 - i. The word "Primary" causes a great deal of understandable confusion with this field; this should be the diagnosis of the most acute (primary) problem ***not necessarily the first*** problem that was wrong with the patient, or their initial complaint.
 - b. eSituation.12 - Provider's Secondary Impressions [Choose all that Apply]
- 2. Incident/Patient Disposition
 - a. eSituation.13 - Initial Patient Acuity (*Intended to be prior to EMS care*)
 - b. eDisposition.19 - Final Patient Acuity (*Intended to be after EMS care*)
 - c. eDisposition.12 - Incident/Patient Disposition
 - d. eDisposition.16 - EMS Transport Method
 - e. Transport Mode (*e.g. use of lights and sirens*)
 - i. eDisposition.17 - Transport Mode from Scene
 - ii. eDisposition.18 - Additional Transport Mode Descriptors
 - f. eDisposition.01 - Destination/Transferred To, Name
 - i. Intended by NEMSIS to be the destination facility or the Agency transferred to, although many ePCR systems only collect this as the destination facility because of the complexity of mixing facilities and services in the same field.

Assessments and Exams

1. Exams

By definition, use of NEMSIS eExam fields is optional; they are, however, available for both state and local EMS system use.

 - a. Many systems do not require use of these fields as they can be time-consuming to enter, often too detailed (e.g. there is no value for whole arm, it would need to be entered as shoulder, upper arm, elbow, forearm and wrist with separate exam findings for each component, meaning a single exam finding of paralysis for an arm would take ten steps to enter) and the same information is often reflected in the provider's narrative.
 - b. However, there *is* some utility in targeted use of these fields for certain situations such as stroke, spinal exams, and trauma without needing to enter all the fields in each record.
2. Capacity Assessment Group

This can be used to support documentation of patient capacity for refusal of care and/or transport, participation in advanced spinal assessments, or support for treatment decisions by EMS providers.

Note: The Capacity Assessment Group does not provide a legal definition of capacity and should not be used as such. It is intended only to assist the EMS provider in documenting the most basic exam and history findings in order to determine capacity. Many additional factors must be considered when determining capacity, including the situation, patient medical history, medical conditions, and consultation with on-line medical control.

 - a. Barriers and situational confounders [Both only single entry]
 - i. eHistory.01 - Barriers to Patient Care
 - ii. eHistory.17 - Alcohol/Drug Use Indicators
 - b. Glasgow Coma Score (GCS) Vitals Group [*see Vitals section*] [serial entries allowed]
 - c. eVitals.26 - Level of Responsiveness (AVPU) [serial entries allowed]
 - d. eExam.19 - Mental Status Assessment [serial entries allowed]
 - e. eExam.20 - Neurological Assessment [serial entries allowed]
3. Stroke Assessments
 - a. Initial Vitals
 - b. eSituation.18 - Date/Time Last Known Well (Stroke/CVA)
 - c. Stroke Score Group
 - d. eExam.19 - Mental Status Assessment
 - e. eExam.20 - Neurological Assessment (*Speech, facial droop, arm drift, unilateral weakness*)
 - f. eVitals.31 - Reperfusion Checklist (*May not apply if service area does not use due to lack of consensus on a standard reperfusion checklist, or acceptance by EMS if used*)
4. Spinal Injury/Exam
 - a. Capacity Assessment Group
 - b. Back and Spine Assessment Group
 - i. eExam.13 - Back and Spine Assessment Finding Location
 - ii. eExam.14 - Back and Spine Assessment
 - c. Extremity Assessment Group
 - i. eExam.15 - Extremity Assessment Finding Location
 - ii. eExam.16 - Extremities Assessment
5. 12-lead ECG Acquisition
 - a. eTimes.06 - Unit Arrived on Scene Date/Time

- b. eTimes.07 - Arrived at Patient Date/Time
 - c. ECG Rhythm Group [see **Vitals** section]
 - d. Attach 12-lead graphic ePCR (through direct integration linkage with ECG monitor or attachment of scanned printout as allowed/available in software)
 - e. 12-lead-ECG Procedure-documented under Procedures Performed Group
6. Trauma/Injury
- The exam fields have many useful values for documenting trauma (deformity, bleeding, burns, etc.). Use of targeted documentation of injured areas can be helpful, particularly in cases of more serious trauma. Because of the endless possible variations where this could be used, specific fields will not be defined here.

Note: Exam fields use a specific and useful Pertinent Negative called "Exam Finding Not Present." This can be used to document that the provider actually performed the assessment, but did not find any injury/abnormality.

Vitals

- 1. Vitals Date/Time Group
 - a. eVitals.01 - Date/Time Vital Signs Taken
 - b. eVitals.02 - Obtained Prior to this Unit's EMS Care
- 2. Glasgow Coma Score (GCS) Group
 - a. Vitals Date/Time Group
 - b. eVitals.19 - Glasgow Coma Score-Eye
 - c. eVitals.20 - Glasgow Coma Score-Verbal
 - d. eVitals.21 - Glasgow Coma Score-Motor
 - e. eVitals.22 - Glasgow Coma Score-Qualifier
 - f. eVitals.23 - Total Glasgow Coma Score
- 3. ECG Rhythm Group
 - a. Vitals Date/Time Group
 - b. eVitals.03 - Cardiac Rhythm/Electrocardiography (ECG)
 - c. eVitals.04 - ECG Type
 - d. eVitals.05 - Method of ECG Interpretation
- 4. Temperature Group
 - a. Vitals Date/Time Group
 - b. eVitals.24 - Temperature
 - c. eVitals.25 - Temperature Method
- 5. Pain Scale Group
 - a. Vitals Date/Time Group
 - b. eVitals.27 - Pain Scale Score
 - c. eVitals.28 - Pain Scale Type
- 6. Stroke Score Group
 - a. Vitals Date/Time Group
 - b. eVitals.29 - Stroke Scale Score
 - c. eVitals.30 - Stroke Scale Type
- 7. Additional Vitals Options

All should have a value in the Vitals Date/Time Group and can be documented individually or as an add-on to basic, standard, or full vitals

 - a. eVitals.09 - Mean Arterial Pressure
 - b. eVitals.13 - Pulse Rhythm
 - c. eVitals.15 - Respiratory Effort

- d. eVitals.16 - End Tidal Carbon Dioxide (ETCO₂)
- e. eVitals.17 - Carbon Monoxide (CO)
- f. eVitals.18 - Blood glucose Level
- g. eVitals.26 - Level of Responsiveness (AVPU)
- h. Vitals.32 - APGAR
- 8. Routine Vitals – Includes the following vital signs:
 - a. Vitals Date/Time Group
 - b. Blood Pressure
 - c. eVitals.06 - SBP (Systolic Blood Pressure)
 - d. eVitals.07 - DBP (Diastolic Blood Pressure)
 - e. eVitals.10 - Heart Rate
 - f. eVitals.12 - Pulse Oximetry
 - g. eVitals.14 - Respiratory Rate
 - h. eVitals.26 - Level of Responsiveness (AVPU)
 - i. Pain Scale Group
- 9. Initial Vitals
 - a. Routine Vitals
 - b. eVitals.18 - Blood glucose Level
 - c. Glasgow Coma Score (GCS) Group
 - d. Temperature Group
- 10. Full Vitals
 - a. Initial Vitals
 - b. eVitals.13 - Pulse Rhythm
 - c. eVitals.15 - Respiratory Effort
 - d. eVitals.16 - End Tidal Carbon Dioxide (ETCO₂) (*If available and applicable*)
 - e. ECG Rhythm Group (*If available and applicable*)

Medications Given

- 1. eMedications.01 - Date/Time Medication Administered
- 2. eMedications.02 - Medication Administered Prior to this Unit's EMS Care
- 3. eMedications.03 - Medication Given
 - a. Pertinent Negatives (medication qualifiers) allowed
 - i. Contraindication Noted
 - ii. Medication Already Taken
 - iii. Denied By Order
 - iv. Refused
 - v. Medication Allergy
 - vi. Unable to Complete
- 4. eMedications.04 - Medication Administered Route
- 5. eMedications.05 - Medication Dosage
- 6. eMedications.06 - Medication Dosage Units
- 7. eMedications.07 - Response to Medication [*see Definitions of Medication Response below*]
- 8. eMedications.08 - Medication Complication
- 9. eMedications.09 - Medication Crew (Health care Professionals) ID (*Name or license #*)
- 10. eMedications.10 - Role/Type of Person Administering Medication (*License level*)

Procedures Performed

- 1. eProcedures.01 - Date/Time Procedure Performed
- 2. eProcedures.02 - Procedure Performed Prior to this Unit's EMS Care

3. eProcedures.03 – Procedure
 - a. Pertinent Negatives Allowed
 - i. Contraindication Noted
 - ii. Refused
 - iii. Denied By Order
 - iv. Unable to Complete
4. eProcedures.04 - Size of Procedure Equipment
5. eProcedures.05 - Number of Procedure Attempts (*This should always be "1" with each attempt at a procedure documented separately with appropriate date/time stamp*)
6. eProcedures.06 - Procedure Successful
7. eProcedures.07 - Procedure Complication
8. eProcedures.08 - Response to Procedure [*see Definitions for Response to Procedures below*]
9. eProcedures.09 - Procedure Crew Members ID
10. eProcedures.10 - Role/Type of Person Performing the Procedure
11. eProcedures.13 - Vascular Access Location (*If applicable*)

Narrative

The use of the narrative is essential to an effective and complete Patient Care Record. It summarizes the incident history and care in a manner that is easily digested between caregivers for continuity of care and provides a place for EMS to document facts that do not fit into fixed data fields [see **Narrative** section under **Notes and Educational Pearls** (below) for more detail]

Notes and Educational Pearls

Documenting Signs and Symptoms Versus Provider Impressions

1. Signs and Symptoms
 - a. Signs and Symptoms should support the provider impressions, treatment guidelines, and overall care given.
 - i. A symptom is something the patient experiences and tells the provider; it is subjective.
 - ii. A sign is something the provider sees; it is objective.
 - b. Symptoms should not be confused with provider impressions. The provider impressions are the EMS working field diagnosis of the patient's actual medical condition.
2. Provider Impressions
 - a. There is often a great deal of confusion on the part of EMS providers about the difference between symptoms and provider impressions. Provider impressions should be *supported* by symptoms but not *be* the symptoms except on *rare* occasions where they may be the same (e.g. weakness when no etiology for the weakness can be determined by the EMS provider).
 - b. Correctly documenting impressions is essential to many aspects of EMS data use, such as EMS reimbursement, reports of incident types, specialty registries (e.g. CARES) and CPI reviews. EMS agencies could *literally lose money or equipment and staffing resources* if the providers are incorrectly entering provider impressions. Addressing this issue should be an essential part of the record quality assurance and CQI process and documentation training.
 - c. Example of documenting symptoms versus impressions:
 - i. An opiate overdose patient who received naloxone and had a positive response: This patient would have possible *Symptoms* of altered mental

status, unconscious, respiratory distress, and respiratory failure or apnea. All four of these symptoms are available as provider impressions, however the correct impression for this patient would be whatever variation of “Drug Overdose Opiates or Heroin” impression(s) are setup in the local ePCR system being used. This impression will specifically define the call as an overdose with opiates, rather than a case where one of the symptoms was also used as an impression when the use of naloxone and other assessments and diagnostic tools could not determine an etiology for the symptom(s).

Narrative

The various data fields within the ePCR are important as they provide a means of uniformly entering incident data that can be used for importing into billing software or hospital records; transmitting between EMS systems or creating descriptive reports; or conducting research. In most cases, at a local, state, or national level, if something wasn’t documented in the appropriate data field, it didn’t happen or exist. However, the narrative plays several essential roles in the PCR.

1. Role of the Narrative

- a. Provides an efficient and effective means to share patient information for continuity of care between EMS services and EMS and hospital staff. The narrative summarizes the incident history and care in a manner that is easily digested when shared between caregivers.
- b. Provides a place for EMS to document facts that do not fit into fixed data fields. Specifically, this would include the detailed history of the scene, what the patient may have done or said or other aspects of the call that only the provider saw, heard, or did. The narrative is the place for the EMS provider to “paint the picture” for all others to more fully understand the incident.
- c. Provides a standard means to add essential details about medical history, exams, treatments, patient response, and changes in patient condition that can’t otherwise be effectively or clearly communicated.

2. Narrative Formats

Documentation by EMS providers demonstrates a wide variation of training and practice reinforcement. Most training programs provide limited instruction on how to properly document operational and clinical processes, and almost no practice. Most providers learn this skill on the job, and often proficient mentors are sparse. Therefore, it is essential that the EMS provider uses a standard format to ensure they are consistent and complete in their documentation. There are three standard formats for EMS documentation. EMS providers should choose the best match for them, master the format, and be consistent in its use.

- a. **Medical Narrative:** This format is the one most new EMS providers use as it is intuitive and easy to learn. Some more experienced providers use it as they find telling the story from start to finish works best to organize their thoughts. A drawback to this method is that it is easy to forget to include facts because of the lack of structure.
- b. **SOAP:** This format stands for **S**ubjective, **O**bjective, **A**ssessment, and **P**lan. This is a format that is very common in the medical field.
- c. **CHART:** This format stands for **C**omplaint, **H**istory, **A**ssessment, **R**x (Treatment) and **T**ransport. Each section’s content is clearly defined and consistent in format. It minimizes the likelihood of forgetting information and ensures documentation is consistent between records and providers. CHART is

the format most recommended as best practice by EMS legal authorities and is considered the standard in many EMS systems. A variation is DCHART, where the "D" stands for **D**ispatch (reason).

Medications Given Showing Positive Action Using Pertinent Negatives

For medications that are required by protocol (e.g. aspirin for cardiac chest pain), *pertinent negatives* should be used to show that a medication protocol was considered but was satisfied by other than provider action.

Example: EMS is called to a patient for cardiac chest pain. The patient has already taken 324 mg of aspirin by the time EMS arrives per 911 pre-arrival instructions. EMS providers should document this as a medication given, prior-to-arrival, with the best estimated time, and qualify the medication as "Medication Already Taken" using the pertinent negative.

Definitions for Response to Medications

1. Improved:
 - a. The medication had its intended therapeutic effect and the patient's symptoms decreased or clinical condition improved or resolved (the word "effective" could be generally be substituted for "improved").
 - b. If a patient had the intended therapeutic response to the medication, but a side effect that caused a clinical deterioration in another body system, then "Improved" should be chosen and the side effects documented as a complication (e.g. nitroglycerin improved chest pain but dropped the blood pressure).
2. Unchanged:
 - a. The medication was ineffective and had no intended therapeutic effect or had a sub- therapeutic and unnoticeable effect
AND
 - b. The patient condition did not deteriorate
3. Worse:
 - a. The patient condition deteriorated or continued to deteriorate because either the medication:
 - i. Was ineffective and had no intended therapeutic effect
or
 - ii. Had a sub-therapeutic effect that was unable to stop or reverse the decline in patient condition
or
 - iii. Was the wrong medication for the clinical situation and the therapeutic effect caused the condition to worsen (e.g. giving glucose to a patient with hyperglycemia/diabetic ketoacidosis)

Definitions for Response to Procedures

1. Not Applicable:

The nature of the procedure has no direct expected clinical response (e.g. patient assessment, 12-lead ECG acquisition).
2. Improved:
 - a. The procedure performed had the intended effective outcome and/or the patient's symptoms decreased or clinical condition improved or resolved (e.g. defibrillation resolved VF into a perfusing rhythm; intubation controlled the airway and allowed effective management of breathing).

- b. An effective procedure that caused an improvement in the patient condition may also have resulted in a procedure complication and the complication should be documented (e.g. intubation caused minor airway trauma, but the intubation successfully secured the airway).
- 3. Unchanged:
 - a. The procedure performed did not have the clinical effect intended, but did not directly worsen the patient's symptoms or clinical condition (e.g. attempted defibrillation and the person remained in VF);
or
 - b. Had a sub-therapeutic effect and the symptoms continued (e.g. a bandage applied to a bleeding wound failed to stop the bleeding);
or
 - c. The nature of the procedure has no direct expected clinical response (e.g. patient assessment).
Note: "Not Applicable" would also be appropriate to choose for these cases.
- 4. Worse:
 - a. The results of the procedure performed lead to a worsening of the patient's symptoms or condition (e.g. defibrillation converted VF into asystole, application of a splint caused significant increase in pain or loss of sensation and pulses).
 - b. In the case of worsening condition, documentation of the procedure complications may also be appropriate.
Note: Just because a patient got worse, doesn't necessarily mean the provider performed the procedure incorrectly.

NEMSIS Data Standards and Limitations

1. NEMSIS is a national dataset and standard used by all EMS software systems. Currently there are three versions of the data standard available for documentation and in which data is stored:
 - a. NEMSIS Version 2.2.1 (v2.2.1)
 - i. Adopted in 2006, there have been no changes since release.
 - ii. Most states or systems have used this standard since its release, and the majority of most states' data available since approximately 2016 is in this format.
 - iii. NEMSIS accepted v2.2.1 data through 12/31/2016, and some states may continue to collect data in this standard until they transition to NEMSIS v3 standards.
 - b. NEMSIS Version 3 (v3)
 - i. NEMSIS v3 was created and finalized in 2011 to replace v2.2.1 in order to allow the dataset to become more flexible for updates and adopt technical standards making linkage to other health records possible.
 - a. NEMSIS v3.3.4 was released in March 2014 and was the first version in production where live data was collected by services and states and subsequently submitted to NEMSIS. NEMSIS will continue to accept v3.3.4 data until 12/31/2017.
 - b. NEMSIS v3.4, released in March 2015, included both changed elements and many added values to existing elements. NEMSIS has been accepting data from this version concurrently with V3.3.4 data. As of 01/01/2018, v3.4 will be the only standard and V3.3.4 will be phased out. All documentation guidelines found in this document are based on the NEMSIS v3.4 dataset

and standard.

2. Mandatory and Required Elements
 - b. *Mandatory*: NEMSIS makes certain elements or fields mandatory so, if not included, the record cannot be properly stored or moved electronically. These fields require real data and do not accept Nil (Blank) values, Not Values, or Pertinent Negatives.
 - c. *Required*: NEMSIS requires these elements or fields to be completed or the record cannot be properly stored or moved electronically. However, required fields allow Nil (blank) values, Not Values, or Pertinent Negatives to be entered and submitted.
 - d. State and local systems may have Mandatory or Required fields that are not Mandatory or Required by NEMSIS. The manager for these systems should be contacted for a list of these fields.
3. Not Values, Nil, and Pertinent Negatives
 - b. Not Values (NV), Nil, and Pertinent Negatives (PN) are values that are attributes of certain NEMSIS elements designed to clarify a null data entry or qualify data entry into the element with which the NV, Nil, or PN is associated.
 - c. Not Values available are "Not Applicable" and "Not Recorded."
 - i. Some NEMSIS rules require one of these values to be entered when data is imported or exported if there is no other data in a field (e.g. at least one medication given, it must have a value, if no medications are given, then the software system must insert "Not Applicable" in the medications field when exporting).
 - ii. At times the EMS provider use of "Not Applicable" is appropriate documentation (e.g. using "Not Applicable" under *eInjury.03 - Trauma Center Criteria*, which combines step 1 and 2 of CDC's Guidelines for Field Triage of Injured Patients, when transporting a patient with a simple sprained ankle).
 - d. Nil Values are blank values
 - i. Values can be left blank, which can either be an accidental or purposeful omission of data.
 - ii. Value fields can appropriately and purposefully be left blank if there was nothing to enter (e.g. a procedure field left blank if no patient was encountered).
 - e. Pertinent Negatives are attributes or qualifiers for both elements and fields. There are 11 possible Pertinent Negative values and the available list for each field varies as appropriate to the field. Two examples of the use of Pertinent Negatives are:
 - i. Documenting non-administration of aspirin for chest pain by the EMS provider with the Pertinent Negative of "Medication Already Taken" to show evidence that this treatment requirement was met.
 - ii. Documenting assessment of, and lack of, a gunshot wound to the chest with the qualifier of "Chest --> gunshot wound --> Exam Finding Not Present" in the examination section (previously you could only document a positive finding of a gunshot wound with no way to document that you looked and did not find one).
4. NEMSIS Element and Value Name Formats
 - a. NEMSIS Elements and Fields are organized into groups with other related elements and fields
 - i. There are two parent datasets: Demographic (designated by a "d") and

- EMS (designated by an “e”). The majority of the documentation in any ePCR falls in the “e” section. The Demographic dataset is intended to be descriptive of the EMS agencies and system characteristics for correlation at a larger research level, rather than for use in operational CPI reviews.
- ii. The element numbering structure reflects the dataset and the text group name of the element.
 - b. Example: “eVitals.06 - SBP (Systolic Blood Pressure)” where “e” is the EMS dataset and “Vitals” is the dataset grouping for all elements related to Vitals and the number is the number assigned to a specific element.
 - i. “eVitals.06” is used to store the data in the background and “SBP (Systolic Blood Pressure)” is what providers and reviewers see.
 - ii. Values are designated by a code and text name.
 - The codes are generally derived from various sources such as ICD-10, SNOMED, or RxNorm and are used to store and move the data in the system’s background.
 - Codes are not seen by the EMS provider in the ePCR, but rather the provider will see text names.

Some software systems allow the visible text name to be modified or relabeled to meet local standards or nomenclature. This feature can help improve data quality by making documentation easier for the provider.
 - An example of a value code and name for cardiac chest pain, found under the element “eProtocols.01 - Protocols Used” is “9914117—Medical-Cardiac Chest Pain”.
 - c. All minimum general documentation guideline requirements are identified using the NEMSIS element, values codes, and names to allow application across a variety of ePCR software labels for these fields.
5. Custom Elements, Fields and Values
- b. The NEMSIS Standard provides a data format for software vendors to create custom elements or values requested by states or local systems.
 - c. States or local systems may create new elements or value extensions for existing NEMSIS elements to meet regional needs (e.g. adding additional protocol name values not on the NEMSIS list).

Airway Confirmation Fields

Specific use of the NEMSIS airway confirmation fields in documentation will not be detailed at this time due to current operational and technical challenges all states, local systems, and ePCR software vendors are experiencing.

The NEMSIS airway confirmation fields were closely modeled on the “Recommended Guidelines for Uniform Reporting of Data from Out-of-Hospital Airway Management: Position Statement of the National Association of EMS Physicians” and the fields and values could provide excellent and appropriately useful data to evaluate airway management. However, the technical structure of the fields has made their practical use limited as all the data is collected as a separate, self-contained group, rather than as part of the procedures group. This means EMS providers would need to enter much of

the same information twice in the ePCR, in both the procedures area and airway confirmation section (when, who did it, what device was used, and complications).

Furthermore, the airway group can only be entered once per ePRC, so the fields cannot be used again if more than one airway was required (e.g. one airway became ineffective and needed to be replaced with a different type of airway). Many states and ePCR software vendors have been struggling with how to make these fields functional for use by only using a portion of them or looking to add mirrored custom values that are directly linked to procedures performed. However, solutions are currently far from practical, functional, effective, or uniform in how they are being implemented or used across various systems.

II. 2021 Scope of Practice Medications

The following appendix contains the medication doses for respective protocol or indication

Drug (A-Z)	Protocol or Indication	Adult Dosage -Typical ranges -Repeat as directed or indicated	Pediatric Dosage -Typical ranges -Repeat as directed or indicated -Do not exceed adult dose unless indicated
Acetaminophen	-Pain management -Seizure (febrile) -Shock (sepsis)	15mg/kg PR/IV/IO/PO (max dose 1000mg)	15mg/kg PR/IV/IO/PO (max dose 1000mg)
Activate charcoal	-Poisoning (universal) -Calcium channel blocker overdose	1g/kg PO/NG	1g/kg PO/NG
Adenosine	Tachycardia with pulse	6mg IV (proximal site) followed by 10ml fluid bolus, repeat 12mg IV, as needed, up to three doses total	0.1mg/kg (maximum of 6mg), may repeat at 0.2mg/kg (maximum of 12 mg IV), as needed up to three doses total
Albuterol	-Anaphylaxis -Bronchospasm -Airway respiratory irritants	2.5–5mg nebulized or 6 puffs MDI	2.5–5mg nebulized or 6 puffs MDI
Amiodarone (preferred antiarrhythmic)	Tachycardia with pulse	150mg IV/IO over 10 minutes, may repeat	5mg/kg IV/IO (max of 150mg) over 10 minutes
	-Ventricular fibrillation -Pulseless ventricular tachycardia	5mg/kg IV/IO (max 300mg)	5mg/kg IV/IO (max 300mg)
Aspirin	Acute Coronary Syndrome	162 to 325mg chewable	Seek expert on-line medical control consultation
Atropine	-Bradycardia -Beta blocker overdose -Calcium channel blocker overdose	1mg IV/IO q5minutes (maximum total dose of 3mg)	0.02mg/kg IV/IO q5 minutes (maximum total dose of 3mg)
	Acetylcholinesterase inhibitor exposure	Refer to tables	Refer to tables
Calcium gluconate (preferred preparation)	-Hyperglycemia (hyperkalemia) -Resuscitation -Eclampsia -Preeclampsia -Calcium channel blocker overdose	3 grams IV/IO push over 2 mins	60 mg/kg max dose 3 grams IV/IO push over 2 mins
Dextrose (D10)	Hypoglycemia	4 ml/kg max of 100ml; repeat PRN	4 ml/kg max of 100ml; repeat PRN

Diazepam	Seizure	0.2 mg/kg PR max dose of 10mg	0.2 mg/kg PR max dose of 10mg
	Acetylcholinesterase inhibitors	Refer to tables	Refer to tables
Diltiazem	Tachycardia with pulse	0.25mg/kg slowly IV/IO over 2 minutes; after 15 minutes a second dose of diltiazem 0.35 mg/kg IV may be given if needed. For patients older than 65, recommend maximum initial dose of diltiazem 10 mg IV and a maximum second dose of 20mg	
Diphenhydramine	-Anaphylaxis -Allergic reaction -Nausea and vomiting -Poisoning	1mg/kg IM/IV/PO (maximum dose of 25mg)	1mg/kg IM/IV/PO (maximum dose of 25mg)
Dexamethasone	-Croup -Bronchospasm -Allergic reaction	0.5 mg/kg max 16 mg	0.5 mg/kg max 16 mg
Epinephrine 1:1,000	Anaphylaxis	>30 kg (66lbs.) Adult auto-injector or 0.3 mg IM; repeat q 5-15 mins PRN	<30 kg (66 lbs.) Pediatric auto-injector or 0.15 mg IM; repeat q 5-15 mins PRN
	Bronchospasm	0.3 mg IM; repeat q 5-15 mins PRN	0.15 mg IM; repeat q 5-15 mins PRN
	Bronchiolitis	n/a	2.5–5 ml nebulized
	Croup	n/a	2.5–5 ml nebulized
Epinephrine 1:10,000	Resuscitation	1mg IV/IO q 3-5 mins	0.01 mg/kg IV/IO q 3-5 mins max 1mg
Epinephrine 1:10,000 as Push Dose Pressor (mix 1ml of epinephrine 1:10,000 in 9ml of NS yielding a new mixture of 100 mcg in 10 ml which is also 10 mcg in every 1 ml.	-Bradycardia -Distributive shock -Cardiogenic shock	5–20 mcg per 3-5 minutes titrated to MAP 65; by volume, this is 0.2 ml to 1ml	Seek expert on-line medical control consultation
Famotidine	-Anaphylaxis -Allergic reaction	40 mg PO 20mg IV	0.5 mg/kg orally

			0.25 mg/kg IV injected over at least 2 minutes
Fentanyl (preferred opioid)	Pain management	0.5–1mcg/kg IV/IM/IN (maximum initial dose of 100mcg)	0.5–1mcg/kg IV/IM/IN (maximum initial dose of 100mcg)
Glucagon	Hypoglycemia	1mg IM/IN	<20 kg: 0.5 mg IM/IN >20kg: 1 mg IM/IN
	-Beta blocker overdose -Calcium channel blocker overdose	5mg IVIO, can be repeated in 5-10 minutes as necessary	1mg IVP (25-40kg) or 0.5mg IVP (≤25kg) every 5-10 minutes as necessary
Glucose (oral)	-Altered mental status -Hypoglycemia	25 grams PO	0.5–1mg/kg PO
Haloperidol	-Agitated patient -Violent patient -End-of-life and palliative care	5mg IM/IV	0.15 mg/kg max of 5 mg
Hydrocortisone	Shock	2mg/kg IV/IO/IM max of 100 mg	2mg/kg IV/IO/IM max of 100 mg
Hydroxocobalamin	Cyanide exposure	5 grams IV/IO administered over 15 minutes	70mg/kg (max 5g) IV/IO administered over 15 minutes
Ibuprofen	-Pain -Seizure (febrile)	10mg/kg PO max 600 mg	10mg/kg PO max 600 mg
Ipratropium	-Bronchospasm -Airway respiratory irritant	0.5 mg nebulized max of 1.5 mg total dosage	0.5 mg nebulized max of 1.5 mg total dosage
Ketamine	-Agitated patient -Combative patient	1–2 mg/kg IV or 3–4 mg/kg IM	1–2 mg/kg IV or 3–4 mg/kg IM
	Pain control	0.5mg/kg IV/IO/IM/IN (max initial dose of 25mg, maximum cumulative dose of 100mg)	0.5mg/kg IV/IO/IM/IN (max initial dose of 25mg, maximum cumulative dose of 100mg)
Ketorolac	-Pain management -Seizures (febrile)	15mg IV/IO or 30mg IM	0.5mg/kg IV/IO (max dose of 15mg) or 1mg/kg IM (max dose of 30mg)
Lidocaine	-Tachycardia with pulse -Resuscitation	1 mg/kg IV/IO may be repeated q 5 mins max total dose of 3 mg/kg	1 mg/kg IV/IO may be repeated q 5 mins max total dose of 3 mg/kg
	Analgesia for IO insertion	1mg/kg max 40 mg IO	1mg/kg max 40 mg IO

Magnesium	-Tachycardia with pulse -Bronchospasm	40mg/kg IV (max dose of 2g) over 10 minutes	40mg/kg IV (max dose of 2g) over 10 minutes
-----------	--	---	---

	Resuscitation	40mg/kg IV (max dose of 2g) bolus	40mg/kg IV (max dose of 2g) bolus
	-Eclampsia -Pre-eclampsia	4g IV (20% solution) over 20 min, followed by 1g/hour IV	Seek expert on-line medical control consultation
Methylprednisolone	-Shock -Bronchospasm	2mg/kg IV/IO (maximum 125mg)	2mg/kg IV/IO (maximum 125mg)
Metoprolol	Tachycardia with pulse	5mg IV over 1–2 mins; repeat q 5mins x 3 doses	Seek expert on-line medical control consultation
Midazolam (preferred benzodiazepine)	Agitated patient Violent patient End of life and palliative care	2mg IM/IN/IV/IO	0.1 mg/kg max 2mg IM/IN/IV/IO
	-Seizure -Hyperthermia (seizure) -Overdose (seizure)	4mg IV/IO/IN 10 mg IM	0.1 mg/kg IV/IO/IN 0.25 mg/kg IM
Naloxone	-Altered mental status -Opioid overdose	Adult auto-injector Or 0.4-2mg IV/IM/ETT or up to 4mg IN	Pediatric auto-injector Or 0.4-2mg IV/IM/ETT or up to 4mg IN
Nitroglycerin	Acute coronary syndrome	0.4mg SL, can repeat 3-5 minutes as long as SBP>100mgHg	Seek expert on-line medical control consultation
	Congestive heart failure with hypertension	1.2 mg SL, can repeat 3-5 minutes as long as SBP>100mgHg	Seek expert on-line medical control consultation
Nitrous oxide	Analgesia	20–50% concentrated mixed with oxygen and self-administered with constant monitoring	20–50% concentrated mixed with oxygen and self-administered with constant monitoring
Norepinephrine	-Bradycardia -Shock	0.05–0.5mcg/kg/minute titrated to MAP greater than 65mgHg	0.05–0.5mcg/kg/minute titrated to age appropriate BP
Ondansetron (preferred antiemetic)	Nausea and vomiting	4mg IV/PO/SL or ODT	0.15mg/kg IV/PO (max dose of 4mg)
Oxygen	Universal care for dyspnea or distress	Target of achieving greater than 93% saturation for most acutely ill patients	Target of achieving greater than 93% saturation for most acutely ill patients
Pralidoxime (2-PAM)	Acetylcholinesterase inhibitors exposure	Refer to tables	Refer to table

Procainamide	Tachycardia with pulse	20-50mg/min until arrhythmia suppressed, hypotension ensues, or QRS duration increases greater than 50% or maximum dose 17mg/kg given	Seek expert on-line medical control consultation
Sodium bicarbonate	-Hyperkalemia -Poisoning -Beta blocker overdose	1mEq/kg IV/IO max 50 mEq over 5 mins	1mEq/kg IV/IO max 50 mEq over 5 mins
	Resuscitation	1mEq/kg IV/IO max 50 mEq bolus	1mEq/kg IV/IO max 50 mEq bolus
Sodium thiosulfate	Cyanide exposure	12.5g IV	0.5g/kg IV max of 12.5g

III. Medications

The project team considered the use of Institute for Safe Medication Practices (ISMP) Tall Man Letters methodology to avoid the miscommunication of lookalike drug names. Upon review of the list and the limited number of medications carried by EMS, as well as the expected use of this document, it was elected not to institute this measure into our medication list. We recommend EMS agencies consider incorporating these measures into practice where appropriate.

Additional information regarding Tall Man Letters can be found on the ISMP website: <http://www.ismp.org/tools/tallmanletters.pdf> and the US Food and Drug Administration website: <http://www.fda.gov/Drugs/DrugSafety/MedicationErrors/ucm164587.htm>.

Acetazolamide

Name: Diamox Sequels®

Class: Carbonic anhydrase inhibitors

Pharmacologic Action: Inhibits hydrogen ion excretion in renal tubule, increasing sodium, potassium, bicarbonate, and water excretion and producing alkaline diuresis.

Indications: Acute mountain sickness

Contraindications: Known hypokalemia/hyponatremia, hypersensitivity to acetazolamide or sulfa, liver disease, renal disease, cirrhosis, long term administration in patients with chronic, noncongestive angle- closure glaucoma

Acetaminophen

Name: There are multiple over-the-counter medications, as well as scheduled drugs, that include acetaminophen (Tylenol®) as an active ingredient.

Class: Analgesics, antipyretic, other

Pharmacologic Action: May work peripherally to block pain impulse generation; may also inhibit prostaglandin synthesis in CNS

Indications: Pain control, fever control

Contraindications: Hypersensitivity, severe acute liver disease

Acetic acid (vinegar)

Name: Vinegar

Class: Other

Pharmacologic Action: Stabilizes nematocyst discharge in non-United States jellyfish thus decreasing pain.

Indications: Pain control for jellyfish envenomation (outside of the United States (US))

Contraindications: May increase nematocyst discharge for US jellyfish and therefore should be used outside of the US only

Acetylcysteine

Name: Mucomyst®, Acetadote®

Class: Antidotes, other

Pharmacologic Action: Acts as sulfhydryl group donor to restore liver glutathione; may also scavenge free radicals to prevent delayed hepatotoxicity as antioxidant; encourages sulfation pathway of metabolism for acetaminophen

Indications: Antidote for acetaminophen overdose

Contraindications: Acute asthma

WARNING: Nausea and vomiting are common adverse effects following the oral administration of acetylcysteine

Activated Charcoal Name: Actidose-Aqua®

Class: Antidotes, other

Pharmacologic Action: Adsorbs a variety of drugs and chemicals (e.g. physical binding of a molecule to the surface of charcoal particles); desorption of bound particles may occur unless the ratio of charcoal to toxin is extremely high

Indications: Overdose and poisoning

Contraindications: Unprotected airway (beware of aspiration), caustic ingestions, intestinal obstruction

Adenosine

Name: Antidysrhythmics

Pharmacologic Action: Slows conduction through AV node and interrupts AV reentry pathways, which restore normal sinus symptoms

Indications: Conversion of regular, narrow complex tachycardia; stable supraventricular tachycardia (SVT) or regular, monomorphic wide complex tachycardia

Contraindications: Hypersensitivity, second or third degree AV Block (except those on pacemakers), sick sinus syndrome, atrial flutter or fibrillation, ventricular tachycardia

Albuterol

Name: Proventil®, Ventolin®, Proair®, Accuneb®

Class: Beta-2 agonist

Pharmacologic Action: Beta-2 receptor agonist with some beta-1 activity; relaxes bronchial smooth muscle with little effect on heart rate

Indications: Bronchospastic lung disease

Contraindications: Hypersensitivity, tachycardia secondary to heart condition

Amiodarone

Name: Pacerone®, Cordarone®, Nexterone®

Class: Class III antidysrhythmics

Pharmacologic Action: Class III antidysrhythmic agent, which inhibits adrenergic stimulation; affects sodium, potassium, and calcium channels; markedly prolongs action potential and repolarization; decreases AV conduction and sinus node function

Indications: Management of regular wide complex tachycardia in stable patients, irregular wide complex tachycardia in stable patients, and as antidysrhythmic for the management of ventricular fibrillation (VF) and pulseless ventricular tachycardia (VT)

Contraindications: Hypersensitivity, severe sinus node dysfunction, second degree or third degree heart block or bradycardia causing syncope (except with functioning artificial pacemaker), cardiogenic shock

WARNING: Avoid during breastfeeding.

Amyl Nitrite

Name: Component of the Cyanide Antidote Kit®

Class: Cyanide antidote

Pharmacologic Action: Reacts with hemoglobin to form methemoglobin, an oxidized form of hemoglobin incapable of oxygen transport but with high affinity for cyanide. Cyanide preferentially binds to methemoglobin over cytochrome a3, forming the nontoxic cyanomethemoglobin

Indications: Acute cyanide toxicity

Contraindications: None in the case of suspected pure cyanide toxicity noted, documented hypersensitivity, suspected or confirmed smoke inhalation and/or carbon monoxide poisoning
WARNING: There is a risk of worsening hypoxia due to methemoglobin formation.

Aspirin

Name: Multiple over-the-counter medications, as well as scheduled drugs, include aspirin as an active ingredient. These include, but are not limited to, Bayer Buffered Aspirin®, Alka-Seltzer with Aspirin®, Ascriptin®, Bayer Women's Low Dose®, Ecotrin®

Class: Antiplatelet agent, non-steroidal anti-inflammatory drug (NSAID)

Pharmacologic Action: Inhibits synthesis of prostaglandin by cyclooxygenase; inhibits platelet aggregation; has antipyretic and analgesic activity

Indications: Antiplatelet agent for the care of patients suspected of suffering from an acute coronary syndrome

Contraindications: Hypersensitivity to aspirin or NSAIDs (aspirin-associated hypersensitivity reactions include aspirin-induced urticarial or aspirin-intolerant asthma), bleeding GI ulcers, hemolytic anemia from pyruvate kinase (PK) and glucose-6-phosphate dehydrogenase (G6PD) deficiency, hemophilia, hemorrhagic diathesis, hemorrhoids, lactating mother, nasal polyps associated with asthma, sarcoidosis, thrombocytopenia, ulcerative colitis

Atropine

Name: Atropen®, a component of Mark I® kits and DuoDote®

Class: Anticholinergic, toxicity antidotes

Pharmacologic Action: Competitively inhibits action of acetylcholinesterase on autonomic effectors innervated by postganglionic nerves

Indications: Management of nerve agent toxicity, symptomatic bradycardia (primary or related to toxin ingestion), organophosphate and carbamate insecticide toxicity

Note: Ineffective in hypothermic bradycardia

Contraindications: No absolute contraindications for ACLS, documented hypersensitivity in non- ACLS, nerve agent, and organophosphate scenarios

Relative contraindications: Narrow-angle glaucoma, GI obstruction, severe ulcerative colitis, toxic megacolon, bladder outlet obstruction, myasthenia gravis, hemorrhage with cardiovascular instability, thyrotoxicosis

Calcium Chloride

Name: Calcium Chloride

Class: Antidotes, other; calcium salts

Pharmacologic Action: Bone mineral component; cofactor in enzymatic reactions; essential for neurotransmission, muscle contraction, and many signal transduction pathways

Indications: For use in topical burns (hydrofluoric acid) or for use in calcium channel blocker overdose

Contraindications: Hypercalcemia, documented hypersensitivity, life-threatening cardiac arrhythmias may occur in known or suspected severe hypokalemia

WARNING: There is a risk for digitalis toxicity. Be cautious of peripheral IV use as significant tissue necrosis at injection site may occur.

Calcium Gluconate

Name: Gluconate®

Class: Antidotes, calcium salts, other

Pharmacologic Action: Bone mineral component; cofactor in enzymatic reactions; essential for neurotransmission, muscle contraction, and many signal transduction pathways

Indications: For use in topical burns (hydrofluoric acid) or for use in calcium channel blocker overdose

Contraindications: Hypercalcemia, documented hypersensitivity, sarcoidosis, life-threatening cardiac arrhythmias may occur in known or suspected severe hypokalemia

WARNING: There is a risk for digitalis toxicity.

Cimetidine

Name: Tagamet®

Class: Histamine H2 antagonist

Pharmacologic Action: Blocks H2-receptors of gastric parietal cells, leading to inhibition of gastric secretions

Indications: For the management of gastric or duodenal ulcers, gastroesophageal reflux, as an adjunct in the treatment of urticarial and/or pruritus in patients suffering from allergic reaction

Contraindications: Hypersensitivity to cimetidine or other H2-receptor antagonists

Dexamethasone

Name: Decadron®, Dexasone®

Class: Corticosteroid, anti-inflammatory drugs

Pharmacologic Action: Potent glucocorticoid with minimal to no mineralocorticoid activity; decreases inflammation by suppressing migration of polymorphonuclear leukocytes (PMNs) and reducing capillary permeability; stabilizes cell and lysosomal membranes; increases surfactant synthesis; increases serum vitamin A concentration, and inhibits prostaglandin and proinflammatory cytokines; suppresses lymphocyte proliferation through direct cytolysis; inhibits mitosis; breaks down granulocyte aggregates; and improves pulmonary microcirculation

Indications: Used in the management of croup and bronchospasm, as well as the management of patients suffering from high altitude cerebral edema (HACE)

Contraindications: Documented hypersensitivity, systemic fungal infection, cerebral malaria

Dextrose

Name: D50W, DGlucose®, glucose

Class: Glucose-elevating agents; metabolic and endocrine; other

Pharmacologic Action: Parenteral dextrose is oxidized to carbon dioxide and water, and provides 3.4 kilocalories/gram of d-glucose

Indications: Used for the management of hypoglycemia

Contraindications: Hyperglycemia, anuria, diabetic coma, intracranial or intraspinal hemorrhage, dehydrated patients with delirium, glucose-galactose malabsorption syndrome, and documented hypersensitivity

Diazepam

Name: Valium®, Diastat®, AcuDial®

Class: Benzodiazepine, anticonvulsants, skeletal muscle relaxants, anxiolytic

Pharmacologic Action: Modulates postsynaptic effects of GABA-A transmission, resulting in an increase in presynaptic inhibition. Appears to act on part of the limbic system, as well as on the thalamus and hypothalamus, to induce a calming effect

Indications: For use in agitated or violent patients, as well as for the management of seizures

Contraindications: Documented hypersensitivity, severe respiratory depression

Diltiazem

Name: Includes Cardizem®, Dilacor®, Diltiaz®

Class: Calcium channel blocker, antidysrhythmic type IV

Pharmacologic Action: Inhibits extracellular calcium ion influx across membranes of myocardial cells and vascular smooth muscle cells, resulting in inhibition of cardiac and vascular smooth muscle contraction and thereby dilating main coronary and systemic arteries; no effect on serum calcium concentrations; substantial inhibitory effects on cardiac conduction system, acting principally at AV node, with some effects at sinus node

Indications: For management of narrow complex tachycardias

Contraindications: Documented hypersensitivity, Wolff-Parkinson-White syndrome, Lown-Ganong-Levine syndrome, symptomatic severe hypotension (systolic BP < 90 mm Hg), sick sinus syndrome (if no pacemaker), second and third degree heart block (if no pacemaker present), and complete heart block.

Contraindications for IV administration: Use in newborns (because of benzyl alcohol), concomitant beta-blocker therapy, cardiogenic shock, ventricular tachycardia (must determine whether origin is supraventricular or ventricular)

Diphenhydramine Name: Benadryl®

Class: Antihistamine—first generation

Pharmacologic Action: Histamine H1-receptor antagonist of effector cells in respiratory tract, blood vessels, and GI smooth muscle

Indications: For urticarial and/or pruritis in the management of patients suffering from allergic reaction as well as for the management of patients suffering from dystonia or akathisia

Contraindications: Documented hypersensitivity, use controversial in lower respiratory tract disease (such as acute asthma), premature infants and neonates

Dopamine

Name: Intropin®

Class: Inotropic agent; catecholamine; pressor

Pharmacologic Action: Endogenous catecholamine, acting on both dopaminergic and adrenergic neurons; low dose stimulates mainly dopaminergic receptors, producing renal and mesenteric vasodilation; higher dose stimulates both beta-1-adrenergic and dopaminergic receptors, producing cardiac stimulation and renal vasodilation; large dose stimulates alpha-adrenergic receptors

Indications: As a pressor agent used in the management of shock

Contraindications: Hypersensitivity to dopamine, pheochromocytoma, ventricular fibrillation, uncorrected tachyarrhythmias

WARNING: Dopamine is a vesicant and can cause severe tissue damage if extravasation occurs.

Droperidol

Name: Inapsine®

Class: Antiemetic agents; antipsychotic

Pharmacologic Action: Antiemesis: dopamine receptor blockade in brain, predominantly dopamine-2 receptor; when reuptake is prevented, a strong antidopaminergic, antiserotonergic response occurs; droperidol reduces motor activity, anxiety, and causes sedation; also possesses adrenergic-blocking, antifibrillatory, antihistaminic, and anticonvulsive properties

Indications: For use in the patient with acute delirium or psychosis

Contraindications: Hypersensitivity, known or suspected prolonged QT interval; QTc interval > 450 msec in females or > 440 msec in males

WARNING: Use with caution in patients with bradycardia, cardiac disease, concurrent MAO inhibitor therapy, Class I and Class III dysrhythmics or other drugs that prolong the QT interval and cause electrolyte disturbances due to its adverse cardiovascular effects, i.e. QT prolongation, hypotension, tachycardia, and torsades de pointes.

Epinephrine

Name: EpiPen®, TwinJect®, Adrenaclick®, Auvi-Q, Adrenalin®, AsthmaNefrin®, Vaponefrin®

Class Alpha/beta adrenergic agonist

Pharmacologic Action: Strong alpha-adrenergic effects, which cause an increase in cardiac output and heart rate, a decrease in renal perfusion and peripheral vascular resistance, and a variable effect on BP, resulting in systemic vasoconstriction and increased vascular permeability; strong beta-1- and moderate beta-2-adrenergic effects, resulting in bronchial smooth muscle relaxation; secondary relaxation effect on smooth muscle of stomach, intestine, uterus, and urinary bladder

Indications: For use in the management of patients suffering anaphylaxis, shock, cardiac arrest, bradycardia, or—in the nebulized form—for croup or bronchiolitis; and IM form for refractory acute asthma

Contraindications: Hypersensitivity, cardiac dilatation and coronary insufficiency

Famotidine Name: Pepcid®

Class: Histamine H2 antagonist

Pharmacologic Action: Blocks H2 receptors of gastric parietal cells, leading to inhibition of gastric secretions

Indications: For the management of gastric or duodenal ulcers, and gastroesophageal reflux, as an adjunct in the treatment of urticarial and/or pruritus in patients suffering from allergic reaction

Contraindications Hypersensitivity to famotidine or other H2-receptor antagonists

Fentanyl

Name: Currently only available in the generic form (formerly Sublimaze®)

Class: Synthetic opioid, opioid analgesics

Pharmacologic Action: Narcotic agonist-analgesic of opiate receptors; inhibits ascending pain pathways, thus altering response to pain; increases pain threshold; produces analgesia, respiratory depression, and sedation

Indications: Management of acute pain

Contraindications: Hypersensitivity

WARNING: Should be used with caution in the elderly and in patients with hypotension, suspected gastrointestinal obstruction, head injury, and concomitant CNS depressants.

Glucagon

Name: GlucaGen®, Glucagon Emergency Kit®, GlucaGen HypoKit®

Class: Hypoglycemia antidotes, glucose-elevating agents, other antidotes (e.g. beta-blocker or calcium channel blocker overdose)

Pharmacologic Action: Insulin antagonist; stimulates cAMP synthesis to accelerate hepatic glycogenolysis and gluconeogenesis; glucagon also relaxes smooth muscles of GI tract

Indications: For the management of hypoglycemic patients as well as patients suffering symptomatic bradycardia after beta blocker or calcium channel blocker overdose

Contraindications: Hypersensitivity, pheochromocytoma, insulinoma

WARNING: Nausea and vomiting are common adverse effects following the administration of glucagon.

Haloperidol

Name: Haldol®, Haldol Decanoate®, Haloperidol LA®, Peridol®

Class: First generation antipsychotic

Pharmacologic Action: Antagonizes dopamine-1 and dopamine-2 receptors in brain; depresses reticular activating system and inhibits release of hypothalamic and hypophyseal hormones

Indications: For the management of acute psychosis or agitated/violent behavior refractory to non-pharmacologic interventions

Contraindications: Documented hypersensitivity, Severe CNS depression (including coma), neuroleptic malignant syndrome, poorly controlled seizure disorder, Parkinson's disease

WARNING: Risk of sudden death, torsades de pointes, and prolonged QT interval from off-label IV administration of higher than recommended dose. Continuous ECG cardiac monitoring is required if administering IV.

Helium Gas

Mixture Name: Heliox®

Class: Optional method of oxygen delivery

Pharmacology: Less resistant than atmospheric air which may reduce the patient's work of breathing by increasing tendency to laminar flow and reducing resistance to turbulent flow

Indications: Persistent or severe bronchospasm in non-intubated patients with obstructive airway disease or pediatric patients with croup that is unresponsive to all other evidence-based medical interventions.

Contraindications: None

Hydralazine

Name: No listed brand name

Class: Vasodilator

Pharmacology: Direct vasodilator at the level of arterioles, with little effect on veins; decreases systemic resistance

Indications: Severe hypertension with pre-eclampsia symptoms

Contraindications: Hypersensitivity, coronary artery disease, mitral valve rheumatic heart disease. Use with caution in CVA, known renal disease, hypotension

Hydrocortisone succinate

Name: Cortef®, SoluCortef®

Class: Corticosteroid

Pharmacologic Action: Glucocorticoid; elicits mild mineralocorticoid activity and moderate anti-inflammatory effects; controls or prevents inflammation by controlling rate of protein

synthesis, suppressing migration of polymorphonuclear leukocytes (PMNs) and fibroblasts, and reversing capillary permeability

Indications: For the management of adrenal insufficiency

Contraindications: Untreated serious infections (except tuberculous, meningitis, or septic shock), idiopathic thrombocytopenic purpura, intrathecal administration (injection), documented hypersensitivity

Hydromorphone

Name: Dilaudid®

Class: Synthetic opiate, opioid analgesic

Pharmacology: Narcotic agonist-analgesic of opiate receptors; inhibits ascending pain pathways, thus altering response to pain; increases pain threshold; produces analgesia, respiratory depression, and sedation

Indications: Management of acute pain

Contraindications: Hypersensitivity

WARNING: Should be used with caution in the elderly and in patients with hypotension, suspected gastrointestinal obstruction, head injury, and concomitant CNS depressants.

Hydroxocobalamin

Name: Cyanokit®

Class: Cyanide antidote

Pharmacologic Action: Vitamin B12 with hydroxyl group complexed to cobalt which can be displaced by cyanide resulting in cyanocobalamin that is renally excreted

Indications: For the management of cyanide toxicity

Contraindications: Documented hypersensitivity

WARNING: Will cause discoloration of the skin and urine, can interfere with pulse oximetry. Due to its interference with certain diagnostic blood tests, the performance of prehospital phlebotomy is preferable prior to the administration of hydroxocobalamin.

Ibuprofen

Name: There are multiple over-the-counter medications that include ibuprofen, such as Advil®, Motrin®

Class: Non-steroidal anti-inflammatory drug (NSAID)

Pharmacologic Action: Inhibits synthesis of prostaglandins in body tissues by inhibiting at least 2 cyclo-oxygenase (COX) isoenzymes, COX-1 and COX-2; may inhibit chemotaxis, alter lymphocyte activity, decrease proinflammatory cytokine activity, and inhibit neutrophil aggregation; these effects may contribute to anti-inflammatory activity

Indications: For the acute management of pain or as an antipyretic

Contraindications: Aspirin allergy; perioperative pain in setting of coronary artery bypass graft (CABG) surgery; preterm infants with untreated proven or suspected infection; bleeding with active intracranial hemorrhage or GI bleed; thrombocytopenia, coagulation defects, proven or necrotizing enterocolitis, significant renal impairment, congenital heart disease where patency or the patent ductus arteriosus (PDA) is necessary for pulmonary or systemic blood flow

Ipratropium

Name: Atrovent®

Class: Anticholinergics, respiratory

Pharmacologic Action: Anticholinergic (parasympatholytic) agent; inhibits vagally mediated reflexes by antagonizing acetylcholine action; prevents increase in intracellular calcium concentration that is caused by interaction of acetylcholine with muscarinic receptors on

bronchial smooth muscle

Indications: For the management of asthma and COPD

Contraindications: Documented hypersensitivity to ipratropium, atropine, or derivatives.

Isopropyl Alcohol

Name: No brand name available

Class: Secondary alcohol

Pharmacology: In addition to traditional role as antiseptic, may be used as antiemetic

Indications: Nausea and vomiting

Contraindications: None

Ketamine

Name: Ketalar®

Class: General anesthetics, systemic

Pharmacologic Action: Produces dissociative anesthesia; blocks N-methyl D-aspartate (NMDA) receptor

Indications: For the management of agitated or violent behavior

Contraindications: Hypersensitivity

Relative, controversial contraindications: Head trauma, intracranial mass or hemorrhage, hypertension, angina, and stroke, underlying psychiatric disorder

WARNING: Overdose may lead to panic attacks and aggressive behavior; rarely seizures, increased ICP, and cardiac arrest. Very similar in chemical makeup to PCP (phencyclidine), but it is shorter acting and less toxic

Ketorolac

Name: Toradol®

Class: Non-steroidal anti-inflammatory drug (NSAID)

Pharmacologic Action: Inhibits synthesis of prostaglandins in body tissues by inhibiting at least 2 cyclo-oxygenase (COX) isoenzymes, COX-1 and COX-2; may inhibit chemotaxis, alter lymphocyte activity, decrease proinflammatory cytokine activity, and inhibit neutrophil aggregation; these effects may contribute to anti-inflammatory activity

Indications: For the acute management of moderately severe pain

Contraindications: Allergy to aspirin, ketorolac, or other NSAIDs; women who are in active labor or are breastfeeding, significant renal impairment particularly when associated with volume depletion, previous or current GI bleeding, intracranial bleeding, coagulation defects, patients with a high risk of bleeding

Labetalol

Name: Trandate®

Class: Beta blockers, alpha activity

Pharmacology: Nonselective beta blocker with intrinsic sympathomimetic activity; also alpha blocker

Indications: Severe hypertension with pre-eclampsia symptoms

Contraindications: Asthma or obstructive airway disease; severe bradycardia; second-degree or third-degree heart block (without pacemaker); cardiogenic shock; bronchial asthma; uncompensated cardiac failure; hypersensitivity; sinus bradycardia; sick sinus syndrome without permanent pacemaker; conditions associated with prolonged and severe hypotension. Use with caution in patients taking calcium channel blockers. Hypotension with or without syncope may occur; monitor. Consider pre-existing conditions, such as, sick sinus syndrome before initiating therapy. Use caution in patients with history of severe anaphylaxis to allergens; patients taking beta-blockers may become more sensitive to repeated challenges;

treatment with epinephrine in patients taking beta-blockers may be ineffective or promote undesirable effects. Use with caution in patients with myasthenia gravis, psoriasis, or psychiatric illness (may cause or exacerbate CNS depression).

Lidocaine

Name: Lidocaine CV®, Lidopen®, Xylocaine®

Class: Class Ib antidysrhythmics

Pharmacologic Action: Class 1b antidysrhythmic; combines with fast sodium channels and thereby inhibits recovery after repolarization, resulting in decreasing myocardial excitability and conduction velocity

Indications: For the management of refractory or recurrent ventricular fibrillation or pulseless VT

Contraindications: Hypersensitivity to lidocaine or amide-type local anesthetic, Adams-Stokes syndrome, SA/AV/intraventricular heart block in the absence of artificial pacemaker; CHF, cardiogenic shock, second and third degree heart block (if no pacemaker is present), Wolff-Parkinson-White Syndrome

Lorazepam

Name: Ativan®

Class: Anticonvulsants, other; antianxiety agent; anxiolytics; benzodiazepines

Pharmacologic Action: Sedative hypnotic with short onset of effects and relatively long half-life; by increasing the action of gamma-aminobutyric acid (GABA), which is a major inhibitory neurotransmitter in the brain, lorazepam may depress all levels of the CNS, including limbic and reticular formation

Indications: For the management of seizures, uncontrolled shivering in hypothermia, and for the management of agitated or violent patients suffering behavioral emergencies

Contraindications: Documented hypersensitivity, acute narrow angle glaucoma, severe respiratory depression, sleep apnea

Magnesium sulfate

Name: MgSO₄

Class: Class V antidysrhythmic, electrolyte

Pharmacologic Action: Depresses CNS, blocks peripheral neuromuscular transmission, produces anticonvulsant effects; decreases amount of acetylcholine released at end-plate by motor nerve impulse; slows rate of sino-atrial (SA) node impulse formation in myocardium and prolongs conduction time; promotes movement of calcium, potassium, and sodium in and out of cells and stabilizes excitable membranes

Indications: For the management of torsades de pointes or for severe bronchoconstriction with impending respiratory failure, seizure during the third trimester of pregnancy, or in the postpartum patient

Contraindications: Hypersensitivity, myocardial damage, diabetic coma, heart block, hypermagnesemia, hypercalcemia

Methylprednisolone

Name: Medrol®, Medrol Dosepak®, DepoMedrol®, SoluMedrol®

Class: Corticosteroid, anti-inflammatory agent

Pharmacologic Action: Potent glucocorticoid with minimal to no mineralocorticoid activity; modulates carbohydrate, protein, and lipid metabolism and maintenance of fluid and electrolyte homeostasis; controls or prevents inflammation by controlling rate of protein synthesis, suppressing

migration of polymorphonuclear leukocytes (PMNs) and fibroblasts, reversing capillary permeability, and stabilizing lysosomes at cellular level

Indications: For the management of acute bronchospastic disease as well as for adrenal insufficiency

Contraindications: Untreated serious infections, documented hypersensitivity, IM route is contraindicated in idiopathic thrombocytopenic purpura, traumatic brain injury (high doses)

Metoclopramide

Name: Reglan®, Metozolv ODT®

Class: Antiemetic agent, prokinetic agent

Pharmacologic Action: Blocks dopamine receptors (at high dose) and serotonin receptors in chemoreceptor trigger zone of CNS and sensitizes tissues to acetylcholine; increases upper GI motility but not secretions; increases lower esophageal sphincter tone

Indications: For the management of nausea and vomiting

Contraindications: Hypersensitivity to metoclopramide or procainamide, GI hemorrhage, mechanical obstruction, perforation, history of seizures, pheochromocytoma; other drugs causing extrapyramidal symptoms (e.g. phenothiazines, butyrophenones)

Metoprolol

Name: Lopressor®, Toprol XL®

Class: Beta blocker, beta-1 selective

Pharmacologic Action: Blocks response to beta-adrenergic stimulation; cardio selective for beta-1 receptors at low doses, with little or no effect on beta-2 receptors

Indications: For management of narrow complex tachycardias

Contraindications: Hypersensitivity; *when administered for hypertension or angina:* sinus bradycardia, second or third degree AV block, cardiogenic shock, sick sinus syndrome (unless permanent pacemaker in place), severe peripheral vascular disease, pheochromocytoma; *when administered for myocardial infarction:* severe sinus bradycardia with heart rate <45 beats/minute, systolic BP <100 mmHg, significant first-degree heart block (PR interval at least 0.24 seconds), moderate-to-severe cardiac failure

WARNING: May cause 1st, 2nd, or 3rd degree AV block

Midazolam

Name: Versed®

Class: Anticonvulsants, other; antianxiety agent; anxiolytics; benzodiazepines

Pharmacologic Action: Binds receptors at several sites within the CNS, including the limbic system and reticular formation; effects may be mediated through gamma-aminobutyric acid (GABA) receptor system; increase in neuronal membrane permeability to chloride ions enhances the inhibitory effects of GABA; the shift in chloride ions causes hyperpolarization (less excitability) and stabilization of the neuronal membrane

Indications: For the management of seizures, uncontrolled shivering in hypothermia, and for the management of agitated or violent patients suffering behavioral emergencies

Contraindications: Documented hypersensitivity, severe respiratory depression, sleep apnea

WARNING: May cause respiratory depression, arrest, or apnea

Morphine Sulfate

Name: MS Contin®, Avinza®, Depodur®, Duramorph®, Infumorph®, Astramorph®, Kadian®, MSO4

Class: Opioid analgesic

Pharmacologic Action: Narcotic agonist-analgesic of opiate receptors; inhibits ascending pain pathways, thus altering response to pain; produces analgesia, respiratory depression, and

sedation; suppresses cough by acting centrally in medulla

Indications: Management of acute pain

Contraindications: Hypersensitivity, paralytic ileus, toxin-mediated diarrhea, respiratory depression, acute or severe bronchial asthma, upper airway obstruction, GI obstruction (extended release), hypercarbia (immediate release tablets/solution), upper airway obstruction (epidural/intrathecal), heart failure due to chronic lung disease, head injuries, brain tumors, deliriums tremens, seizure disorders, during labor when premature birth anticipated (injectable formulation), cardiac arrhythmia, increased intracranial or cerebrospinal pressure, acute alcoholism, use after biliary tract surgery, surgical anastomosis (suppository formulation)

Naloxone

Name: Narcan®, EVZIO®

Class: Opioid reversal agent

Pharmacologic Action: Competitive opioid antagonist; synthetic congener of oxymorphone

Indications: Reversal of acute opioid toxicity

Contraindications: Hypersensitivity

WARNING: Administration of naloxone can result in the sudden onset of opiate withdrawal (agitation, tachycardia, pulmonary edema, nausea, vomiting, and, in neonates, seizures).

Nifedipine

Name: Procardia®, Adalat CC®, Nifedical®

Class: Calcium channel blocker

Pharmacologic Action: Calcium-channel blocker; inhibits transmembrane influx of extracellular calcium ions across myocardial and vascular smooth muscle cell membranes without changing serum calcium concentrations; this results in inhibition of cardiac and vascular smooth muscle contraction, thereby dilating main coronary and systemic arteries; vasodilation with decreased peripheral resistance and increased heart rate

Indications: For the management of high altitude pulmonary edema (HAPE)

Contraindications: Hypersensitivity to nifedipine or other calcium-channel blockers, cardiogenic shock, concomitant administration with strong CYP3A4 inducers (e.g. rifampin, rifabutin, phenobarbital, phenytoin, carbamazepine, St. John's wort) significantly reduces nifedipine efficacy, immediate release preparation (sublingually or orally) for urgent or emergent hypertension

Nitrous

Oxide

Name: N₂O

Class: Weak inhalational anesthetic

Pharmacologic Action: Its analgesic mechanism of action is described as opioid in nature and may involve a number of spinal neuromodulators; the anxiolytic effect is similar to that of benzodiazepine and may involve gamma aminobutyric (GABA) receptors; the anesthesia mechanism may involve GABA and possibly N-methyl-D-aspartate receptors as well; in general, the effect of nitrous oxide ceases as soon as the inhalation stops, with no residual effect

Indications: Analgesia in the patient who is capable of self-administration of this medication

Contraindications: Significant respiratory compromise, suspected abnormal air-filled cavities (e.g. pneumothorax, bowel obstruction, air embolism)

Relative contraindications: History of stroke, hypotension, pregnancy, known cardiac conditions, known vitamin B12 deficiency

Nitroglycerin

Name: Nitrostat®, Nitrolingual Pumpspray®, NitroQuick®

Class: Nitrates, anti-anginal

Pharmacologic Action: Organic nitrate which causes systemic venodilation, decreasing preload. Cellular mechanism: nitrate enters vascular smooth muscle and converted to nitric oxide (NO) leading to activation of cyclic guanosine monophosphate (cGMP) and vasodilation. Relaxes smooth muscle via dose-dependent dilation of arterial and venous beds to reduce both preload and afterload, and myocardial O₂ demand. Also improves coronary collateral circulation. Lower BP, increases heart rate, occasional paradoxical bradycardia

Indications: As an anti-anginal medication for the management of chest pain as well as a reducer of preload for patients suffering from acute pulmonary edema

Contraindications: Hypersensitivity, acute myocardial infarction, severe anemia, recent use of erectile dysfunction medications sildenafil (Viagra® within last 24 hours), tadalafil (Cialis® within last 48 hours), vardenafil (Levitra® within last 48 hours), or other phosphodiesterase-5 inhibitors. There is potential for dangerous hypotension, narrow angle glaucoma (controversial: may not be clinically significant). Nitrates are contraindicated in the presence of hypotension (SBP < 90 mm Hg or ≥30 mm Hg below baseline), extreme bradycardia (< 50 bpm), tachycardia in the absence of heart failure (> 100 bpm), and right ventricular infarction

Norepinephrine

Name: Levophed®, Levarterenol®

Class: Alpha/beta adrenergic agonist

Pharmacologic Action: Strong beta-1 and alpha-adrenergic effects and moderate beta-2 effects, which increase cardiac output and heart rate, decrease renal perfusion and peripheral vascular resistance, and cause variable BP effects

Indications: As a pressor agent used in the management of shock

Contraindications: Hypersensitivity, hypotension due to blood volume deficit, peripheral vascular thrombosis (except for lifesaving procedures)

Relative contraindications: concomitant use with some general anesthetics: chloroform, trichloroethylene, cyclopropane, halothane

WARNING: Norepinephrine is a vesicant and can cause severe tissue damage if extravasation occurs. Do not use in the same IV line as alkaline solutions as these may deactivate it.

Olanzapine

Name: Zyprexa®

Class: Antipsychotic, second generation, antimanic agents

Pharmacologic Action: May act through combination of dopamine and serotonin type 2 receptor site antagonism

Indications: For the management of agitated or violent patients suffering a behavioral emergency

Contraindications: Documented hypersensitivity

WARNING: Patients are at risk for severe sedation (including coma) or delirium after each injection and must be observed for at least 3 hours in registered facility with ready access to emergency response services. Patients are at significant risk of severe sedation when olanzapine is administered with benzodiazepines or to patients who have are taking benzodiazepines.

Ondansetron

Name: Zofran®, Zofran ODT®, Zuplenz®

Class: Antiemetic, selective 5-HT₃ antagonist

Pharmacologic Action: Mechanism not fully characterized; selective 5-HT₃ receptor antagonist; binds to 5-HT₃ receptors both in periphery and in CNS, with primary effects in GI tract. Has no effect on dopamine receptors and therefore does not cause extrapyramidal symptoms

Indications: For the management of nausea or vomiting

Note: ECG monitoring is recommended in patients who have electrolyte abnormalities, CHF, or bradyarrhythmias or who are also receiving other medications that cause QT prolongation.

Contraindications: Hypersensitivity, coadministration with apomorphine; combination reported to cause profound hypotension and loss of consciousness

WARNING: May cause dose-dependent QT prolongation, avoid in patients with congenital long QT syndrome

Oxymetazoline

Name: Afrin®, Duramist Plus®, Dristan 12 Hr®, Sinarest 12 Hour®, Vicks Sinus 12 Hour®

Class: Decongestants, intranasal

Pharmacologic Action: Alpha-adrenergic agonist; stimulates alpha-adrenergic receptors and produces vasoconstriction in the arterioles of the nasal mucosa

Indications: For the management of epistaxis in the patient suffering facial trauma

Contraindications: Hypersensitivity

Potassium iodide

Name: Pima Syrup®, SSKI®, ThyroSafe®, ThyroShield®

Class: Antidotes, other; antithyroid agents

Pharmacologic Action: As a thyroid protective agent: Systemically circulating potassium iodide is readily taken up by thyroid gland by sodium or iodide transporter in basal membrane; blocking the thyroid uptake of radioactive isotopes of iodine; concentration gradient of thyroid gland to plasma is 20-50:1

Indications: Indicated during environmental radiation emergency to block uptake of radioactive iodine isotopes in thyroid and reduce risk of thyroid cancer

Contraindications: Iodine sensitivity (although allergy to radiocontrast media, contact dermatitis from iodine-containing antibacterials, allergy to seafood should not be considered evidence of potassium iodide allergy), hyperthyroidism, respiratory failure

Pralidoxime chloride (2-PAM)

Name: Protopam®, 2PAM Antidote®, Pralidoxime Auto Injector®, a component of Mark I® kits and DuoDote®

Class: Cholinergic, toxicity antidote

Pharmacologic Action: Binds to organophosphates and breaks alkyl phosphate-cholinesterase bond to restore activity of acetylcholinesterase

Indications: For the management of toxicity caused by organophosphate insecticides and related nerve gases (e.g. tabun, sarin, soman)

Contraindications: Documented hypersensitivity

Procainamide

Name: Pronestyl®,

Procanbid® **Class:** Class Ia
antidysrhythmic

Pharmacologic Action: Class Ia (membrane stabilizing) antidysrhythmic agent; inhibits recovery after repolarization resulting in decreasing myocardial excitability and conduction velocity. Direct membrane depressant that decreases conduction velocity, prolongs refractoriness, decreases automaticity and reduces repolarization abnormalities

Indications: For the management of stable patients with regular, wide complex tachycardia

Contraindications: Hypersensitivity to procainamide or other ingredients, complete heart block, second or third degree AV block, systemic lupus erythematosus (SLE), torsades de pointes

Relative contraindication: Patients with QT prolongation

Prochlorperazine

Name: Compazine®

Class: Antiemetic agent; antipsychotics, phenothiazine

Pharmacologic Action: Antiemetic: antidopaminergic effect, blocking dopamine receptors in the brain, blocking vagus nerve in GI tract. Antipsychotic: Blocking mesolimbic dopamine receptors, and blocking alpha-adrenergic receptors (D1 and D2) in brain

Indications: For the management of nausea and vomiting

Contraindications: Documented hypersensitivity to phenothiazines, coma, severe CNS depression, concurrent use of large amounts of CNS depressants, poorly controlled seizure disorder, subcortical brain damage, pediatric surgery, children < 2 years or weighing < 9 kg

Sildenafil

Name: Revatio®, Viagra®

Class: Pulmonary artery hypertension therapy, PDE-5 inhibitors; phosphodiesterase-5 enzyme inhibitor

Pharmacologic Action: Inhibits PDE-5, increasing cyclic guanosine monophosphate (cGMP) to allow smooth-muscle relaxation

Indications: As an adjunct to descent in the management of high altitude pulmonary edema (HAPE)

Contraindications: Concomitant use of organic nitrates in any form (e.g. nitroglycerin, isosorbide, illicit "poppers") either regularly or intermittently, increases risk of severe or potentially fatal hypotension, hypersensitivity

WARNING: Hypotension may occur due to vasodilation

Sodium Bicarbonate

Name: Bicarb

Class: Antidote, other

Pharmacologic Action: Increases blood and urinary pH by releasing a bicarbonate ion, which in turn neutralizes hydrogen ion concentrations

Indications: For the management of cardiac arrest in cases in which either hyperkalemia or tricyclic antidepressant (TCA) overdose are suspected as contributory, QRS prolongation in known or suspected TCA overdose

Contraindications: Documented hypersensitivity, severe pulmonary edema, known alkalosis, hyponatremia, or hypocalcemia

Sodium Nitrite

Name: Nithiodote®

Class: Cyanide antidote

Pharmacologic Action: Nitrites create methemoglobins to bind to cyanide

Indications: For the management of cyanide toxicity

Contraindications: Documented hypersensitivity, suspected or confirmed smoke inhalation and/or carbon monoxide poisoning

WARNING: There is a risk of worsening hypoxia due to methemoglobin formation. In addition, sodium nitrite can cause serious adverse reactions and death from hypotension and methemoglobin formation. Monitor to ensure adequate perfusion and oxygenation during treatment with sodium nitrite.

Sodium Thiosulfate

Name: Nithiodote®

Class: Cyanide antidote

Pharmacologic Action: Thiosulfate is sulfur donor utilized by rhodenase to convert cyanide to less toxic thiocyanate

Indications: For the management of cyanide toxicity

Contraindications: Documented hypersensitivity

Sorbitol

Name: Sorbitol

Class: Laxatives, osmotic

Pharmacologic Action: Polyalcoholic sugar with hyperosmotic effects

Indications: Administered for the management of patients suffering from toxic ingestions

Contraindications: Acute abdominal pain, nausea, vomiting, or other symptoms of appendicitis or undiagnosed abdominal pain, documented hypersensitivity

WARNING: Sorbitol is no longer recommended to be given with activated charcoal

Tadalafil

Name: Cialis®, Adcirca®

Class: Pulmonary artery hypertension therapy, PDE-5 inhibitors; phosphodiesterase-5 enzyme inhibitor

Pharmacologic Action: Pulmonary arterial hypertension (PAH): inhibits PDE-5, increasing cyclic guanosine monophosphate (cGMP) to allow relaxation of pulmonary vascular smooth-muscle cells and vasodilation of pulmonary vasculature

Indications: As an adjunct to descent in the management of high altitude pulmonary edema (HAPE)

Contraindications: Concomitant use of any form of organic nitrates (e.g. nitroglycerin, isosorbide dinitrate, isosorbide mononitrate, illicit "poppers"), either regularly or intermittently; may potentiate hypotensive effect of nitrates. Hypersensitivity, including Stevens-Johnson syndrome and exfoliative dermatitis

WARNING: Hypotension may occur due to vasodilation

Ziprasidone

Name: Geodon®

Class: Second generation antipsychotic

Pharmacologic Action: Acts as antagonist at dopamine-2 and serotonin type 1 and 2 (5HT1D, 5HT2A) receptors; acts as agonist at serotonin 5HT1A receptor; moderately inhibits reuptake of norepinephrine and serotonin; has alpha-blocking and antihistaminic activity

Indications: For the management of agitated or violent patients suffering a behavioral emergency

Contraindications: Documented hypersensitivity, any drugs or conditions that prolong QT interval, recent acute myocardial infarction, uncompensated heart failure

IV. Approved Abbreviations

The following is the project's list of approved medical abbreviations used in this document. The drug.com article "[Medical Abbreviations on Pharmacy Prescriptions](#)" is considered the reference of authority.

Abbreviation	Description
ACS	Acute coronary syndrome
AED	Automatic external defibrillator
A-FIB	Atrial fibrillation
ALS	Advanced life support
AMS	Altered mental status
ASA	Aspirin
AV	Atrioventricular
AVPU	Neurological status measure: alert, verbal, pain, unresponsive
BiPAP	Bi-level positive airway pressure
BLS	Basic life support
BP	Blood pressure
BPM	Beats per minute
BSA	Body surface area
BSI	Body substance isolation
BVM	Bag-valve-mask
CABG	Coronary artery bypass graft
CAD	Coronary artery disease
CARES	Cardiac Arrest Registry to Enhance Survival
CC	Chief complaint
CDC	Centers for Disease Control and Prevention
CHF	Congestive heart failure
CNS	Central nervous system
CO	Carbon monoxide
CO ₂	Carbon dioxide
COPD	Chronic obstructive pulmonary disease
CP	Chest pain
CPAP	Continuous positive airway pressure
CPI	Continuous performance improvement
CPR	Cardiopulmonary resuscitation


Abbreviation	Description
C-SECTION	Caesarean section
C-SPINE	Cervical spine
CT	Cat scan, cardiac technician
CVA	Cerebrovascular accident (stroke)
D5W	5% dextrose in water
DKA	Diabetic ketoacidosis
DNI	Do not intubate
DNR	Do not resuscitate
DT	Delirium tremens
Dx	Diagnosis
ECPR	Extracorporeal cardiopulmonary resuscitation
EEG	Electroencephalogram
EENT	Eye, ear, nose, and throat
EGD	Extraglottic device
ECG	Electrocardiogram
EMS	Emergency medical services
EMT	Emergency medical technician
ePCR	Electronic patient call/care record/report
ET	Endotracheal
ETA	Estimated time of arrival
ETCO ₂	End-tidal CO ₂
ETOH	Ethanol (alcohol)
ETT	Endotracheal tube
FBAO	Foreign body airway obstruction
FiO ₂	Fraction of inspired oxygen
g	Gram(s)
GI	Gastrointestinal
gtts	Drops
GU	Gastrourinary
GYN	Gynecology (gynecological)
HFNC	High flow nasal cannula
HR	Heart rate (hour)
ICU	Intensive care unit
IM	Intramuscular
IO	Intraosseous

Abbreviation	Description
IPPB	Intermittent positive pressure breathing
IV	Intravenous
IVP	Intravenous push
J	Joules
JVD	Jugular vein distension
kg	Kilogram
KVO	Keep vein open
L	Liter
LMA	Laryngeal mask airway
LPM	Liters per minutes
LR	Lactated Ringer's
MAT	Multifocal atrial tachycardia
mcg	Microgram(s)
MED	Medicine
mg	Milligram(s)
mg/dL	Milligrams per deciliter
MI	Myocardial infarction (heart attack)
mL	Milliliter
mmHg	Millimeters of mercury
mmol	Millimole
MOLST	Medical orders for life-sustaining treatment
MS	Mental status
msec	Millisecond
MVC	Motor vehicle crash
N/V	Nausea/vomiting
NC	Nasal cannula
NRB	Non-rebreather
NS	Normal saline
NSR	Normal sinus rhythm
OB/GYN	Obstetrics/gynecology
O ₂	Oxygen
P	Pulse
PAC	Premature atrial contraction
PCR	Patient call/care record/report
PE	Pulmonary embolus

Abbreviation	Description
PEA	Pulseless electrical activity
PO	Orally
POLST	Physician orders for life-sustaining treatment
PPE	Personal protection equipment
prn	As needed
PVC	Premature ventricular contraction
q	Every (e.g. q 3–5 minutes)
RR	Respiratory rate
RSI	Rapid sequence intubation
Rx	Medicine
sat	Saturation
SBP	Systolic blood pressure
SC	Subcutaneous
SCBA	Self-contained breathing apparatus
SCUBA	Self-contained under-water breathing apparatus
SGD	Supraglottic device
SL	Sublingual
SOB	Shortness of breath
ST	Sinus tachycardia
SVT	Supraventricular tachycardia
T	Temperature
TBSA	Total body surface area
TCA	Tricyclic antidepressants
TIA	Transient ischemic attack
TID	Three times a day
TKO	To keep open
VF	Ventricular fibrillation
VS	Vital signs
VT	Ventricular tachycardia
yo	Years old (years of age)


v. Burn Body Surface Area

Burn Size Chart 1



Source: Used with permission, University of Utah Burn Center

Burn Size Chart 2



Source: American Heart Association, *Pediatric Advanced Life Support Textbook*, 2013

Percentage of Total Body Surface Area by Age, Anatomic Structure, and Body Habitus

<i>Adult</i>	
Anatomic Structure	Surface Area
Anterior head	4.5%
Posterior head	4.5%
Anterior torso	18%
Posterior torso	18%
Anterior leg, each	9%
Posterior leg, each	9%
Anterior arm, each	4.5%
Posterior arm, each	4.5%
Genitalia, perineum	1%

<i>Child</i>	
Anatomic Structure	Surface Area
Anterior head	9%
Posterior head	9%
Anterior torso	18%
Posterior torso	18%
Anterior leg, each	6.75%
Posterior leg, each	6.75%
Anterior arm, each	4.5%
Posterior arm, each	4.5%
Genitalia, perineum	1%

<i>Adult – Obese 80 kg</i>	
Anatomic Structure	Surface Area
Head and neck	2%
Anterior torso	25%
Posterior torso	25%
Leg, each	20%
Arm, each	5%
Genitalia, perineum	0%

<i>Infant 10 kg</i>	
Anatomic Structure	Surface Area
Head and neck	20%
Anterior torso	16%
Posterior torso	16%
Leg, each	16%
Arm, each	8%
Genitalia, perineum	1%

VI. Neurologic Status Assessment

Neurologic status assessment involves establishing a baseline and then trending any change in patient neurologic status. Glasgow Coma Score (GCS) is frequently used, but there are often errors in applying and calculating this score. With this in consideration, Glasgow Coma Score may not be more valid than a simpler field approach. Either AVPU (Alert, Verbal, Painful, Unresponsive—see below) or only the motor component of the GCS may more effectively serve in this capacity.

Glasgow Coma Score			
	Points	Pediatric	Adult
Eyes	1	No eye opening	No eye opening
	2	Eye opening to pain	Eye opening to pain
	3	Eye opening to verbal	Eye opening to verbal
	4	Eyes open spontaneously	Eyes open spontaneously
Verbal	1	No vocalization	No verbal response
	2	Inconsolable, agitated	Incomprehensible sounds
	3	Inconsistently consolable, moaning	Inappropriate words
	4	Cries but consolable, inappropriate interactions	Confused
	5	Smiles, oriented to sounds, follows objects, interacts	Oriented
Motor	1	No motor response	No motor response
	2	Extension to pain	Extension to pain
	3	Flexion to pain	Flexion to pain
	4	Withdraws from pain	Withdraws from pain
	5	Localizes pain	Localizes pain
	6	Obeys commands	Obeys commands

AVPU

A: The patient is alert

V: The patient responds to verbal stimulus

P: The patient responds to painful stimulus

U: The patient is completely unresponsive

VII. Abnormal Vital Signs

Age	Heart Rate	Resp Rate	Systolic BP	Temp (°C)
0 d – 1 m	> 205	> 60	< 60	<36 or >38
≥ 1 m - 3 m	> 205	> 60	< 70	<36 or >38
≥ 3 m - 1 r	> 190	> 60	< 70	<36 or >38.5
≥ 1 y - 2 y	> 190	> 40	< 70 + (age in yr × 2)	<36 or >38.5
≥ 2 y - 4 y	> 140	> 40	< 70 + (age in yr × 2)	<36 or >38.5
≥ 4 y - 6 y	> 140	> 34	< 70 + (age in yr × 2)	<36 or >38.5
≥ 6 y - 10 y	> 140	> 30	< 70 + (age in yr × 2)	<36 or >38.5
≥ 10 y - 13 y	> 100	> 30	< 90	<36 or >38.5
> 13 y	> 100	>16	< 90	<36 or >38.5

VIII. Evidence-Based Guidelines: GRADE Methodology

An Overview of GRADE Methodology

Although engagement in quality EMS research has increased significantly, the demand for evidence-based quality prehospital research continues to exceed its availability. The need for evidence-based prehospital patient care protocols was clearly recognized by the Institute of Medicine of the National Academies and clearly stated in 2007 in *The Future of Emergency Care: Emergency Medical Services at the Crossroads*.

The Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) methodology is a transparent process where the available research is reviewed and assessed by a panel of subject matter experts. Following this thorough review process, the available research is reviewed and graded for its validity based upon the assessment of the workgroup, and an evidence-based guideline (EBG) is developed based upon the outcome of the workgroup.

The Federal Interagency Committee on Emergency Medical Services (FICEMS) and the National EMS Advisory Council (NEMSAC) approved the National Prehospital Evidence-Based Guideline Model Process for the development, implementation, and evaluation of evidence-based guidelines. This model process recommends the use of the GRADE methodology for the guideline development tool. The six process steps of the GRADE EBG development tool are:


- Assemble the expert panel and provide GRADE training.
- Define the EBG content area and establish the specific clinical questions to address in patient, intervention, comparison, and outcome (PICO) format.
- Prioritize outcomes to facilitate systematic literature searches.
- Create GRADE tables (or evidence profiles) for each PICO question.
- Vet and endorse GRADE evidence tables and draft recommendations.
- Synthesize recommendations into an EMS protocol and visual algorithm.

The current evidence-based guidelines cited in this document were created for and released by NHTSA; however, the GRADE methodology is not proprietary to NHTSA or any other organization. Local, regional, and state EMS agencies and EMS systems are encouraged to support the ongoing need for quality prehospital care, improved patient outcome, and the growing demand for EBGs for EMS.

References:

Brown KM. The development of evidence-based prehospital guidelines using a GRADE-based methodology, *Prehospital Emergency Care*, 2014, Suppl 1:3-14, 2014

IX. 2011 Guidelines for Field Triage of Injured Patients



Source: Adapted from American College of Surgeons. Resources for the optimal care of the injured patient. Chicago, IL: American College of Surgeons; 2006. Footnotes (see following page) have been added to enhance understanding of field triage by persons outside the acute injury care field.

<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6101a1.htm>

* The upper limit of respiratory rate in infants is greater than 29 breaths per minute to maintain a higher level of overtriage for infants

- [†] Trauma centers are designated Level I–IV, with Level I representing the highest level of trauma care available.
- [§] Any injury noted in Steps Two and Three triggers a "yes" response.
- [¶] Age less than 15 years.
- **** Intrusion refers to interior compartment intrusion, as opposed to deformation which refers to exterior damage.
- ^{††} Includes pedestrians or bicyclists thrown or run over by a motor vehicle or those with estimated impact greater than 20 mph with a motor vehicle.
- ^{§§} Local or regional protocols should be used to determine the most appropriate level of trauma center; appropriate center need not be Level I.
- ^{¶¶} Age greater than 55 years.
- ***** Patients with both burns and concomitant trauma for whom the burn injury poses the greatest risk for morbidity and mortality should be transferred to a burn center. If the nonburn trauma presents a greater immediate risk, the patient may be stabilized in a trauma center and then transferred to a burn center.
- ^{†††} Injuries such as an open fracture or fracture with neurovascular compromise.
- ^{§§§} Emergency medical services.
- ^{¶¶¶} Patients who do not meet any of the triage criteria in Steps One through Four should be transported to the most appropriate medical facility as outlined in local EMS protocols.