





## **Tick Surveillance Report**

Epidemiological Week: 47

Collection Date: 11/17/2025

Division of Public Health Bureau of Communicable Disease Vectorborne Disease Unit



# (j) About

The Wisconsin Department of Health Services conducts **field collection of ticks** routinely during the tick season and this report summarizes the ticks collected. Sites in three counties serve as sentinel sites and tick collection occurs approximately every other week throughout the tick season, typically March through November. Field collection activities highlighted in this report focus on **blacklegged ticks (also known as deer ticks and** *Ixodes scapularis***)** because this species of ticks is the most important vector of human tickborne infections in Wisconsin.

Field collection and tick images submitted through the **DHS Tick Identification Service** are one of several indicators DHS uses to monitor tick activity and trends. This data may not reflect the threat of encountering a tick in all areas of the state. Ticks are most active in Wisconsin from spring through fall, but you should use caution throughout the year and <u>take steps to prevent tick bites</u>.

## Field Collection Activity Tracker

Overall blacklegged tick activity is based on the average number of each tick life stage found while conducting tick surveillance. Usually, only nymphs and adult female ticks can spread illness.

	Adults	Nymphs	Larvae	
Life stage	J.	No.		
<b>Current Activity</b>	Moderate	Minimal	Minimal	
Trajectory	<b>Decreasing</b>	Stable	Stable	

Moderate

High


Very High

Low


Minimal

Activity Scale





DHS conducts bi-weekly sampling of host-seeking ticks at three sentinel sites in Wisconsin. During each visit, a white denim tick drag is pulled through vegetation along a 750 meter transect and drags are check for ticks at 15 meter intervals. All host-seeking ticks found on drags are collected and stored in 70% ethanol.



Data shows total number of each life stage of the blacklegged tick found during each visit. Adult blacklegged ticks are displayed as **Female** and **Male**.

		Other ticks			
County	Larva	Nymph	Adult Female	Adult Male	
Iowa	0	0	2	3	0
Waupaca	0	0	3	9	0
Lincoln	0	0	2	0	0
Average	0	0	2.33	4	0

**Notes:** 




## Trends by Sentinel Site County, 2021-2025


Total number of blacklegged ticks, collected on 750 meters during each visit to three sentinel surveillance sites in odd–numbered CDC epidemiological weeks (Epi week 9, 11, 13, 15, 17...).




#### Total number of adult blacklegged ticks by epi week



#### Total number of nymphal blacklegged ticks by epi week



### Total number of larval blacklegged ticks by epi week



The graph shows the total number of submissions to the Wisconsin Department of Health Services Tick Identification Service since Epidemiological Week 1 (January 1-4). Data collected through the Tick Identification Service is shown below for submissions from users in Wisconsin only.

