



INFANT FORMULAS: Contract Standard

Provided by the Wisconsin WIC Program to Partially Breastfed and Fully Formula Fed Infants (Birth to 12 Months of Age)

Formula	Use	Protein Sources	CHO Sources	Fat Sources	Product Form, Size, Amount	By Age, Maximum Number Cans Per Month and Prepared Ounces Per Day ¹					
						0-3 Mos		4-5 Mos		6-11 Mos ²	
						cans	oz	cans	oz	cans	oz
Similac Advance Abbott Nutrition	Use: Nutritionally complete cow's milk-based, iron fortified infant formula for routine feeding in infants with normal GI tracts. Contains OptiGRO™, an exclusive blend of DHA, lutein and vitamin E. Kosher and Halal. Standard Dilution: 20 kcal/fl oz , 1 scoop to 2 oz. water	(8% kcals) Intact cow's milk protein	(42% kcals) Lactose, Prebiotic: galactooligosaccharides	(50% kcals) High oleic safflower, soy, and coconut oils (0.15% DHA; 0.40% ARA) Oil ratio: 40:30:29	Powder 12.4 oz. can 90 fl oz	9	27	10	30	7	21
					Concentrate 13 fl oz. carton 26 fl oz.	31	26	34	29	24	20
					Ready to Feed 32 fl oz.	26	27	28	29	20	21
Similac Soy Isomil Abbott Nutrition	Use: Nutritionally complete soy-based infant formula for infants with feeding problems such as fussiness and gas; for infants whose parents choose a milk-free formula as a first feeding or as a supplement to breastfeeding; and for infants of vegetarian families. A lactose-free feeding for infants with disorders for which lactose should be avoided, including lactase deficiency, lactose intolerance, and galactosemia. Contains OptiGRO™, an exclusive blend of DHA, lutein and vitamin E. Kosher and Halal. Standard Dilution: 20 kcal/fl oz, 1 scoop to 2 oz. water	(10% kcals) Soy protein isolate, L-methionine	(41% kcals) Corn syrup, sugar, (Ratio: 80:20) Prebiotic: fructooligosaccharides	(49% kcals) High oleic safflower, soy, and coconut oils (0.15% DHA, 0.40% ARA) Oil ratio: 40:30:29	Powder 12.4 oz. can 90 fl. oz.	9	27	10	30	7	21
					Concentrate 13 fl. oz. carton 26 fl. oz.	31	26	34	29	24	20
					Ready to Feed 32 fl. oz.	26	27	28	29	20	21
Similac Total Comfort Abbott Nutrition	Use: Nutritionally complete infant formula with partially hydrolyzed whey protein as an alternative to standard intact protein, milk-based formulas. For mild tolerance symptoms such as fussiness and gas due to lactose sensitivity. Contains OptiGRO™, an exclusive blend of DHA, Lutein, and Vitamin E. Kosher and Halal. Standard Dilution: 20 kcal/fl oz., 1 scoop to 2 oz. water	(9% kcals) Whey protein hydrolysate	(42% kcals) Maltodextrin, sugar, (Ratio: 80:20) Prebiotic: galactooligosaccharides	(49% kcals) High oleic safflower, soy and coconut oils (0.15% DHA, 0.40% ARA) Oil ratios: 40:30:29	Powder 12.6 oz. can 90 fl. oz.	9	27	10	30	7	21
Similac Sensitive Abbott Nutrition	Use: Nutritionally complete infant feeding that is an alternative to standard milk-based formulas. For mild tolerance symptoms such as fussiness and gas due to lactose sensitivity. Suitable for infants with lactose	(9% kcals) Milk protein isolate	(42% kcals) Maltodextrin, sugar(Ratio:80:20)	(49% kcals) High oleic safflower, soy, and	Powder 12.5 oz. can 90 fl. oz	9	27	10	30	7	21

Formula	Use	Protein Sources	CHO Sources	Fat Sources	Product Form, Size, Amount	By Age, Maximum Number Cans Per Month and Prepared Ounces Per Day ¹					
						0-3 Mos		4-5 Mos		6-11 Mos ²	
						cans	oz	cans	oz	cans	oz
	sensitivity. Contain OptiGRO™, an exclusive blend of DHA, lutein, and vitamin E. Kosher and Halal. Standard Dilution: 20 kcal/fl oz., 1 scoop to 2 oz. water		Prebiotic: galactooligosaccharides	coconut oils (0.15% DHA, 0.40% ARA) Oil ratio: 40:30:29	Ready to Feed 32 fl oz.	26	27	28	29	20	21

¹ Partially breastfed babies receive less formula; the amount is adjusted to help mothers maintain an adequate milk supply.

² Infants 6-11 months of age may receive the same monthly maximum as infants 4 through 5 months of age when infant foods are contraindicated for medical reasons.

³ Infant cases with serious infections secondary to *Enterobacter sakazakii* have been reported in the literature that were traced back to powdered infant formulas contaminated with this organism. Preterm infants and those with underlying medical conditions may be at higher risk of developing infection. FDA recommends sterile, commercial Ready-To-Feed formulas (RTF) for immunocompromised infants including preterm infants. WIC routinely provides cans of powdered formula. RTF standard formula may be provided when the local WIC Project RD verifies that the water supply is unsafe to drink, if the caregiver has difficulty mixing powder or concentrate properly, if RTF better accommodates the participant's condition (e.g., immunocompromised), or if RTF improves the participant's compliance in consuming the prescribed formula.

Go to: www.dhs.wisconsin.gov/wic/nonDiscrimination.htm for the U.S. Department of Agriculture non-discrimination statement.